


Institute for Language, Logic and Information

**SUBALGEBRAS OF DIAGONIZABLE ALGEBRAS
OF THEORIES CONTAINING ARITHMETIC**

V.Yu. Shavrukov

ITLI Prepublication Series


X-91-03



University of Amsterdam

The ITLI Prepublication Series

- 1986
 86-01 The Institute of Language, Logic and Information
 86-02 Peter van Emde Boas A Semantical Model for Integration and Modularization of Rules
 86-03 Johan van Benthem Categorical Grammar and Lambda Calculus
 86-04 Reinhard Muskens A Relational Formulation of the Theory of Types
 86-05 Kenneth A. Bowen, Dick de Jongh Some Complete Logics for Branched Time, Part I Well-founded Time,
 86-06 Johan van Benthem Logical Syntax Forward looking Operators
 1987
 87-01 Jeroen Groenendijk, Martin Stokhof Type shifting Rules and the Semantics of Interrogatives
 87-02 Renate Bartsch Frame Representations and Discourse Representations
 87-03 Jan Willem Klop, Roel de Vrijer Unique Normal Forms for Lambda Calculus with Surjective Pairing
 87-04 Johan van Benthem Polyadic quantifiers
 87-05 Víctor Sánchez Valencia Traditional Logicians and de Morgan's Example
 87-06 Eleonore Oversteegen Temporal Adverbials in the Two Track Theory of Time
 87-07 Johan van Benthem Categorical Grammar and Type Theory
 87-08 Renate Bartsch The Construction of Properties under Perspectives
 87-09 Herman Hendriks Type Change in Semantics: The Scope of Quantification and Coordination
 1988
 LP-88-01 Michiel van Lambalgen *Logic, Semantics and Philosophy of Language:* Algorithmic Information Theory
 LP-88-02 Yde Venema Expressiveness and Completeness of an Interval Tense Logic
 LP-88-03 Year Report 1987
 LP-88-04 Reinhard Muskens Going partial in Montague Grammar
 LP-88-05 Johan van Benthem Logical Constants across Varying Types
 LP-88-06 Johan van Benthem Semantic Parallels in Natural Language and Computation
 LP-88-07 Renate Bartsch Tenses, Aspects, and their Scopes in Discourse
 LP-88-08 Jeroen Groenendijk, Martin Stokhof Context and Information in Dynamic Semantics
 LP-88-09 Theo M.V. Janssen A mathematical model for the CAT framework of Eurotra
 LP-88-10 Anneke Kleppe A Blissymbolics Translation Program
 ML-88-01 Jaap van Oosten *Mathematical Logic and Foundations:* Lifschitz' Realizability
 ML-88-02 M.D.G. Swaen The Arithmetical Fragment of Martin Löf's Type Theories with weak Σ -elimination
 ML-88-03 Dick de Jongh, Frank Veltman Provability Logics for Relative Interpretability
 ML-88-04 A.S. Troelstra On the Early History of Intuitionistic Logic
 ML-88-05 A.S. Troelstra Remarks on Intuitionism and the Philosophy of Mathematics
 CT-88-01 Ming Li, Paul M.B. Vitanyi *Computation and Complexity Theory:* Two Decades of Applied Kolmogorov Complexity
 CT-88-02 Michiel H.M. Smid General Lower Bounds for the Partitioning of Range Trees
 CT-88-03 Michiel H.M. Smid, Mark H. Overmars Maintaining Multiple Representations of
 Leen Torenvliet, Peter van Emde Boas Dynamic Data Structures
 CT-88-04 Dick de Jongh, Lex Hendriks Computations in Fragments of Intuitionistic Propositional Logic
 Gerard R. Renardel de Lavalette
 CT-88-05 Peter van Emde Boas Machine Models and Simulations (revised version)
 CT-88-06 Michiel H.M. Smid A Data Structure for the Union-find Problem having good Single-Operation Complexity
 CT-88-07 Johan van Benthem Time, Logic and Computation
 CT-88-08 Michiel H.M. Smid, Mark H. Overmars Multiple Representations of Dynamic Data Structures
 Leen Torenvliet, Peter van Emde Boas
 CT-88-09 Theo M.V. Janssen Towards a Universal Parsing Algorithm for Functional Grammar
 CT-88-10 Edith Spaan, Leen Torenvliet, Peter van Emde Boas Nondeterminism, Fairness and a Fundamental Analogy
 CT-88-11 Sieger van Dennecheuvel, Peter van Emde Boas Towards implementing RL
 X-88-01 Marc Jumelet *Other prepublications:* On Solovay's Completeness Theorem
 1989
 LP-89-01 Johan van Benthem *Logic, Semantics and Philosophy of Language:* The Fine-Structure of Categorical Semantics
 LP-89-02 Jeroen Groenendijk, Martin Stokhof Dynamic Predicate Logic, towards a compositional,
 non-representational semantics of discourse
 LP-89-03 Yde Venema Two-dimensional Modal Logics for Relation Algebras and Temporal Logic of Intervals
 LP-89-04 Johan van Benthem Language in Action
 LP-89-05 Johan van Benthem Modal Logic as a Theory of Information
 LP-89-06 Andreja Prijatelj Intensional Lambek Calculi: Theory and Application
 LP-89-07 Heinrich Wansing The Adequacy Problem for Sequential Propositional Logic
 LP-89-08 Víctor Sánchez Valencia Peirce's Propositional Logic: From Algebra to Graphs
 LP-89-09 Zhisheng Huang Dependency of Belief in Distributed Systems
 ML-89-01 Dick de Jongh, Albert Visser *Mathematical Logic and Foundations:* Explicit Fixed Points for Interpretability Logic
 ML-89-02 Roel de Vrijer Extending the Lambda Calculus with Surjective Pairing is conservative
 ML-89-03 Dick de Jongh, Franco Montagna Rosser Orderings and Free Variables
 ML-89-04 Dick de Jongh, Marc Jumelet, Franco Montagna On the Proof of Solovay's Theorem
 ML-89-05 Rineke Verbrugge Σ -completeness and Bounded Arithmetic
 ML-89-06 Michiel van Lambalgen The Axiomatization of Randomness
 ML-89-07 Dirk Roorda Elementary Inductive Definitions in HA: from Strictly Positive towards Monotone
 ML-89-08 Dirk Roorda Investigations into Classical Linear Logic
 ML-89-09 Alessandra Carbone Provable Fixed points in $\text{IA}_0 + \Omega_1$
 CT-89-01 Michiel H.M. Smid *Computation and Complexity Theory:* Dynamic Deferred Data Structures
 CT-89-02 Peter van Emde Boas Machine Models and Simulations
 CT-89-03 Ming Li, Herman Neuféglise, Leen Torenvliet, Peter van Emde Boas On Space Efficient Simulations
 CT-89-04 Harry Buhrman, Leen Torenvliet A Comparison of Reductions on Nondeterministic Space
 CT-89-05 Pieter H. Hartel, Michiel H.M. Smid A Parallel Functional Implementation of Range Queries
 Leen Torenvliet, Willem G. Vree
 CT-89-06 H.W. Lenstra, Jr. Finding Isomorphisms between Finite Fields
 CT-89-07 Ming Li, Paul M.B. Vitanyi A Theory of Learning Simple Concepts under Simple Distributions and
 Average Case Complexity for the Universal Distribution (Prel. Version)
 CT-89-08 Harry Buhrman, Steven Homer Honest Reductions, Completeness and
 Leen Torenvliet Nondeterministic Complexity Classes
 CT-89-09 Harry Buhrman, Edith Spaan, Leen Torenvliet On Adaptive Resource Bounded Computations
 CT-89-10 Sieger van Dennecheuvel The Rule Language RL/1
 CT-89-11 Zhisheng Huang, Sieger van Dennecheuvel Towards Functional Classification of Recursive Query Processing
 Peter van Emde Boas
 X-89-01 Marianne Kalsbeek *Other Prepublications:* An Orey Sentence for Predicative Arithmetic
 X-89-02 G. Wagemakers New Foundations: a Survey of Quine's Set Theory
 X-89-03 A.S. Troelstra Index of the Heyting Nachlass
 X-89-04 Jeroen Groenendijk, Martin Stokhof Dynamic Montague Grammar, a first sketch
 X-89-05 Maarten de Rijke The Modal Theory of Inequality
 X-89-06 Peter van Emde Boas Een Relationele Semantiek voor Conceptueel Modelleren: Het RL-project
 1990 SEE INSIDE BACK COVER



Instituut voor Taal, Logica en Informatie
Institute for Language, Logic and
Information

Faculteit der Wiskunde en Informatica
(Department of Mathematics and Computer Science)
Plantage Muidergracht 24
1018TV Amsterdam

Faculteit der Wijsbegeerte
(Department of Philosophy)
Nieuwe Doelenstraat 15
1012CP Amsterdam

SUBALGEBRAS OF DIAGONIZABLE ALGEBRAS
OF THEORIES CONTAINING ARITHMETIC

V.Yu. Shavrukov
Steklov Mathematical Institute
Academy of Sciences of the USSR
Vavilova Str. 42, 117966 Moscow GSP-1, USSR

Department of Mechanics and Mathematics, Moscow State University

Contents

§ 0. Introduction	1
§ 1. Preliminaries	3
§ 2. On conservativity in L	13
§ 3. A family of Kripke models	24
§ 4. Finite credibility extent	30
§ 5. The strong disjunction property and apt formulae	44
§ 6. Σ_1 -ill theories of infinite credibility extent	57
§ 7. Σ_1 -sound theories	72
§ 8. An application	81
§ 9. A question of arithmetical complexity	84
§ 10. Arbitrary subalgebras. Σ_1 -ill theories	93
§ 11. Arbitrary subalgebras. Σ_1 -sound theories	104
References	112

§ 0. Introduction

The study of the diagonalizable algebras of theories containing arithmetic was initiated in the early seventies by Macintyre & Simmons [28] and Magari [29]. In 1976 Solovay [47] characterized the equational theory of the diagonalizable algebra of Peano arithmetic. This theory was shown to be adequately described by the well-known modal logic L . Later on, Montagna [31], Artemov [2], Visser [48] and Boolos [12] strengthened this result somewhat by independently demonstrating that the free diagonalizable algebra on countably many generators is (isomorphic to) a subalgebra of the diagonalizable algebra of Peano arithmetic. The equational theories of diagonalizable algebras of other strong enough theories were calculated by Visser [49] (cf. also Artemov [3]). These are given by the series L , $L+\Box 1$, $L+\Box\Box 1$, ... Among the recent investigations of the subject we may also mention Montagna's paper [32] which undertakes a systematic inquiry into generalizations of Goldfarb's Principle.

Nonetheless, the information on diagonalizable algebras of theories currently available is dejectingly scarce and therefore leaves ample scope to further research. Thus, for instance, it would be natural to attempt a closer look at subalgebras of these algebras. This is the theme of the present paper. It is predominantly devoted to the question which diagonalizable algebras can be embedded into the diagonalizable algebra of a theory. For the easier case of embeddings with r.e. range we obtain a complete solution. It turns out that a short list of most obvious restrictions constitutes a characterization of r.e. subalgebras of the diagonalizable algebra of a theory. Partial results in this (or at least in a closely parallel) direction were obtained by Jumelet [27]. In fact, the work of Jumelet was my main source of ideas and inspiration.

The plan of the paper is as follows. § 1 recollects the necessary definitions and earlier results. It also contains a result on the length of proofs which, in view of a construction in § 11, does not look absolutely out of place here. In §§ 2, 3 and

5 we carry out some modal-logical work relevant for subsequent progress. As a by-product of this we obtain a uniform version of the Craig Interpolation Lemma for L . The main result of the paper is to be found in §§ 4, 6 and 7 where r.e. subalgebras of diagonalizable algebras of a wide class of theories are characterized. This takes us three §§ because we use three slightly different approaches to handle particular kinds of theories. Here we employ extensions of techniques developed by Solovay [47], Artemov [2], Boolos [12], Jumelet [27] and Beklemishev [5]. §§ 8 - 11 are of marginal interest. In § 8 we apply the result of § 7 to give an alternative proof to a lemma in Simmons [41]. Unfortunately the application will not require the full strength of our methods. A question concerning the arithmetical complexity of sentences needed to model a diagonalizable algebra in arithmetic is treated in § 9. In the last two §§ we find out whether our characterization of subalgebras of diagonalizable algebras of theories extends from r.e. to arbitrary subalgebras. It is shown in § 10 that for the case of Σ_1 -ill theories an easy generalization is possible. As regards Σ_1 -sound theories, the situation appears to be more complex and an example is given in § 11 that partially justifies our failure to describe subalgebras of diagonalizable algebras of these theories.

I would like to thank Lev Beklemishev for numerous stimulating ideas and invaluable comments. Without his help the present paper could have hardly been written. In particular, Lev Beklemishev brought my attention to a neat trick in Beklemishev [5] which a key idea for the argument in § 6 was derived from. Thanks are also due to Professors Sergei Artemov and Aleksandr Chagrov, Marc Jumelet and Andrei Muchnik for interesting discussions.

We assume that the reader is familiar with Smoryński [46] or at least with Solovay [47]. Knowledge of (rudiments of) diagonalizable algebra theory and modal logic, especially of L , should also be very helpful. For these matters, good references are Magari [29], Bernardi [8] and Bellissima [7].

§ 1. Preliminaries

1.A. Arithmetic

We shall study r.e. consistent theories whose language comprises that of primitive recursive arithmetic. Given a set Γ of arithmetic formulae, $\Delta_0(\Gamma)$ denotes the closure of Γ under Boolean combinations and primitive recursively bounded quantification. Let

$$\Sigma_0 = \Pi_0 = \Delta_0(\text{atomic arithmetic formulae})$$

and define Σ_{n+1} to be the closure of Π_n under lattice combinations and existential quantification. The class Π_n is defined analogously. We shall say that a formula φ is Σ_n over a theory T if there exists a Σ_n formula which T proves φ equivalent to. Finally, φ is Δ_n over T if it is both Σ_n and Π_n over T .

For Γ a set of arithmetic formulae, a theory T is said to be Γ -sound if each theorem of T which is in Γ is true. A theory is Γ -ill if it is not Γ -sound.

In compliance with a recent tradition of not involving much more arithmetic than is actually needed we take $I\Sigma_1$ as our base theory. In other words it is assumed throughout the paper that every theory under study contains induction for Σ_1 formulae as well as the basic axioms P^- (cf. Paris & Kirby [36]) and defining equations for primitive recursive function symbols. Note that our theory $I\Sigma_1$ proves the same theorems as the theory PRA of Smoryński [46]. The theory $I\Sigma_1$ of Paris & Kirby formulated in the language $(\leq, 0, S, +, \cdot)$ is very much the same as ours. That is not just to say that our variant of $I\Sigma_1$ is conservative over that of Paris & Kirby. What is more, every formula of the language of primitive recursive arithmetic translates easily and $I\Sigma_1$ -equivalently into the smaller language.

The following facts about $I\Sigma_1$ are well worth being kept in mind: The provably recursive functions of $I\Sigma_1$ are exactly the primitive recursive ones (Mints [30]); $I\Sigma_1$ proves induction (and therefore the least number principle) for $\Delta_0(\Sigma_1)$ formulae and each $\Delta_0(\Sigma_1)$ formula is Δ_2 over $I\Sigma_1$ (Hájek & Kucera [24]); every

$\Delta_0(\Sigma_n)$ sentence is equivalent to a Boolean combination of Σ_n sentences.

We assume that every theory comes equipped with a primitive recursive way α to recognize its axioms with which we associate a Δ_0 formula $\text{Prf}_\alpha(y,x)$, the *proof predicate (of T)*, to express that y is a (say, Hilbert-style) proof of x from the (extralogical) axioms given by α (cf. e.g. Feferman [16]). $\text{Pr}_\alpha(x)$, the *provability predicate (of T)*, is short for $\exists y \text{Prf}_\alpha(y,x)$. In what follows we shall be omitting the subscript α since no confusion is likely.

Each formula and, in general, each syntactical object is identified with its gödelnumber. The *numeral for n* , i.e. (the gödelnumber of) a zero followed by n strokes is denoted by \bar{n} . Finally, if $\varphi(x_1, \dots, x_m)$ is a formula then $\overline{\varphi(x_1, \dots, x_m)}$ is the primitive recursive term honestly representing the function which sends (n_1, \dots, n_m) to the numeral for $\varphi(\bar{n}_1, \dots, \bar{n}_m)$.

The least $n \in \omega$ s.t. T proves $\text{Pr}^n(\bar{1})$ is called the *credibility extent of T*. (We let $\text{Pr}^1(\bar{1}) \equiv \text{Pr}(\bar{1})$ and $\text{Pr}^{n+1}(\bar{1}) \equiv \overline{\text{Pr}(\text{Pr}^n(\bar{1}))}$). If no such $n \in \omega$ exists then T is said to be of *infinite credibility extent*. Note that if T is Σ_1 -sound then clearly its credibility extent is infinite. On the other hand, the credibility extent of a Σ_1 -ill theory does not only depend on the set of theorems of T, but also on the primitive recursive way α which the axioms of T are presented. Thus, Beklemishev [5] shows that if a Σ_1 -ill theory T contains full induction then a particular choice of α can make the credibility extent of T anything from 1 to ∞ .

1.B. The modal logic L

The modal logic L (whose other names are K4W (Seegerberg [40]), G (Solovay [47]), GL (Artemov [3]) and PRL (Smoryński [46])) was presumably first introduced by Smiley [42] whose motive for doing so was investigation of ethics rather than of provability in formal systems. The language of L consists of an infinite stock of propositional letters p_0, p_1, \dots , the usual propositional connectives and a unary modal operator \Box . In addi-

tion to the axioms and rules of the classical propositional logic, L contains the following axiom schemata:

$$\begin{aligned} \Box(A \rightarrow B) &\rightarrow \Box A \rightarrow \Box B \\ \Box A &\rightarrow \Box \Box A \\ \Box(\Box A \rightarrow A) &\rightarrow \Box A \end{aligned}$$

and the *necessitation rule*: from A infer $\Box A$.

For A a modal formula we write $\Diamond A$ short for $\neg \Box \neg A$ and $\Box^+ A$ short for $A \wedge \Box A$.

We write $\vdash_L A$ to mean that the formula A is derivable in L . $\vdash_L \Box^+ A \rightarrow \Box^+ B$ will usually be abbreviated as $A \vdash_L B$. Note that since $\vdash_L A$ is equivalent to $\vdash_L \Box^+ A$ (cf. Magari [29]), our notation is coherent in that $\vdash_L A$ if and only if $\top \vdash_L A$. Trivially,

$$\begin{aligned} \vdash_L A \rightarrow B &\text{ implies } A \vdash_L B ; \\ A &\vdash_L A ; \\ A \vdash_L B \text{ and } B &\vdash_L C \text{ imply } A \vdash_L C ; \\ A \vdash_L B &\text{ implies } A \vdash_L \Box B \text{ and } \Box A \vdash_L \Box B \text{ etc.} \end{aligned}$$

\vec{p}, \vec{q} etc. will be treated as variables ranging over finite (possibly empty) tuples of propositional letters.

Kripke semantics has long been known as a mighty weapon in the study of modal logic. We describe a variant of it suited for our purposes. A triple $\mathcal{K} = (K, R, *)$ is a (Kripke) \vec{p} -model if K , the *domain of \mathcal{K}* , is a non-empty set (of nodes); R , the *accessibility relation*, is a strict partial order on K s.t. R^{-1} is well founded and $*$ is a *forcing relation* between elements of K and those modal formulae all of whose propositional letters are among those in \vec{p} . $*$ should satisfy the usual commutativity conditions for Boolean connectives and for each $a \in K$ and each modal formula $A(\vec{p})$ one has $a * \Box A(\vec{p})$ if and only if $b * A(\vec{p})$ for all $b \in K$ s.t. $a R b$. We write $\mathcal{K} \models A$ (A holds in \mathcal{K}) if $a * A$, all $a \in K$.

By a *model* we mean a \vec{p} -model for some tuple \vec{p} . A model $\mathcal{K} = (K, R, *)$ is *finite* if so is K . \mathcal{K} is *rooted* if there exists a node $b \in K$ satisfying $b R a$, all $a \in K$ s.t. $a \neq b$. This b

is then called the *root* of K . A rooted model K is *treelike* if R is a tree on K . For K a rooted model, we write $K * A$ (K forces A ; A is forced in K ; K is a model of A) if the root of K forces A . Clearly $K \models A$ if and only if $K * \Box^+ A$.

It is well known that if a formula A is derivable in L then it holds in every model provided that the forcing relation is defined on A . Various specializations of the converse are also true. Thus, if a formula is forced in every finite rooted model, or even in every finite treelike model, then it is derivable in L (see e.g. Segerberg [40] or Solovay [47]; we shall be referring to this fact as the *Completeness Theorem for L*). The decidability of L follows (cf. also Bernardi [8]).

1.C. Diagonalizable algebras

A *diagonalizable algebra* (Magari [29]) is a pair (\mathfrak{A}, \Box) where \mathfrak{A} is a Boolean algebra with the usual operations $\wedge, \vee, \neg, \rightarrow, \top$ and \perp endowed with an operator \Box satisfying the following identities:

$$\begin{aligned} \Box(x \rightarrow y) &\rightarrow \Box x \rightarrow \Box y = \\ &\Box x \rightarrow \Box \Box x = \\ \Box(\Box x \rightarrow x) &\rightarrow \Box x = \\ &\Box \top = \top \end{aligned}$$

The confusion between modal-logical and algebraic notation is meant to stress the fact that a diagonalizable equation is an identity of the variety of diagonalizable algebras if and only if the corresponding modal formula is derivable in L (see Montagna [31]).

A Boolean filter f of a diagonalizable algebra \mathfrak{D} is a τ -filter if $x \in f$ implies $\Box x \in f$ for each element x of \mathfrak{D} . If a filter f is τ - then there exists the quotient algebra \mathfrak{D}/f . Conversely, the elements that are sent to τ by a homomorphism of diagonalizable algebras constitute a τ -filter (cf. Magari [29] or Bernardi [8]). For each subset X of a diagonalizable algebra \mathfrak{D} there exists the smallest τ -filter $t(X)$ containing X . Thus we can define $\mathfrak{D}/_X$, the *quotient (algebra) of \mathfrak{D} modulo X* , to be

$\mathfrak{D}/\tau(x)$.

Whenever we shall need to construct a particular example of diagonalizable algebra we shall produce an algebra of the form \mathbf{F}/\mathcal{E} where \mathbf{F} is the free diagonalizable algebra on an appropriate set of generators $\{p_i\}_{i \in I}$ (this algebra may be identified with the set of modal formulas using the generators as propositional letters modulo L-provable equivalence) and \mathcal{E} is a set of elements of \mathbf{F} , that is, of formulas in $\{p_i\}_{i \in I}$. Note that for a formula A in $\{p_i\}_{i \in I}$ one has $A = \tau$ in \mathbf{F}/\mathcal{E} if and only if there exists a finite subset \mathcal{F} of \mathcal{E} s.t. $\bigwedge \mathcal{F} \vdash_{\mathbf{L}} A$.

The *height* of a diagonalizable algebra \mathfrak{D} is defined as the least $n \in \omega$ s.t. $\Box^n \perp = \tau$. If for all $n \in \omega$ one has $\Box^n \perp \neq \tau$ then the height of \mathfrak{D} is *infinite*. \mathfrak{D} is ω -consistent if $\perp \neq \tau$ and $x = \tau$ whenever $\Box x = \tau$ for each element x of \mathfrak{D} . ω -consistency obviously implies infinite height. If $\Box x \vee \Box y = \tau$ implies $\Box x = \tau$ or $\Box y = \tau$ then \mathfrak{D} is said to possess the *disjunction property*. Clearly the height, ω -consistency and the disjunction property are inherited by subalgebras. One can show that among homomorphic images of a diagonalizable algebra of infinite height there always are diagonalizable algebras with the strong disjunction property.

A \perp -generated diagonalizable algebra is determined by its height up to isomorphism. Note that the disjunction property is shared by all the \perp -generated diagonalizable algebras whereas the only ω -consistent \perp -generated diagonalizable algebra is the free \perp -generated diagonalizable algebra.

A mapping $\nu: \omega \rightarrow \mathfrak{D}$ s.t. $\text{rng } \nu$ generates the (denumerable) diagonalizable algebra \mathfrak{D} is called a *numeration* of \mathfrak{D} . A numeration ν is *positive* if the set of diagonalizable polynomials $A(\vec{p})$ satisfying $A(\nu 0, \nu 1, \dots) = \tau$ is r.e. A numeration ν is *locally positive* if for each $n \in \omega$ the set of diagonalizable polynomials $A(p_0, \dots, p_n)$ satisfying $A(\nu 0, \dots, \nu n) = \tau$ is r.e. An algebra \mathfrak{D} is (locally) *positive* if a (locally) positive numeration of it exists. Clearly \mathfrak{D} is locally positive if and only if each of its finitely generated subalgebras is positive; a finitely generated diagonalizable algebra is positive if and only if

it is locally positive.

1.D. Diagonalizable algebras and arithmetic

The example of a diagonalizable algebra which motivates the definition is constructed from a theory T of the kind described in 1.A. The Boolean algebra \mathfrak{A} is taken to be the Lindenbaum Sentence Algebra of T , i.e. the set of sentences of T modulo T -provable equivalence and for the mapping \square one takes the provability predicate of T , that is, for φ a sentence, $\square\varphi \equiv \text{Pr}(\overline{\varphi})$. The well-known properties of $\text{Pr}(\cdot)$ guarantee that the algebra obtained in this way is a diagonalizable algebra. (In particular, the identity $\square(\square x \rightarrow x) \rightarrow \square x = \top$ disguises a formalized version of Löb's Theorem.) This diagonalizable algebra is called the *diagonalizable algebra of T* and is denoted by \mathfrak{D}_T . The concept was originally introduced by Macintyre & Simmons [28] without a name. The name was supplied later by Magari [29].

If Γ is a set of arithmetic sentences closed under Boolean operations and \square then \mathfrak{D}_T^Γ is the corresponding subalgebra of \mathfrak{D}_T . A subalgebra of \mathfrak{D}_T is *r.e.* if the underlying set of sentences is. The usual gödelnumbering of sentences gives rise to a positive numeration of each r.e. subalgebra of \mathfrak{D}_T including \mathfrak{D}_T itself.

Clearly the height of \mathfrak{D}_T is equal to the credibility extent of T .

In diagonalizable algebras (and even in diagonalizable algebras of infinite height) neither of ω -consistency and the disjunction property implies the other. The situation in diagonalizable algebras of theories is different. In fact, the following are equivalent:

- (i) T is Σ_1 -sound;
- (ii) $T \vdash \sigma \vee \tau$ implies $T \vdash \sigma$ or $T \vdash \tau$ for each pair of Σ_1 sentences σ and τ ;
- (iii) T decides every sentence which is Δ_1 over T ;
- (iv) \mathfrak{D}_T is ω -consistent;
- (v) The credibility extent of T is greater than 1 and \mathfrak{D}_T possesses the disjunction property.

(i) \Leftrightarrow (ii) \Leftrightarrow (iii) is proved in Jensen & Ehrenfeucht [26] and Guaspari [23] (cf. also Friedman [19] and Smoryński [45] and [46]). The remaining equivalences are folklore and are typical applications of *Goldfarb's Principle*:

Let σ be a Σ_1 sentence and let $T \vdash \text{Pr}(\bar{1}) \rightarrow \sigma$. Then there exists a sentence τ (which can be chosen Σ_1 or Π_1) s.t. $T \vdash \sigma \leftrightarrow \text{Pr}(\bar{\tau})$.

(cf. Visser [49], Bernardi & Mirolli [9], Montagna [32] or Montagna & Sommaruga [34]).

As evidenced by (iv) \Leftrightarrow (v) it will, for the purposes of our paper, be convenient to conjunct ω -consistency and the disjunction property under the name of the *strong disjunction property* which is clearly equivalent to

$$\perp \neq \tau, \text{ and } \Box x \vee \Box y = \tau \text{ implies } x = \tau \text{ or } y = \tau.$$

Before doing so however we shall take a final look at each one of the former separately.

For the remainder of the \S we shall be confusing modal and arithmetic notation.

In 1971 Parikh [35] proved that the implication of the statement (iv) for Σ_1 -sound theories ($T \vdash \Box \varphi \Rightarrow T \vdash \varphi$) may take rather long to materialize. That is, for each provably recursive function g of T there exists a sentence φ and a proof p of $\Box \varphi$ in T s.t. no number $\leq g(p)$ is a proof of φ in T .

We shall prove the same for the disjunction property. Our proof leans heavily on techniques of de Jongh & Montagna [25] and Carbone [13] and an idea in Carbone & Montagna [14].

1.1. Proposition. Let g be a provably recursive function of a Σ_1 -sound theory T .

(a) There exist (Σ_1) sentences σ_1 and σ_2 and a proof p_0 of $\Box \sigma_1 \vee \Box \sigma_2$ in T s.t. $T \vdash \sigma_1$, $T \vdash \sigma_2$ and no $p_1 \leq g(p_0)$ is a proof of $\Box \sigma_1$ or of $\Box \sigma_2$ in T .


(b) There exist (Σ_1) sentences τ_1 and τ_2 and a proof q_0 of

$\Box\tau_1 \vee \Box\tau_2$ in T s.t. $T \vdash \tau_1$, $T \text{ non } \vdash \tau_2$ and no $q_1 \leq g(q_0)$ is a proof of $\Box\tau_1$ in T .

Proof. First we fix a pair of (Σ_1) sentences α and β s.t.

$$\begin{aligned} T \vdash (\Box\alpha \vee \Box\beta) &\leftrightarrow \\ \Box\Box\alpha &\leftrightarrow \\ \Box\Box\beta &\leftrightarrow \Box\Box\perp \end{aligned}$$

Sentences α and β satisfying these conditions could be produced with the help of Solovay's [47] Second Theorem applied to the following Kripke model (at each node, only the letters forced are shown):



(This model also appeared in Visser [49] to accomplish a similar task).

Now let $\Box\varphi <_g \Box\psi$ denote the formula saying that there exists a proof p of φ in T s.t. no $q \leq g(p)$ is a proof of ψ in T .

(a) By self-reference find a sentence ρ_1 s.t.

$$(1) T \vdash \rho_1 \leftrightarrow. [\Box(\Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta)) <_g \Box\Box(\rho_1 \vee \alpha)] \wedge \\ \wedge [\Box(\Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta)) <_g \Box\Box(\rho_1 \vee \beta)]$$

We have

$$(2) T \vdash \Box(\Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta)) \rightarrow. \Box^+\rho_1 \vee \Box^+\neg\rho_1$$

(the antecedent implies that the
r.h.s. of (1) is decidable and so
 ρ_1 is decidable)

$$(3) T \vdash \neg\rho_1 \rightarrow. \\ \rightarrow. \Box(\Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta)) \rightarrow (\Box\Box(\rho_1 \vee \alpha) \vee \Box\Box(\rho_1 \vee \beta))$$

(by (1))

- (4) $\mathbb{T} \vdash \Box^+ \neg \rho_1 \rightarrow$
 $\rightarrow \Box^+(\Box(\Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta)) \rightarrow \Box\Box(\rho_1 \vee \alpha) \vee \Box\Box(\rho_1 \vee \beta))$
(by (3))
 $\rightarrow \Box^+(\Box(\Box\alpha \vee \Box\beta) \rightarrow \Box\Box\alpha \vee \Box\Box\beta)$
 $\rightarrow \Box^+(\Box\Box\Box\perp \rightarrow \Box\Box\perp)$ (by the choice of α and β)
- (5) $\mathbb{T} \vdash \Box^+ \neg \rho_1 \rightarrow \Box\Box\Box\perp$ (from (4) by Löb's Theorem)
 $\rightarrow \Box\Box\perp$ (by (4))
- (6) $\mathbb{T} \vdash \Box\Box\perp \rightarrow \Box\alpha \vee \Box\beta$ (by the choice of α and β)
 $\rightarrow \Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta)$
- (7) $\mathbb{T} \vdash \Box^+ \neg \rho_1 \rightarrow \Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta)$
(by (5) and (6))
- (8) $\mathbb{T} \vdash \Box\Box\Box\rho_1 \rightarrow \Box\Box(\Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta))$
 $\rightarrow \Box(\Box\rho_1 \vee \Box^+ \neg \rho_1)$
(by (2))
 $\rightarrow \Box(\Box(\rho_1 \vee \alpha) \vee \Box(\rho_1 \vee \beta))$
(by (7))
 $\rightarrow \Box\rho_1 \vee \Box^+ \neg \rho_1$
(by (2))
 $\rightarrow \Box\rho_1 \vee \Box\Box\perp$
(by (5))
 $\rightarrow \Box\Box\rho_1$
- (9) $\mathbb{T} \vdash \Box\Box\rho_1$ (from (8) by Löb's Theorem)
- (10) $\mathbb{T} \vdash \rho_1$ (from (9) by Σ_1 -soundness)

By (10) also the r.h.s. of (1) is provable and hence by Σ_1 -soundness true. Now let $\sigma_1 \equiv \rho_1 \vee \alpha$, $\sigma_2 \equiv \rho_1 \vee \beta$ and note that (a) is proved.

(b) Construct a sentence ρ_2 s.t.

$$\mathbb{T} \vdash \rho_2 \leftrightarrow \Box(\Box(\rho_2 \vee \alpha) \vee \Box\beta) <_g \Box\Box(\rho_2 \vee \alpha)$$

and show that $\mathbb{T} \vdash \rho_2$ in perfect analogy with the proof of (a). Then take $\tau_1 \equiv \rho_2 \vee \alpha$ and $\tau_2 \equiv \beta$. ■

After the research underlying the present paper was essentially completed I learnt that Proposition 1.1 fell corollary to very general recent results of Montagna [33].

Leaving alone the problem of actually constructing from a proof of $\Box\varphi \vee \Box\psi$ that of one of the disjuncts, one might at least ask which one of those is true. The next proposition shows that this generally also is a very difficult question.

1.2. Proposition. *If T is a Σ_1 -sound theory then there is no provably recursive function of T which, given a proof in T of a sentence of the form $\Box\varphi \vee \Box\psi$, picks a true disjunct (even if one restricts the task to Σ_1 sentences φ and ψ).*

Proof. Suppose g were such a function. That is, if p is a proof of a sentence of the form $\Box\varphi \vee \Box\psi$ with φ and ψ in Σ_1 then

$$g(p) = 0 \Rightarrow T \vdash \varphi \quad \text{and} \quad g(p) = 1 \Rightarrow T \vdash \psi.$$

Clearly we can assume w.l.o.g. that

$$T \vdash \forall x (g(x) = 0 \vee g(x) = 1)$$

We introduce two ad hoc "modal" operators:

$$\begin{aligned} \Box_0\varphi &\equiv \Box\varphi \wedge g(\text{the least proof of } \varphi) = 0 \\ \Box_1\varphi &\equiv \Box\varphi \wedge g(\text{the least proof of } \varphi) = 1 \end{aligned}$$

Next we define by parallel self-reference:

$$\begin{aligned} T \vdash \sigma &\leftrightarrow \Box_1(\Box\sigma \vee \Box\tau) \\ T \vdash \tau &\leftrightarrow \Box_0(\Box\sigma \vee \Box\tau) \end{aligned}$$

We have

$$\begin{aligned} T \vdash \Box(\Box\sigma \vee \Box\tau) &\rightarrow (\Box_0 \vee \Box_1)(\Box\sigma \vee \Box\tau) \\ &\rightarrow \sigma \vee \tau \\ &\rightarrow \Box\sigma \vee \Box\tau \quad (\sigma \text{ and } \tau \text{ are } \Sigma_1) \end{aligned}$$

and hence

$$T \vdash \Box\sigma \vee \Box\tau$$

by Löb's Theorem. Now if $g(\text{the least proof of } \Box\sigma \vee \Box\tau) = 0$ then σ is false and therefore by Σ_1 -soundness $T \text{ non } \vdash \sigma$ contrary to the assumptions on g ; $g(\text{the least proof of } \Box\sigma \vee \Box\tau) = 1$ contradicts the assumptions in the symmetric manner. ■

§ 2. On conservativity in L

2.1. Definition. The degree $d(A)$ of a modal formula A is defined inductively:

$$\begin{aligned} d(p_i) &= d(\perp) = d(\top) = 0 \\ d(\neg A) &= d(A) \\ d(A \wedge B) &= d(A \vee B) = d(A \rightarrow B) = \max(d(A), d(B)) \\ d(\Box A) &= 1 + d(A) \end{aligned}$$

Thus, formulae of degree 0 are precisely the \Box -free formulae.

Let \vec{p} be a finite tuple of propositional letters. Formulae of degree $\leq n$ containing no letters other than in \vec{p} constitute (modulo L-equivalence) a finite Boolean algebra which we denote by $F^n(\vec{p})$. Elements of $F^n(\vec{p})$ will be persistently confused with modal formulas representing these elements. We also let $A^n(\vec{p})$ denote the set of atoms of $F^n(\vec{p})$. Clearly $F^n(\vec{p})$ is a subalgebra of $F^m(\vec{q})$ whenever $n \leq m$ and $\vec{p} \subseteq \vec{q}$. It is convenient to think of the modal operator \Box as sending elements of $F^n(\vec{p})$ to those of $F^{n+1}(\vec{p})$.

F and $F(\vec{p})$ denote the diagonalizable algebras of all formulae and of all formulae whose propositional letters are in \vec{p} respectively.

2.2. Lemma. Consider elements of $F^{n+1}(\vec{p})$ of the form

$$\alpha \wedge \Box \bigvee \gamma \wedge \bigwedge \diamond \gamma$$

with α ranging over $A^0(\vec{p})$ and γ ranging over subsets of $A^n(\vec{p})$. Call such formulas types. (Here $\diamond \gamma \equiv \{ \diamond C \mid C \in \gamma \}$)

(a) The conjunction of two distinct types is (L-equivalent to) \perp .

(b) Each formula in $F^{n+1}(\vec{p})$ is (L-equivalent to) a disjunction of types.

(c) Each formula in $A^{n+1}(\vec{p})$ is (L-equivalent to) a type.

(d) Each type either belongs to $A^{n+1}(\vec{p})$ or is (L-equivalent to) \perp .

Proof. (a) It is straightforward to show that

$$\begin{aligned} \vdash_{\mathbf{L}} (\alpha_1 \wedge \Box \mathbb{W} \gamma_1 \wedge \mathbb{M} \diamond \gamma_1) \wedge (\alpha_2 \wedge \Box \mathbb{W} \gamma_2 \wedge \mathbb{M} \diamond \gamma_2) &\leftrightarrow. \\ &\leftrightarrow. (\alpha_1 \wedge \alpha_2) \wedge (\Box \mathbb{W} \gamma_1 \wedge \Box \mathbb{W} \gamma_2) \wedge (\mathbb{M} \diamond \gamma_1 \wedge \mathbb{M} \diamond \gamma_2) \\ &\leftrightarrow. (\alpha_1 \wedge \alpha_2) \wedge \Box \mathbb{W} (\gamma_1 \cap \gamma_2) \wedge \mathbb{M} \diamond (\gamma_1 \cup \gamma_2) \end{aligned}$$

and the claim follows by an easy Kripke model argument.

(b) By the definition of $\mathbf{F}^{n+1}(\vec{p})$ every formula therein can be thought of as a lattice combination of elements α of $\mathbf{A}^0(\vec{p})$ and formulas of the form $\Box C$ and $\neg \Box C$ with $C \in \mathbf{F}^n(\vec{p})$ or, equivalently, $\Box \mathbb{W} \gamma$ and $\neg \Box \mathbb{W} \gamma$ with $\gamma \subseteq \mathbf{A}^n(\vec{p})$. Thus to prove the claim it will suffice to show that α , $\Box \mathbb{W} \gamma$ and $\neg \Box \mathbb{W} \gamma$ are L-equivalent to appropriate disjunctions of types and that the conjunction of two disjunctions of types can be L-equivalently brought into the form of a disjunction of types. This is unproblematic:

$$\begin{aligned} \vdash_{\mathbf{L}} \alpha &\leftrightarrow \mathbb{W} \{ \alpha \wedge \Box \mathbb{W} \delta \wedge \mathbb{M} \diamond \delta \mid \delta \subseteq \mathbf{A}^n(\vec{p}) \} \\ \vdash_{\mathbf{L}} \Box \mathbb{W} \gamma &\leftrightarrow \mathbb{W} \{ \beta \wedge \Box \mathbb{W} \delta \wedge \mathbb{M} \diamond \delta \mid \beta \in \mathbf{A}^0(\vec{p}), \delta \subseteq \gamma \} \\ \vdash_{\mathbf{L}} \neg \Box \mathbb{W} \gamma &\leftrightarrow \\ &\leftrightarrow \mathbb{W} \{ \beta \wedge \Box \mathbb{W} \delta \wedge \mathbb{M} \diamond \delta \mid \beta \in \mathbf{A}^0(\vec{p}), \delta \subseteq \mathbf{A}^n(\vec{p}), \delta \text{ non } \subseteq \gamma \} \end{aligned}$$

and, finally,

$$\begin{aligned} \vdash_{\mathbf{L}} \mathbb{W}_i (\alpha_i \wedge \Box \mathbb{W} \gamma_i \wedge \mathbb{M} \diamond \gamma_i) \wedge \mathbb{W}_j (\alpha_j \wedge \Box \mathbb{W} \gamma_j \wedge \mathbb{M} \diamond \gamma_j) &\leftrightarrow \\ &\leftrightarrow \mathbb{W}_{i,j} ((\alpha_i \wedge \Box \mathbb{W} \gamma_i \wedge \mathbb{M} \diamond \gamma_i) \wedge (\alpha_j \wedge \Box \mathbb{W} \gamma_j \wedge \mathbb{M} \diamond \gamma_j)) \end{aligned}$$

Since by (a) the conjunction of two types is L-equivalent to 1 and/or to a type we are done.

(c) and (d) follow easily from (a) and (b). ■

The types of Lemma 2.2 are essentially the same as the normal form formulas of Fine [17] and the n -S-characters of Gleit & Goldfarb [20].

2.3. Definition. Let \mathcal{K} be a rooted model. The unique element of $\mathbf{A}^n(\vec{p})$ forced in \mathcal{K} is called the (n, \vec{p}) -character of \mathcal{K} . If the (n, \vec{p}) -characters of two rooted models coincide then these models are said to be (n, \vec{p}) -twins.

2.4. Definition. If $\mathcal{K} = (K, R, *)$ is a Kripke model and $a \in K$ then $\mathcal{K}[a]$, the a -cone of \mathcal{K} , is the rooted model whose domain is the set $\{a\} \cup \{b \in K \mid a R b\}$ and the accessibility and forcing relations are R and $*$ restricted to this set respectively. A *proper cone* of \mathcal{K} is the a -cone of \mathcal{K} for some $a \in K$ which is not the root of \mathcal{K} .

The following lemma, although simple, will render us a number of valuable services. It should be compared with Theorem 1 of Fine [17].

2.5. Lemma (Fine Lemma).

(a) Two rooted \vec{p} -models are $(n+1, \vec{p})$ -twins iff
 (i) they are $(0, \vec{p})$ -twins and
 (ii) each proper cone of one of these models has got an (n, \vec{p}) -twin among the proper cones of the other model and vice versa.

(b) $(n+m, \vec{p})$ -twins are (n, \vec{p}) -twins.

Proof. (a) is easy.

(b) is proved by induction on m using (a). ■

2.6. Definition. Let \vec{p} be a finite tuple of propositional letters. A formula A is said to be \vec{p} -conservative over a formula B if for each $C \in \mathcal{F}(\vec{p})$ one has $\vdash_{\mathbf{L}} B \rightarrow C$ whenever $\vdash_{\mathbf{L}} A \rightarrow C$. A is *conservative over* B if it is Λ -conservative over B where Λ is the empty tuple. A is $(\vec{p}-)$ conservative if it is $(\vec{p}-)$ conservative over \top .

Our aim is to show that conservativity is decidable. In fact we shall obtain stronger results.

2.7. Definition. Let $\mathcal{K}_1 = (K_1, R_1, *_1)$ and $\mathcal{K}_2 = (K_2, R_2, *_2)$ be rooted models, $a \in K_1$ and assume K_1 and K_2 disjoint. By saying that we *graft* \mathcal{K}_2 above a (in \mathcal{K}_1) we mean that a new model is constructed whose domain is $K_1 \cup K_2$, the forcing relation

coincides with $\#_1 \cup \#_2$ on propositional letters and the accessibility relation R is defined by putting

$$b R c \Leftrightarrow \\ \Leftrightarrow b R_1 c \text{ or } b R_2 c \text{ or } ((b R_1 a \text{ or } b = a) \text{ and } c \in K_2)$$

Let $\mathcal{K} = (K, R, \#)$ be a rooted \vec{p} -model and $a \in K$. Suppose one grafts an isomorphic copy of the a -cone of \mathcal{K} above $b \in K$ in \mathcal{K} with $b R a$. Then the "old" nodes can be easily shown to force precisely the same modal formulae in the resulting model as they did in \mathcal{K} (cf. Artemov [3]). Suppose $\mathcal{K}' = (K', R', \#')$ is a \vec{p} -model obtained from \mathcal{K} by a finite number of graftings of the sort described and let there exist a forcing relation $\#^+$ extending $\#'$ s.t. $\mathcal{K}^+ = (K', R', \#^+)$ is a \vec{q} -model that forces a formula $A \in \mathbf{F}(\vec{q})$. Then we shall say that \mathcal{K} is *expandable to (a model of) A* and that \mathcal{K}^+ is an *expansion of \mathcal{K} to (a model of) A* .

2.8. Lemma. *Let $\vec{p} \subseteq \vec{q}$. If every finite rooted (treelike) \vec{p} -model of a formula $A \in \mathbf{F}(\vec{p})$ is expandable to a model of $B \in \mathbf{F}(\vec{q})$ then B is \vec{p} -conservative over A .*

Proof. Easy. ■

2.9. Definition. If $A \equiv \alpha \wedge \square \bigvee \gamma \wedge \bigwedge \diamond \gamma$ with $\alpha \in \mathbf{A}^0(\vec{p})$ and $\gamma \subseteq \mathbf{A}^n(\vec{p})$ and \mathcal{K} is a rooted \vec{p} -model forcing A then α is called the *real world part* of A and of \mathcal{K} ; elements of γ are the (n, \vec{p}) -possible worlds of A and of \mathcal{K} and γ itself is the (n, \vec{p}) -possible worlds part of A and of \mathcal{K} . The number of elements in γ is the (n, \vec{p}) -rank of A and of \mathcal{K} .

Clearly the real world part and the (n, \vec{p}) -possible worlds part of A and of each rooted \vec{p} -model are defined uniquely up to L-equivalence.

The following lemma may be thought of as an improvement on the Joint Satisfiability Theorem of Gleit & Goldfarb [20].

2.10. Lemma (Expansion Lemma). *Let $\vec{p} \subseteq \vec{q}$. To every $n \in \omega$ there corresponds an $N \in \omega$ s.t. every finite treelike \vec{p} -model of $B \in \mathbf{A}^N(\vec{p})$ is expandable to a model of $C \in \mathbf{A}^n(\vec{q})$ whenever*

$B \wedge C$ is irrefutable in L .

Proof. The claim is immediate for $n = 0$ (in this case we can take $N = 0$). For the remaining $n \in \omega$ we use induction on the $(n-1, \vec{q})$ -rank of C . When this rank is 0 and $N > 0$ the claim is once again obvious.

Thus, given an $r \neq 0$, we assume for induction₀ hypothesis that each finite treelike \vec{p} -model of $D \in \mathbf{A}^N(\vec{p})$ is expandable to a model of $E \in \mathbf{A}^n(\vec{q})$ once $D \wedge E$ is irrefutable in L and the $(n-1, \vec{q})$ -rank of E is smaller than r .

Now let $C \in \mathbf{A}^n(\vec{q})$ of $(n-1, \vec{q})$ -rank r be forced in a rooted model \mathcal{K} along with $B \in \mathbf{A}^{N+c}(\vec{p})$ and let \mathcal{K} be an arbitrary finite treelike \vec{p} -model of B . The constant c will be specified later. We are going to expand \mathcal{K} to a model of C . To avoid heavy notation we stipulate that \mathcal{K} retains its name throughout the process of expansion despite the changes it undergoes and, at intermediate stages, despite being neither a \vec{p} - nor a \vec{q} -model.

First we consider a particular case when the $(n-1, \vec{q})$ -rank of \mathcal{K} is greater than that of any of its proper cones. In this case we let $c = 1$.

Let a_1, \dots, a_m be the immediate successors of the root of \mathcal{K} . By the Fine Lemma (a) there exists a sequence $\mathcal{K}[b_1], \dots, \mathcal{K}[b_m]$ of proper cones of \mathcal{K} s.t. $\mathcal{K}[b_i]$ is an (N, \vec{p}) -twin of $\mathcal{K}[a_i]$, $1 \leq i \leq m$. Since the $(n-1, \vec{q})$ -rank of each of the $\mathcal{K}[b_i]$'s is smaller than r , the induction₀ hypothesis yields an expansion of $\mathcal{K}[a_i]$ to the (n, \vec{q}) -character of $\mathcal{K}[b_i]$. Now replace each of the $\mathcal{K}[a_i]$'s by the corresponding expansion (this is possible because \mathcal{K} is treelike). Analogously, each proper cone $\mathcal{K}[b]$ of \mathcal{K} has got an (N, \vec{p}) -twin among the proper cones of \mathcal{K} which is expandable to the (n, \vec{q}) -character of $\mathcal{K}[b]$. For each such $\mathcal{K}[b]$, graft a copy of the corresponding expansion above the root of \mathcal{K} . Finally extend the forcing relation at the root of \mathcal{K} in the obvious way.

We show that the resulting model is an (n, \vec{q}) -twin of \mathcal{K} . The real world parts of their (n, \vec{q}) -characters coincide by construction. That the proper cones of the model constructed have

$(n-1, \vec{q})$ -twins among the proper cones of \mathcal{H} follows from the fact that every proper cone of the new model is either an (n, \vec{q}) - (and therefore by the Fine Lemma (b) an $(n-1, \vec{q})$ -) twin of a proper cone of \mathcal{H} or is a proper cone of an (n, \vec{q}) -twin of a proper cone of \mathcal{H} (and hence by the Fine Lemma (a) an $(n-1, \vec{q})$ -twin of a proper cone of \mathcal{H}). As to the opposite direction, recall that we grafted in \mathcal{K} an (n, \vec{q}) -twin to each proper cone of \mathcal{H} . Finally, apply the Fine Lemma (a).

Now we drop the assumption on the $(n-1, \vec{q})$ -ranks of the proper cones of \mathcal{H} and increase c to 3, that is, we assume \mathcal{K} and \mathcal{H} to be $(N+3, \vec{p})$ -twins.

Our plan is as follows. We set the induction₀ hypothesis and the skills we acquired when treating the particular case to work and let these expand as many proper cones of \mathcal{K} as possible to the characters of the corresponding proper cones of \mathcal{H} . What remains unexpanded in \mathcal{K} after this first attack corresponds to proper cones of \mathcal{H} of $(n-1, \vec{q})$ -rank r and hence the $(n-1, \vec{q})$ -possible worlds of these cones have to be the same as those of \mathcal{H} itself. Thus, provided we have implanted all the $(n-1, \vec{q})$ -possible worlds of \mathcal{H} above each of the yet unexpanded nodes of \mathcal{K} , we only have to care that no $(n-1, \vec{q})$ -possible world alien to \mathcal{H} comes into existence when the forcing relation at these nodes is being extended to \vec{q} .

Our first move will be to classify the proper cones of \mathcal{K} . Thus, we call a proper cone $\mathcal{K}[a]$ along with its root a

- *frontier*, if there is an $(N+1, \vec{p})$ -twin $\mathcal{H}[b]$ of $\mathcal{K}[a]$ among the proper cones of \mathcal{H} s.t. the $(n-1, \vec{q})$ -rank of $\mathcal{H}[b]$ is r but each proper cone of $\mathcal{H}[b]$ is of a smaller $(n-1, \vec{q})$ -rank;
- *high*, if a is not frontier and there is an $(N+1, \vec{p})$ -twin of $\mathcal{K}[a]$ among the proper cones of \mathcal{H} of $(n-1, \vec{q})$ -rank smaller than r ;
- *low*, if a is not frontier and every $(N+1, \vec{p})$ -twin of $\mathcal{K}[a]$ among the proper cones of \mathcal{H} is of $(n-1, \vec{q})$ -rank r ;

- *genuinely frontier*, if a is frontier and every node which a is accessible from is low;
- *critically high*, if a is high and every node which a is accessible from is low;
- *essentially low*, if a is low and every node which a is accessible from is also low.

Claim 1. *Each proper cone of K is either frontier or high or low.*

Claim 2. *Of each node of K which is not the root of K exactly one of the following statements is true:*

- (i) a is genuinely frontier;
- (ii) a is critically high;
- (iii) a is essentially low;
- (iv) a is accessible from a genuinely frontier or from a critically high node.

Indeed, Claim 1 is easy. Claim 2 follows from Claim 1 by inspection of our classification.

Informally, we have this picture: To the root of K clings a downward closed collection of essentially low nodes and immediately above this collection there is a one-node-thick layer of genuinely frontier and critically high nodes which separates the essentially low nodes from the rest of the model.

Claim 3. *From each (essentially) low node a frontier node is accessible.*

The proof of Claim 3 explains why we chose c to be so abnormally large:

By the Fine Lemma (a) each low proper cone $K[a]$ has got at least one $(N+2, \vec{p})$ -twin among the proper cones of K . Each of these $(N+2, \vec{p})$ -twins has a proper cone of $(n-1, \vec{q})$ -rank r , or else a would be frontier. Pick one of these $(N+2, \vec{p})$ -twins and a proper cone $K[b]$ of it of $(n-1, \vec{q})$ -rank r s.t. each proper cone of $K[b]$ has a smaller $(n-1, \vec{q})$ -rank. By the Fine Lemma (a) the root of an $(N+1, \vec{p})$ -twin of $K[b]$ should be accessible from a . This root is

by definition a frontier node so Claim 3 is proved.

Let us now start working. The root of \mathcal{K} is as usual unproblematic. Next we replace each genuinely frontier and each critically high proper cone of \mathcal{K} by its expansion to the (n, \vec{q}) -character of one of those of its $(N+1, \vec{p})$ -twins in \mathcal{H} which this proper cone owes its frontier or high statute to respectively. For (genuinely) frontier proper cones such expansions were carried out when treating the easy particular case with $c = 1$ and for expansions of critically high nodes we turn to induction₀ hypothesis. By Claim 2 and since \mathcal{K} is treelike these replacements can not conflict. To each essentially low node a of \mathcal{K} we do the following: extend the forcing relation at a to that at the root of one of the $(N+1, \vec{p})$ -twins of $\mathcal{K}[a]$ in \mathcal{H} and graft above a an expansion of a frontier proper cone $\mathcal{K}[a_0]$ with a_0 accessible from a , which corresponds to a by Claim 3, to the (n, \vec{q}) -character of an $(N+1, \vec{p})$ -twin of $\mathcal{K}[a_0]$ in \mathcal{H} which enjoys $(n-1, \vec{q})$ -rank r but none of its proper cones does. Lastly, for each proper cone $\mathcal{H}[b]$ of \mathcal{H} s.t. every one of its proper cones has $(n-1, \vec{q})$ -rank smaller than r pick an $(N+1, \vec{p})$ -twin in (the original copy of) \mathcal{K} and graft above the root of \mathcal{K} an expansion of this $(N+1, \vec{p})$ -twin to the (n, \vec{q}) -character of $\mathcal{H}[b]$.

It is now easily seen from Claim 2 that \mathcal{K} has been metamorphosed into a \vec{q} -model. We check that the $(n-1, \vec{q})$ -possible worlds parts of \mathcal{H} and of the model constructed are the same.

If there is a proper cone of \mathcal{H} of $(n-1, \vec{q})$ -rank r then at least one of such cones enjoys an (n, \vec{q}) -twin in the modified \mathcal{K} grafted above the root. Since \mathcal{H} is itself of $(n-1, \vec{q})$ -rank r , each $(n-1, \vec{q})$ -possible world of \mathcal{H} is also an $(n-1, \vec{q})$ -possible world of this (n, \vec{q}) -twin and hence of the expanded \mathcal{K} . If there were no proper cones of \mathcal{H} of this $(n-1, \vec{q})$ -rank then we would have grafted in \mathcal{K} an (n, \vec{q}) -twin to each proper cone of \mathcal{H} and anyway this is the easy $c = 1$ case that we dealt away with earlier.

It remains to see that each $(n-1, \vec{q})$ -possible world of \mathcal{K} is that of \mathcal{H} . Expansions of genuinely frontier, critically high and frontier proper cones of \mathcal{K} grafted in \mathcal{K} present, as in the $c =$

1 case, no problem. We show by rootward induction₁ on the essentially low nodes of \mathcal{K} that these only gave rise to $(n-1, \vec{q})$ -possible worlds that are those of \mathcal{H} . Consider an essentially low node a of \mathcal{K} . Recall that there is an expansion of something to the (n, \vec{q}) -character of a proper cone of \mathcal{H} having $(n-1, \vec{q})$ -rank r grafted above a . Hence the $(n-1, \vec{q})$ -possible worlds part of the a -cone of the new \mathcal{K} is the same as that of \mathcal{H} : by induction₁ hypothesis no extra $(n-1, \vec{q})$ -possible world could have crept in. Find now the root b of the $(N+1, \vec{p})$ -twin $\mathcal{H}[b]$ of $\mathcal{K}[a]$ which the forcing relation at a was extended to. Since this $(N+1, \vec{p})$ -twin also had to have $(n-1, \vec{q})$ -rank r and hence the same $(n-1, \vec{q})$ -possible worlds as \mathcal{H} , we see by the Fine Lemma (a) that the a -cone of the modified \mathcal{K} is an (n, \vec{q}) -twin of $\mathcal{H}[b]$ which gives us the desiderata. Thus we have executed the induction₁ step and the proof is complete.

Since the $(n-1, \vec{q})$ -rank of a formula can not be greater than $|\mathbf{F}^{n-1}(\vec{q})|$ our proof yields $N = 1 + 3 \cdot |\mathbf{F}^{n-1}(\vec{q})|$. ■

2.11. Lemma. *Let $\vec{p} \subseteq \vec{q}$. For each formula $B \in \mathbf{F}(\vec{q})$ there exists a formula $C \in \mathbf{F}(\vec{p})$ s.t. $\vDash_{\mathbf{L}} B \rightarrow C$ and a finite tree-like \vec{p} -model is expandable to a model of B iff this model forces C .*

Proof. Let $B \in \mathbf{F}^n(\vec{q})$ and let N be the number which corresponds to n by the Expansion Lemma. Take C to be the disjunction of those elements D of $\mathbf{A}^N(\vec{p})$ whose conjunction with B is irrefutable in \mathbf{L} and use the Expansion Lemma. ■

We are now able to prove the converse to Lemma 2.8.

2.12. Lemma. *Suppose that $\vec{p} \subseteq \vec{q}$ and $B \in \mathbf{F}(\vec{q})$ is \vec{p} -conservative over $A \in \mathbf{F}(\vec{p})$. Then each finite tree-like \vec{p} -model of A is expandable to a model of B .*

Proof. By Lemma 2.11 there exists a formula $C \in \mathbf{F}(\vec{p})$ s.t. $\vDash_{\mathbf{L}} B \rightarrow C$ and each \vec{p} -model of C is expandable to a model of B . Since B is \vec{p} -conservative over A we have that $\vDash_{\mathbf{L}} A \rightarrow C$ and so

each finite treelike \vec{p} -model of A is expandable to a model of B . ■

Smoryński [44] establishes the Craig Interpolation Property for the modal logic L : If $\vdash_L A \rightarrow B$ then there exists a formula C s.t. $\vdash_L A \rightarrow C$ and $\vdash_L C \rightarrow B$ and C only contains propositional letters common to A and B (cf. also Boolos [11] and Gleit & Goldfarb [20]). The following corollary shows that all we need know of B to construct C is what propositional letters A and B have in common.

2.13. Corollary (Uniform Craig Interpolation Lemma for L). *Let $\vec{p} \subseteq \vec{q}$. Given a formula $B \in F(\vec{q})$ we can construct a formula $C \in F(\vec{p})$ s.t. $\vdash_L B \rightarrow C$ and $\vdash_L C \rightarrow D$ whenever \vec{r} is a tuple of propositional letters disjoint from \vec{q} and $D \in F(\vec{p}, \vec{r})$ is s.t. $\vdash_L B \rightarrow D$. Moreover, this formula C is unique up to L -equivalence.*

Proof. Let C be as in Lemma 2.11. Take a formula D meeting the requirements of the present corollary and let $E \in F(\vec{p})$ be the interpolant between B and D provided by the usual Craig Interpolation Lemma. We show $\vdash_L C \rightarrow E$ whence $\vdash_L C \rightarrow D$ follows by modus ponens. For if this were not the case then we would have a finite treelike model forcing $C \wedge \neg E$. By Lemma 2.11 this model would expand to a model of B and thus $B \wedge \neg E$ would be irrefutable in L contradicting the assumption that E is an interpolant.

Uniqueness is left to the reader. ■

Thus if $\vec{p} \subseteq \vec{q}$ and $B \in F(\vec{q})$ then among the formulas in $F(\vec{p})$ implied by B exists the strongest one.

For the case of \vec{p} an empty tuple Corollary 2.13 is essentially proved in Artemov [2] and [3]. The full strength of this corollary will not be needed until § 10.

2.14. Corollary. (\vec{p} -) conservativity is decidable.

Proof. To decide whether a formula A is \vec{p} -conservative over

a formula B construct the formula C provided by the Uniform Craig Interpolation Lemma s.t. $\vdash_{\mathbf{L}} A \rightarrow C$ and $\vdash_{\mathbf{L}} C \rightarrow D$ whenever $\vdash_{\mathbf{L}} A \rightarrow D$ and A and D do not have common propositional letters other than those in \vec{p} . Use the same lemma to see that A is \vec{p} -conservative over B iff $\vdash_{\mathbf{L}} B \rightarrow C$. ■

In what follows formalized versions of certain lemmas of the present § will appear within IS_1 without special notice. In each case the verification that such formalizations are possible is unproblematic and therefore left to the reader.

§ 3. A family of Kripke models

3.1. Definition. Let \mathcal{K} be a finite \vec{p} -model. \mathcal{K} is said to be *differentiated* if for each node a of \mathcal{K} there exists a formula $A \in \mathbf{F}(\vec{p})$ s.t. a is the only node in \mathcal{K} forcing A .

Note that for finite models our definition of differentiated is equivalent to that of Fine [17].

3.2. Definition. Let \mathcal{K} be a finite model. The least such $n \in \omega$ that $\mathcal{K} \models \square^n \perp$ is called the *height* of \mathcal{K} . Thus the height of \mathcal{K} is equal to the number of elements in the largest subset of the domain of \mathcal{K} linearly ordered by the accessibility relation. Clearly if \mathcal{K} is rooted then the height of \mathcal{K} exceeds that of any one of its proper cones.

3.3. Lemma. Let \mathcal{K} be a finite rooted differentiated \vec{p} -model and let $A \in \mathbf{F}(\vec{p})$.

(a) Each proper cone of \mathcal{K} is differentiated.

(b) (Fine [18]) To each finite rooted \vec{p} -model \mathcal{K} there corresponds a finite rooted differentiated \vec{p} -model which forces precisely the same formulas in $\mathbf{F}(\vec{p})$ as \mathcal{K} does.

(c) There exists a formula, which we shall denote $\Psi_{\mathcal{K}}(\vec{p})$ (or just $\Psi_{\mathcal{K}}$), s.t. any rooted differentiated \vec{p} -model \mathcal{H} is isomorphic to \mathcal{K} if and only if $\mathcal{H} \models \Psi_{\mathcal{K}}(\vec{p})$.

(d) $\mathcal{K} \models A$ iff $\vdash_{\mathbf{L}} \Psi_{\mathcal{K}}(\vec{p}) \rightarrow A$.

(e) Either $\vdash_{\mathbf{L}} \Psi_{\mathcal{K}}(\vec{p}) \rightarrow A$ or $\vdash_{\mathbf{L}} \Psi_{\mathcal{K}}(\vec{p}) \rightarrow \neg A$.

Proof. (a) Obvious.

(b) Let $\mathcal{H} = (H, R, *)$. Define an equivalence relation E on H :

$a E b \Leftrightarrow a$ and b force the same formulas in $\mathbf{F}(\vec{p})$.

Define R/E to be the relation on H/E which holds between two E -equivalence classes a and b whenever for each node $a \in a$ there exists a node $b \in b$ s.t. $a R b$. Clearly R/E is transitive and

irreflexive. Let an E -equivalence class a force a propositional letter $p_i \in \vec{p}$ ($a *_E p_i$) if a representative of a forces p_i .

We show by induction on the structure of A that if $a \in a$ then

$$a *_E A \text{ iff } a *_E A.$$

The only interesting induction step occurs when A is of the form $\Box B$. Suppose $a *_E \Box B$. If $a \overset{R}{/}_E \mathfrak{L}$ then for some $b \in \mathfrak{L}$ one has $a R b$ whence $b *_E B$. Hence by the induction hypothesis $\mathfrak{L} *_E B$. Conclude $a *_E \Box B$. The converse direction is equally easy.

Thus, \mathcal{K} and $(\overset{H}{/}_E, \overset{R}{/}_E, *_E)$ force the same modal formulas and trivially the new model is differentiated.

(c) We prove that for $\Psi_{\mathcal{K}}(\vec{p})$ one can take the (n, \vec{p}) -character of \mathcal{K} where n is the height of \mathcal{K} . This we do by induction on the height of \mathcal{K} .

So let the height of \mathcal{K} be $n + 1$ and let \mathcal{H} and \mathcal{K} be $(n+1, \vec{p})$ -twins. We construct a mapping f from the domain of \mathcal{K} to the domain of \mathcal{H} . Let f map the root of \mathcal{K} to that of \mathcal{H} . Next let f take the root of a proper cone $\mathcal{K}[a]$ of \mathcal{K} to the root of its (n, \vec{p}) -twin among the proper cones of \mathcal{H} (which exists by the Fine Lemma (a)). Note that by the induction hypothesis $\mathcal{K}[a]$ is isomorphic to $\mathcal{H}[f(a)]$. Since \mathcal{K} is differentiated f is injective for else there would exist two distinct but isomorphic proper cones of \mathcal{K} . Moreover f is surjective because each proper cone of \mathcal{H} enjoys an (n, \vec{p}) -twin among the proper cones of \mathcal{K} and since \mathcal{H} is differentiated f connects these two.

By the Fine Lemma (a) f preserves forcing of propositional letters. It remains to check that f respects the accessibility relation. Let, in \mathcal{K} , b be accessible from a in which case $\mathcal{K}[b]$ is isomorphic to some proper cone of $\mathcal{H}[f(a)]$ and in particular $\mathcal{K}[b]$ is an (n, \vec{p}) -twin of a proper cone of $\mathcal{H}[f(a)]$. So f must take b to the root of that proper cone and hence $f(b)$ is accessible from $f(a)$. A symmetric argument will establish that b is accessible from a whenever $f(b)$ is accessible from $f(a)$. This shows that f is an isomorphism and completes the proof of (c).

(d) (if) By (b) for any finite rooted \vec{p} -model \mathcal{H} forcing

$\Psi_{\mathcal{K}}(\vec{p})$ we can construct a finite rooted differentiated \vec{p} -model which forces the same formulas as \mathcal{K} . By (c) this model will be isomorphic to \mathcal{K} and will therefore force A . Hence $\mathcal{K} \# A$. By the Completeness Theorem for L we are done.

The (only if) direction is left to the reader.

(e) follows at once from (d). ■

Thus, formulas of the form $\Psi_{\mathcal{K}}(\vec{p})$ are atoms of $F(\vec{p})$. Moreover, it can be shown that each atom of $F(\vec{p})$ has this form.

Lemma 3.3(c) is proved in Artemov [4] for treelike models and a suitably adjusted notion of differentiated. To get differentiated models from Artemov differentiated models one only has to identify nodes that force the same formulae. Confer also Bellissima [7] for a related result.

3.4. Definition. Let $\mathcal{M}(\vec{p}) = (M(\vec{p}), R(\vec{p}), \#_{\vec{p}})$ denote the \vec{p} -model whose domain is constituted by all finite rooted differentiated \vec{p} -models (we shall henceforth denote these by lower case Roman letters) with the accessibility relation defined by

$a R(\vec{p}) b \Leftrightarrow b$ is isomorphic to a proper cone of a

and with $a \#_{\vec{p}} p_i$ iff $a \# p_i$ where $p_i \in \vec{p}$.

The models $\mathcal{M}(\vec{p})$ will be our favourite playground and an important tool for our embeddability results. In fact, these models can be shown isomorphic to the models employed by Grigolia [21] and [22] and Rybakov [39]. We collect some facts about $\mathcal{M}(\vec{p})$.

3.5. Lemma. Let $a, b \in \mathcal{M}(\vec{p})$, $A, B \in F^n(\vec{p})$.

(a) $a R(\vec{p}) b$ iff $\vdash_L \Psi_a(\vec{p}) \rightarrow \diamond \Psi_b(\vec{p})$.

(b) a non $R(\vec{p}) b$ iff $\vdash_L \Psi_a(\vec{p}) \rightarrow \Box \neg \Psi_b(\vec{p})$.

(c) $a \#_{\vec{p}} A$ iff $a \# A$.

(d) $\mathcal{M}(\vec{p})[a]$ is isomorphic to a .

(e) $M(\vec{p})$ is differentiated.

(f) $A \text{ non } \vdash_{\mathbf{L}} B$ iff there exists a node $c \in M(\vec{p})$ s.t. $c \vDash A$ and $c \vDash \neg B$.

(g) If $A \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a(\vec{p})$ and $A \vdash_{\mathbf{L}} B$ then $a \vDash B$.

(h) If $a R(\vec{p}) b$ and $A \vdash_{\mathbf{L}} \neg \Psi_b(\vec{p})$ then $A \vdash_{\mathbf{L}} \neg \Psi_a(\vec{p})$.

Proof. (a) and (b) follow from Lemma 3.3(d).

(c) is established by downward induction on $R(\vec{p})$ (since $R(\vec{p})^{-1}$ is clearly well founded). Assume that (c) holds for all $b \in M(\vec{p})$ s.t. $a R(\vec{p}) b$.

It will suffice to prove the claim for propositional letters and formulas of the form $\Box B$. Since the case of propositional letters is self-evident we turn to $\Box B$.

We have: $a \vDash_{\vec{p}} \Box B$ iff for each proper cone b of a there holds $b \vDash_{\vec{p}} B$, iff for each proper cone b of a there holds $b \vDash B$ (this is by the induction hypothesis), iff $a \vDash \Box B$ q.e.d.

(d) By (c) of the present lemma one has $a \vDash_{\vec{p}} \Psi_a$ ergo $M(\vec{p})[a] \vDash \Psi_a$ and hence by Lemma 3.3(c) $M(\vec{p})[a]$ is isomorphic to a .

(e) By (c) of the present lemma Ψ_a differentiates a from all the other nodes of $M(\vec{p})$.

(f) We only prove (only if). If not $\vdash_{\mathbf{L}} \Box^+ A \rightarrow \Box^+ B$ then there exists a finite rooted \vec{p} -model \mathcal{K} s.t. $\mathcal{K} \vDash A$ and $\mathcal{K} \vDash \neg \Box^+ B$. Thus there is a node d of \mathcal{K} s.t. $d \vDash \Box^+ A$ and $d \text{ non } \vDash B$. Apply to $\mathcal{K}[d]$ Lemma 3.3(b) to obtain the desired $c \in M(\vec{p})$.

(g) By (f) since $A \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a$ there exists a node $c \in M(\vec{p})$ s.t. $c \vDash \Box^+ A$ and $c \vDash \Psi_a$ whence by Lemma 3.3(c) $c = a$ and from $A \vdash_{\mathbf{L}} B$ we get $a \vDash B$.

(h) Suppose $A \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a$. Then by (f) there is a node $c \in M(\vec{p})$ s.t. $c \vDash \Box^+ A$ and $c \vDash \Psi_a$. By Lemma 3.3(c) $c = a$ whence $c R(\vec{p}) b$. So we have $b \vDash \Box^+ A$ and $b \vDash \Psi_b$, therefore by (f) $A \text{ non } \vdash_{\mathbf{L}} \neg \Psi_b$ contrary to assumptions. ■

Lemma 3.5(c) permits us to drop the notational distinction between $\ast_{\vec{p}}$ and \ast .

We shall also need to know something about the interrelations between the models $M(\vec{p})$ with different \vec{p} .

3.6. Definition. Let $\vec{p} \subseteq \vec{q}$. We define a relation \triangleleft between nodes of $M(\vec{q})$ and those of $M(\vec{p})$. For $a \in M(\vec{q})$ and $b \in M(\vec{p})$ put

$$a \triangleleft b \Leftrightarrow a \text{ and } b \text{ force the same formulas in } F(\vec{p}).$$

Thus $a \triangleleft b$ if and only if b is expandable to a model of Ψ_a .

3.7. Lemma. Let $A \in F$, $\vec{p} \subseteq \vec{q}$, $a, b \in M(\vec{p})$, $c \in M(\vec{q})$.

$$(a) \ c \triangleleft a \text{ iff } \Vdash_{\mathbf{L}} \Psi_c(\vec{q}) \rightarrow \Psi_a(\vec{p}).$$

(b) There exists a unique $d \in M(\vec{p})$ s.t. $c \triangleleft d$.

(c) There exist only finitely many $e \in M(\vec{q})$ s.t. $e \triangleleft a$.

$$(d) \ \Vdash_{\mathbf{L}} \Psi_a(\vec{p}) \leftrightarrow \bigvee \{ \Psi_e(\vec{q}) \mid e \in M(\vec{q}), e \triangleleft a \}.$$

(e) If $c \triangleleft a$ and $A \Vdash_{\mathbf{L}} \neg \Psi_a(\vec{p})$ then $A \Vdash_{\mathbf{L}} \neg \Psi_c(\vec{q})$.

(f) If $a R(\vec{p}) b$ and $c \triangleleft a$ then there exists a node $e \in M(\vec{q})$ s.t. $c R(\vec{q}) e$ and $e \triangleleft b$.

(g) If $A \text{ non } \Vdash_{\mathbf{L}} \neg \Psi_a(\vec{p})$ then there exists a node $e \in M(\vec{q})$ s.t. $e \triangleleft a$ and $A \text{ non } \Vdash_{\mathbf{L}} \neg \Psi_e(\vec{q})$.

Proof. (a) (only if) Since $c \triangleleft a$ the node c forces the same elements of $F(\vec{p})$ as a does. In particular $c \Vdash \Psi_a(\vec{p})$. Hence by Lemma 3.3(d) $\Vdash_{\mathbf{L}} \Psi_c(\vec{q}) \rightarrow \Psi_a(\vec{p})$.

(if) Suppose a forces a formula $B \in F(\vec{p})$. Then by Lemma 3.3(d) we have $\Vdash_{\mathbf{L}} \Psi_a \rightarrow B$. Hence $\Vdash_{\mathbf{L}} \Psi_c \rightarrow B$ and therefore c forces B by Lemma 3.3(d).

(b) For existence, apply Lemma 3.3(b) to the model obtained from d by restricting the forcing relation to $F(\vec{p})$. Uniqueness follows by Lemma 3.5(e).

(c) If $e \triangleleft a$ then clearly the height of e is the same as that of a and there can only exist finitely many finite

differentiated \vec{q} -models of a given height.

(d) (\leftarrow) follows from (a).

(\rightarrow) Let \mathcal{K} be a finite rooted \vec{q} -model forcing Ψ_a . By Lemma 3.3(b) there exists a finite rooted differentiated \vec{q} -model forcing the same formulas in $\mathbf{F}(\vec{q})$ as \mathcal{K} does and hence it will force Ψ_e for some $e \in M(\vec{q})$ s.t. $e \triangleleft a$. Therefore $\mathcal{K} \Vdash \Psi_e$ whence $\mathcal{K} \Vdash \bigvee \{ \Psi_e \mid e \in M(\vec{q}), e \triangleleft a \}$. An application of the Completeness Theorem for \mathbf{L} completes the proof.

(e) follows from (a).

(f) By Lemma 3.5(a) we have $\vdash_{\mathbf{L}} \Psi_a \rightarrow \diamond \Psi_b$. Since $c \triangleleft a$ we also have $\vdash_{\mathbf{L}} \Psi_c \rightarrow \diamond \Psi_b$ by (a) of the present lemma. Hence $c \Vdash \diamond \Psi_b$ and there is a node $e \in M(\vec{q})$ s.t. $c R(\vec{q}) e$ and $e \Vdash \Psi_b$. By (a) and Lemma 3.3(d) this implies $e \triangleleft b$.

(g) follows from (d). ■

§ 4. Finite credibility extent

4.1. Theorem. *Let the credibility extent of a theory T be $n \in \omega$. A denumerable diagonalizable algebra \mathfrak{D} is isomorphic to an r.e. subalgebra of \mathfrak{D}_T iff*

- (i) \mathfrak{D} is positive and
- (ii) the height of \mathfrak{D} is n .

The (only if) direction is straightforward. The present § is devoted to the proof of the converse implication. Thus we are given a denumerable positive diagonalizable algebra \mathfrak{D} which we have to show isomorphic to an r.e. subalgebra of \mathfrak{D}_T .

To this end we have to borrow some notation from §§ 2 and 3. But first we simplify it a little bit. The tuple (p_1, \dots, p_i) of propositional letters will usually be represented by the symbol i . So $F^n(i)$ will stand for $F^n(p_1, \dots, p_i)$; M_i will stand for the domain of the model $\mathcal{M}(p_1, \dots, p_i)$ etc. We shall allow ourselves to omit the subscripts in \ast_i and R_i since it will always be clear which model is meant. We stipulate further that 0 is not an element of any of the M_i 's, $0 \in R_i$ (any element of M_i), $0 \triangleleft 0$ and $0 \ast A$ for no modal formula A (thus $a \ast \neg A$ is not the same as $a \text{ non } \ast A$ unless we assume $a \neq 0$). Moreover we shall be confusing the names of sets, relations and properties introduced in §§ 2 and 3 such as M_i , \vdash_L , "a formula A is conservative over a formula B " etc. with the names of their (honest) Δ_0 binumerations in arithmetic.

We are now going to apply a variant of the Solovay construction (see Solovay [47]) to each of the models M_i . We start with $i = 0$.

By self-reference define a Δ_0 function symbol h_0 and a closed ε -term ℓ_0 s.t. $I\Sigma_1$ proves the following clauses:

- (1) $h_0(0) = 0$
 - (2)
- $h_0(x+1) = a$ if (i) $a \in M_0$;
(ii) $a \ast \square_1^0$;

- (iii) $h_0(x) R a$ and
 (iv) $\text{Prf}(x, \ell_0 = \bar{a} \rightarrow \exists y (\overline{h_0(x)} R h_0(y) R \bar{a}))$
 $= h_0(x)$ if no a satisfying (i) - (iv) exists

(3) $\ell_0 = \lim_{x \rightarrow \infty} h_0(x)$ if h_0 reaches a limit
 $= 0$ otherwise

Here $\text{Prf}(\cdot, \cdot)$ is the proof predicate of the theory T under consideration.

4.2. Lemma ($I\Sigma_1$).

- (a) $\forall x \forall y (x \leq y \rightarrow h_0(x) = h_0(y) \vee h_0(x) R h_0(y))$
 (b) $\ell_0 = \lim_{x \rightarrow \infty} h_0(x)$
 (c) $\ell_0 = 0 \vee \ell_0 \# \square^{\bar{n}} \perp$
 (d) $\forall x (h_0(x) = \ell_0 \vee h_0(x) R \ell_0)$
 (e) $\forall a \in M_0 (a \# \square^{\bar{n}} \perp \wedge \ell_0 R a \rightarrow$
 $\rightarrow \neg \text{Pr}(\ell_0 = \bar{a} \rightarrow \exists y (\overline{\ell_0} R h_0(y) R \bar{a})))$
 (f) $\forall a \in M_0 (a \# \square^{\bar{n}} \perp \wedge \ell_0 R a \rightarrow \neg \text{Pr}(\ell_0 \neq \bar{a}))$
 (g) $\ell_0 \neq 0 \rightarrow$
 $\rightarrow \exists x \text{Prf}(x, \ell_0 = \overline{\ell_0} \rightarrow \exists y (\overline{h_0(x)} R h_0(y) R \overline{\ell_0}))$
 (h) $\ell_0 \neq 0 \rightarrow \text{Pr}(\ell_0 \neq \overline{\ell_0})$
 (i) $\ell_0 \neq 0 \rightarrow \text{Pr}(\overline{\ell_0} R \ell_0)$

Proof. (a) follows from inspection of (2) by induction on y .

(b) is immediate from (a) because h can make at most n moves.

(c) By (1) and (2) for each x we either have $h_0(x) = 0$ or $h_0 \# \square^{\bar{n}} \perp$ (this is established by induction on x). Now use (b).

(d) follows from (a) and (b).

(e) Consider an x s.t. $h_0(x) = \ell_0$. By (a) and since R verifiably is a strict partial order there holds $h_0(y) = \ell_0$ for

each $y \geq x$. Therefore there can not exist a proof $y \geq x$ of the formula

$$l_0 = \bar{a} \rightarrow \exists y (\bar{l}_0 R h_0(y) R \bar{a})$$

because then by (2) $h_0(y+1) = a$. Finally, recall that each provable sentence is provable by arbitrarily large proofs.

(f) is immediate from (e).

(g) Since $h_0(0) = 0$ and for some y $h_0(y) = l_0 \neq 0$ there exists an x s.t. $h_0(x) \neq h_0(x+1) = l_0$. By (iv) of (2) this x has to be a proof of the formula

$$l_0 = \bar{l}_0 \rightarrow \exists y (\overline{h_0(x)} R h_0(y) R \bar{l}_0)$$

(h) Once we assume $l_0 \neq 0$ we have by (g) that there is a proof x of the formula

$$l_0 = \bar{l}_0 \rightarrow \exists y (\overline{h_0(x)} R h_0(y) R \bar{l}_0)$$

Clearly by (2) and (d) $h_0(x+1) = l_0$. Moreover since h_0 is Δ_0 and R is a strict partial order we have that (a) formalized implies

$$\Pr(\neg \exists y (\overline{h_0(x)} R h_0(y) R \bar{l}_0))$$

and therefore $\Pr(l_0 \neq \bar{l}_0)$.

(i) follows from (b) and (d) formalized and (h). ■

4.3. Lemma.

(a) For each m s.t. $0 < m \leq n$ one has

$$I\Sigma_1 \vdash \Pr^m(\bar{1}) \leftrightarrow l_0 \neq \square^{\bar{m}} \perp$$

(b) $I\Sigma_1 + \Pr^n(\bar{1}) \vdash l_0 \neq 0$

(c) For no $a \in M_0$ s.t. $a \neq \square^n \perp$ do we have $T \vdash l_0 \neq \bar{a}$.

(d) $N \vDash l_0 = 0$

Proof. (a) Consider $m = 1$.

If $\Pr(\bar{1})$ then by Lemma 4.2(f) we have $l_0 R a$ for no $a \in M_0$. Hence $l_0 \neq \square \perp$.

Conversely, if $\ell_0 \neq \perp$ then by Lemma 4.2(i) $\Pr(\overline{\ell_0 R \ell_0})$ and therefore $\Pr(\overline{\forall a \ell_0 \neq a})$ whence $\Pr(\bar{1})$.

Now use induction on m .

Suppose $m < n$ and $\Pr^{m+1}(\bar{1})$, that is $\Pr(\overline{\Pr^m(\bar{1})})$. By the induction hypothesis this is equivalent to $\Pr(\ell_0 \neq \bar{m}_1)$. In other words, for all $a \in M_0$ s.t. $a \neq \perp$ and $a \text{ non } \neq \perp$ we have $\Pr(\ell_0 \neq \bar{a})$. By Lemma 4.2(f) no such a is accessible from ℓ_0 and hence $\ell_0 \neq \perp^{m+1}$.

As to the converse implication, $\ell_0 \neq \perp^{m+1}$ implies $\ell_0 \neq 0$ whence by Lemma 4.2(i) $\Pr(\overline{\ell_0 R \ell_0})$ which entails $\Pr(\ell_0 \neq \bar{m}_1)$ ergo $\Pr^{m+1}(\bar{1})$.

(b) follows immediately from (a).

(c) Assume $T \vdash \ell_0 \neq \bar{a}$ and reason in IS_1 : One has $\Pr(\ell_0 \neq \bar{a})$ and hence by Lemma 4.2(f) $\ell_0 \text{ non } R a$. In particular $\ell_0 \neq 0$. But then by (a) and by Lemma 4.2.(c) $\Pr^N(\bar{1})$.

Now step outside IS_1 . From $\text{IS}_1 \vdash \Pr^N(\bar{1})$ by Σ_1 -soundness we obtain $T \vdash \Pr^{N-1}(\bar{1})$ which contradicts the assumption on the credibility extent of T .

(d) follows from (c) of the present lemma and Lemma 4.2(c) and (h). ■

Now we have to do "the same" to the models M_i with $i > 0$. From a straightforward rewriting of (1) - (3) with i instead of 0 we could however only extract an embedding into \mathcal{D}_T of the diagonalizable algebra on i generators which is free in the variety of diagonalizable algebras of height $\leq n$. To insure that the extra relations required by the structure of \mathcal{D} be provable in T we have to restrict the range of the Solovay function h_i traveling in M_i (and therefore the possible values of ℓ_i) to a set of nodes smaller than the whole of M_i . The relevant subset of M_i can generally not be singled out by a condition decidable uniformly in i . (It can be done if the algebra \mathcal{D} we are dealing

with is finitely generated.) We therefore use decidable approximations to this set which can be made uniform in i so that the model \mathcal{M}_i is seen by h_i as changing with the passage of time. The approximating conditions on the set of nodes accessible to h will be given the form of the requirement that these nodes be expandable to models of a certain formula. The first variant of Solovay construction employing changing models is due to Jumelet [27]. The models in Jumelet [27] grew; ours will diminish.

It will be important for the success of our enterprises that the models stop changing as soon as it becomes clear that h is going to leave 0 as was the case in Jumelet [27]. A farewell to 0 however can only be bid at a nonstandard moment and so in the meantime we will have obtained the proofs of all the sentences needed to mimick the structure of \mathfrak{D} because we only care about standard proofs.

It should also be kept in mind that we do not want to embed into \mathfrak{D}_T the finitely generated subalgebras of \mathfrak{D} in arbitrary unrelated ways. In fact we would like the embedding of the subalgebra of \mathfrak{D} generated by the first $i + 1$ generators (in some fixed enumeration of those) to prolong the embedding of the subalgebra generated by the first i generators. To achieve this a kind of provable coherence between Solovay functions h_i climbing up different models \mathcal{M}_i is needed. Recall that the model \mathcal{M}_{i+1} is a refinement of \mathcal{M}_i in the sense made precise by Lemma 3.7. Roughly speaking, each node of \mathcal{M}_i falls into several nodes of \mathcal{M}_{i+1} . We want l_{i+1} to be a refinement of l_i in the same sense. Put formally, $l_{i+1} \triangleleft l_i$. Actually in our construction h_{i+1} will move step in step with h_i , that is $h_{i+1}(x) \triangleleft h_i(x)$. It is to maintain this kind of synchronicity that the extra property 4.2(e) which the usual Solovay function does not seem to possess is used.

Since the algebra \mathfrak{D} is positive a positive numeration $\nu: \omega - \{0\} \rightarrow \mathfrak{D}$ is available (here we have only subtracted 0 from $\text{dom } \nu$ for technical convenience). Let $\{A(m)\}_{m \in \omega}$ be a Δ_0 enumeration of the set of diagonalizable polynomials $A(p_1, p_2, \dots)$ that hit τ of \mathfrak{D} when νi is substituted for p_i . We rearrange this

- (iii) $h(i+1, x) R a$;
- (iv) $a \triangleleft h(i, x+1)$;
- (v) if $h(i+1, x) = 0$
then $D(g(x)) \text{ non } \vdash_L \neg \Psi_a$;
- (vi) for each b satisfying (i) - (v) in place
of a one has

$$\begin{aligned} & \forall z \leq x \text{ (Prf } (z, \ell(\bar{i}+1) = \bar{b} \rightarrow \exists y (\bar{h}(i+1, x) R h(\bar{i}+1, y) R \bar{b})) \rightarrow \\ & \rightarrow \exists w \leq z \text{ Prf } (w, \ell(\bar{i}+1) = \bar{a} \rightarrow \exists y (\bar{h}(i+1, x) R h(\bar{i}+1, y) R \bar{a}))) \\ & \text{(vii) } a \text{ is minimal among those } c \text{ that satisfy} \\ & \text{(i) - (vi) in place of } a \text{ (here "minimal" refers to the natural ordering of integers)} \\ & = h(i+1, x) \text{ if no } a \text{ satisfying (i) - (vi) exists} \end{aligned}$$

$$(11) \quad \ell(0) = \ell_0$$

$$(12) \quad \ell(i+1) = \lim_{x \rightarrow \infty} h(i+1, x) \text{ if } h(i+1, \cdot) \text{ reaches a limit} \\ = 0 \text{ otherwise}$$

Of course (vii) of (10) is just another way to say that we do not care what $h(i+1, x+1)$ is as long as it satisfies (i) - (vi). The weakly monotonically increasing function g occurring in (v) of (10) is Δ_0 and will be defined later.

4.5. Lemma ($I\Sigma_1$).

- (a) $\forall i \forall x \forall y (x \leq y \rightarrow h(i, x) = h(i, y) \vee h(i, x) R h(i, y))$
- (b) $\forall i \forall x h(i+1, x) \triangleleft h(i, x)$
- (c) $\forall j \forall i < j \forall x h(j, x) \triangleleft h(i, x)$
- (d) $\forall i \ell(i) = \lim_{x \rightarrow \infty} h(i, x)$
- (e) $\forall i \forall x (h(i, x) R \ell(i) \vee h(i, x) = \ell(i))$
- (f) $\forall j \forall i < j \ell(j) \triangleleft \ell(i)$

Proof. (a) For $i = 0$, use Lemma 4.2(a) and (8) and for $i > 0$ inspect (10) and apply induction on y .

(b) Note that since $0 \triangleleft 0$ the claim holds for $x = 0$ by (9) and assume $h(i+1, x) \triangleleft h(i, x)$ for (Δ_0) induction hypothe-

sis. We shall prove $h(i+1, x+1) \triangleleft h(i, x+1)$. If $h(i, x) = h(i, x+1)$ then the induction step is trivial (see (ii) of (10)). So assume $h(i, x) R h(i, x+1)$.

Case 1. $h(i, x) \neq 0$

Since $h(i, x) R h(i, x+1)$ we have by Lemma 3.7(f) that there exists a node $a \in M_{i+1}$ s.t. $h(i+1, x) R a \triangleleft h(i, x+1)$. The existence of a node a satisfying in addition (vi) and (vii) of (10) follows by the (Δ_0) least number principle applied first to proofs and then to nodes of M_{i+1} . Hence $h(i+1, x+1) = a \triangleleft h(i, x+1)$ so we are done.

Case 2. $h(i, x) = 0$

For $i > 0$, proceed as in Case 1 but use Lemma 3.7(g) instead of 3.7(f). For $i = 0$, recall that by Lemma 4.4(b) we have $D(x) \text{ non } \vdash_L \neg \Psi_a$, all x , all $a \in M_0$ s.t. $a \# \square^N \perp$ and hence $D(g(x)) \text{ non } \vdash_L \neg \Psi_a$. Therefore by Lemma 3.7(d) there holds $D(g(x)) \text{ non } \vdash_L \neg \Psi_b$ for some $b \in M_1$ s.t. $b \triangleleft a$.

(c) is proved with the help of (b) by (Π_1) induction on j .

(d) By Lemma 4.2(b) and (8) pick an x s.t. $\forall y \geq x \ h(0, y) = \ell_0$. By (a) and (c) $\forall y \geq x \ h(i, y) = h(i, x)$ and the claim follows by (12).

(e) follows from (a) and (d).

(f) follows from (c) and (d) without any induction. ■

By (11) and Lemma 4.5(f) we can introduce a sentence $\ell = 0$ as an abbreviation of any of the sentences

$$\ell_0 = 0, \quad \forall i \ \ell(i) = 0 \quad \text{and} \quad \exists i \ \ell(i) = 0$$

4.6. Lemma $(I\Sigma_1 + \ell \neq 0)$.

(a) $\forall i \ \text{Pr} \ (\overline{\ell(i)} R \ell(\bar{i}))$

(b) $\forall i \ \forall a \in M_i \ (a \# \square^N \perp \wedge \ell(i) R a \rightarrow$

$\rightarrow \neg \text{Pr} \ (\ell(\bar{i}) = \bar{a} \rightarrow \exists y \ (\overline{\ell(i)} R h(\bar{i}, y) R \bar{a}))$)

(c) $\forall i \forall a \in M_i (a \# \square_{\perp}^{\bar{n}} \wedge \ell(i) R a \rightarrow \neg \text{Pr}(\overline{\ell(\bar{i}) \neq \bar{a}}))$

Proof. (a) By Lemma 4.5(d) and (e) we only have to prove $\text{Pr}(\overline{\ell(\bar{i}) \neq \ell(\bar{i})})$. From $\ell \neq 0$ we get by (11) and by Lemma 4.2(h) $\text{Pr}(\overline{\ell(\bar{0}) \neq \ell(0)})$ whence by Lemma 4.5(f) formalized $\text{Pr}(\overline{\ell(\bar{i}) \neq \ell(\bar{i})})$.

(b) Please note that the formula

$\forall a \in M_i (a \# \square_{\perp}^{\bar{n}} \wedge \ell(i) R a \rightarrow \overline{\rightarrow \neg \text{Pr}(\ell(\bar{i}) = \bar{a} \rightarrow \exists y (\ell(\bar{i}) R h(\bar{i}, y) R \bar{a}))})$

is $\Delta_0(\Sigma_1)$ over $\text{I}\Sigma_1$ because $\ell \neq 0$ is equivalent to the Σ_1 formula $\exists x h(i, x) \neq 0$, the formula $\ell(i) R a$ is equivalent to the Π_1 formula $\forall x h(i, x) R a$ and the quantifier $\forall a \in M_i$ is primitive recursively bounded by the condition $a \# \square_{\perp}^{\bar{n}}$. In view of this we shall apply induction on i .

For $i = 0$ the claim follows by Lemma 4.2(e) and (11). Assume that it holds for i and suppose a reductio that $\ell(i+1) R a$ and

$\text{Pr}(\overline{\ell(\bar{i}+1) = \bar{a} \rightarrow \exists y (\ell(\bar{i}+1) R h(\bar{i}+1, y) R \bar{a})})$

Let $c \in M_i$ be s.t. $a \triangleleft c$ (see Lemma 3.7(b)) and hence $\ell(i+1) R a \triangleleft c$. By the (Δ_0) least number principle we obtain a node $b \in M_i$ s.t. $\ell(i+1) R b \triangleleft c$ and

$\text{Pr}(\overline{\ell(\bar{i}+1) = \bar{b} \rightarrow \exists y (\ell(\bar{i}+1) R h(\bar{i}+1, y) R \bar{b})})$

satisfying the conditions (vi) and (vii) of (10). Note that $\ell(i) R c$.

Now if $h(i, \cdot)$ were to jump from $\ell(i)$ directly to c then $h(i+1, \cdot)$ would have to jump directly to b because all the conditions (i) - (vii) of (10) would be satisfied (in particular, (v) would hold because $\ell(i+1) \neq 0$). This argument is formalizable in $\text{I}\Sigma_1$ and so we obtain

$\text{Pr}(\overline{\ell(\bar{i}) = \bar{c} \rightarrow \exists y (\ell(\bar{i}) R h(\bar{i}, y) R \bar{c}) \vee \ell(\bar{i}+1) = \bar{b}})$

Combining this with

$$\text{Pr } (\overline{\ell(\bar{i}+1)} = \bar{b} \rightarrow \exists y (\overline{\ell(\bar{i}+1)} R h(\bar{i}+1, y) R \bar{b}))$$

and with Lemma 4.5(b) formalized we get

$$\text{Pr } (\overline{\ell(\bar{i})} = \bar{c} \rightarrow \exists y (\overline{\ell(\bar{i})} R h(\bar{i}, y) R \bar{c}))$$

contrary to the induction hypothesis.

(c) follows immediately from (b). ■

Time is now ripe to define the primitive recursive function g .

- (13) $g(x) = z$ if (i) $z < x$;
(ii) there exists an $i \in \omega$ and a node $a \in M_i$ s.t. $a \# \square^{\bar{n}_i}$,

$$\text{Prf } (\overline{z}, \overline{\ell(\bar{i})} = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a}))$$
and $D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a$;
(iii) z is minimal among those satisfying (i) and (ii)
 $= x$ if no such z exists

4.7. Lemma ($\text{I}\Sigma_1$).

(a) $\forall x \forall y (x \leq y \rightarrow D(g(y)) \vdash_{\mathbf{L}} D(g(x)))$

(b) $\forall i \forall x \forall a \in M_i (a \# \square^{\bar{n}_i} \wedge$

$$\wedge \text{Prf } (x, \overline{\ell(\bar{i})} = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \wedge$$

$$\wedge D(g(x)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a \rightarrow \forall y \geq x g(y) = g(x))$$

(c) $\forall y (g(y) \neq y \rightarrow \exists i \exists a \in M_i (a \# \square^{\bar{n}_i} \wedge$

$$\wedge \text{Pr } (\overline{\ell(\bar{i})} = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \wedge \forall z D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a))$$

Proof. (a) We clearly have $g(x) \leq g(y)$. Now recall Lemma 4.4(a).

(b) Suppose the antecedent holds. By the (Δ_0) least number principle find the least $z \leq x$ satisfying the antecedent in place of x . By (13) it is seen that $\forall y \geq z g(y) = z$.

(c) On inspection of (13) one sees that if $g(y) \neq y$ then

for certain $i \in \omega$, $x < y$ and $a \in M_i$ the antecedent of (b) holds. Since by (b) we have that $\forall z \geq x \ g(z) = g(x)$ it is seen with the help of (a) that $\forall z \ D(g(x)) \vdash_{\mathbf{L}} D(g(z))$ and hence $\forall z \ D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a$ q.e.d. \blacksquare

4.8. Lemma ($\text{I}\Sigma_1$).

(a) $\ell \neq 0 \rightarrow$

$\rightarrow \forall i \ \forall x \ \forall A \in \mathbf{F}(i) \ (h(i, x) = 0 \wedge D(g(x)) \vdash_{\mathbf{L}} A \rightarrow \ell(i) \neq A)$

(b) $\ell = 0 \rightarrow \forall i \ \forall a \in M_i \ (a \neq \square^{\bar{n}} \perp \wedge$

$\wedge \text{Pr}(\ell(\bar{i}) = \bar{a} \rightarrow \exists y \ (0 \ R \ h(\bar{i}, y) \ R \ \bar{a})) \rightarrow \exists x \ D(g(x)) \vdash_{\mathbf{L}} \neg \Psi_a)$

(c) $\ell = 0 \rightarrow$

$\rightarrow \forall i \ \forall a \in M_i \ (a \neq \square^{\bar{n}} \perp \wedge \text{Pr}(\ell(\bar{i}) \neq \bar{a}) \rightarrow \exists x \ D(g(x)) \vdash_{\mathbf{L}} \neg \Psi_a)$

Proof. (a) From $\ell \neq 0$ we have $\ell(i) \neq 0$. Consider the $z \geq x$ s.t. $0 = h(i, z) \neq h(i, z+1)$. By inspection of (v) of (10) one has $D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_{h(i, z+1)}$ and from Lemma 4.7(a) we get $D(g(z)) \vdash_{\mathbf{L}} A$. Now by Lemma 3.5(g) there holds $h(i, z+1) \neq A$ whence by Lemma 4.5(e) $\ell(i) \neq A$.

(b) The proof is much the same as that of Lemma 4.6(b). We proceed by $(\Delta_0(\Sigma_1))$ induction on i . The case when $i = 0$ is immediate by Lemma 4.2(e).

So we assume the claim to hold for i , deny it for $i + 1$ and seek for a contradiction. We have that $\ell = 0$ and for a node $a \in M_{i+1}$ s.t. $a \neq \square^{\bar{n}} \perp$ and a suitable z there holds

$\text{Prf}(z, \ell(\bar{i}+1) = \bar{a} \rightarrow \exists y \ (0 \ R \ h(\bar{i}+1, y) \ R \ \bar{a}))$ and
 $D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a$

Let $c \in M_i$ be s.t. $a \triangleleft c$. By Lemma 3.7(e) one also has

$D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_c$

Note that by Lemma 4.7(b) we have $\forall y \geq z \ D(g(y)) = D(g(z))$ and therefore a satisfies conditions (i) and (iii) - (v) of (10) for all large enough x . Moreover, this fact is formalizable. By the (Δ_0) least number principle we can w.l.o.g. stipulate that a also satisfies (vi) and (vii) of (10). As in the proof of Lemma 4.6(b) we obtain

$$\overline{\text{Pr } (\ell(\bar{i}) = \bar{c} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{c}) \vee \ell(\bar{i}+1) = \bar{a})}$$

whence by Lemma 4.5(c) and since

$$\overline{\text{Pr } (\ell(\bar{i}+1) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}+1, y) R \bar{a}))}$$

one gets

$$\overline{\text{Pr } (\ell(\bar{i}) = \bar{c} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{c}))}$$

which along with $\forall y D(g(z)) \vdash_L D(g(y)) \text{ non } \vdash \neg \Psi_c$ yields the desired contradiction with the induction hypothesis.

(c) follows straightforwardly from (b). ■

4.9. Lemma. $\mathbb{N} \models$ "g is the identity function"

Proof. Suppose not. Then by Lemma 4.7(c) there would exist a node $a \in M_i$ s.t. $a \neq \square^N \perp$,

$$\overline{\text{Pr } (\ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a}))} \quad \text{and} \quad \forall z D(g(z)) \text{ non } \vdash_L \neg \Psi_a$$

By Lemma 4.8(b) this would imply $\ell \neq 0$ contradicting Lemma 4.3(d). ■

Next we define a mapping $^\circ$ from the set of propositional letters $\{p_i\}_{i \in \omega - \{0\}}$ to \mathfrak{D}_T by putting

$$(14) \quad p_i^\circ \equiv \ell(\bar{i}) \neq \overline{p_i}$$

and extend it in the obvious way to every modal formula in these propositional letters.

4.10. Lemma. For each $i \in \omega$ and each modal formula $A(p_1, \dots, p_i)$ there holds

$$\text{I}\Sigma_1 + \text{Pr}^N(\bar{1}) \vdash (A(p_1, \dots, p_i))^\circ \leftrightarrow \ell(\bar{i}) \neq \overline{A(p_1, \dots, p_i)}$$

Proof. We execute induction on the structure of $A(p_1, \dots, p_i)$. The case of propositional letters is handled by Lemma 4.5(f). The induction step is immediate for Boolean connectives.

We turn to \square . Reason in $\text{I}\Sigma_1 + \text{Pr}^N(\bar{1})$:

Suppose $\ell(i) \neq \square A(p_1, \dots, p_i)$. Since \neq is Δ_0 this can be

formalized: $\Pr(\overline{\ell(\bar{i})} * \overline{\square A(p_1, \dots, p_i)})$. From Lemma 4.3(b) we have $\ell \neq 0$ so with the help of Lemma 4.6(a) conclude

$$\Pr(\overline{\ell(\bar{i})} * \overline{A(p_1, \dots, p_i)})$$

Now assume $\ell(i) * \diamond A(p_1, \dots, p_i)$. By Lemma 4.3(b) $\ell \neq 0$ so from Lemmas 4.5(f) and 4.2(c) we get $\ell(i) * \square^n \perp$. There exists a node $a \in M_i$ s.t. $a * \square^n \perp$, $a * A(p_1, \dots, p_i)$ and $\ell(i) R a$ therefore with Lemma 4.6(c) one has $\neg \Pr(\overline{\ell(\bar{i})} \neq \bar{a})$ whence

$$\neg \Pr(\overline{\ell(\bar{i})} \text{ non } * \overline{A(p_1, \dots, p_i)}) \quad \blacksquare$$

4.11. Lemma. *If there holds $A(v_1, v_2, \dots) = \top$ for $A(x_1, x_2, \dots)$ a diagonalizable polynomial then*

$$\text{IS}_1 + \Pr^N(\bar{1}) \vdash (A(p_1, p_2, \dots))^\circ$$

Proof. By the definition of the sequence $\{A(m)\}_{m \in \omega}$ the equality $A(v_1, v_2, \dots) = \top$ implies that $A(k) = A(p_1, p_2, \dots)$ for a suitable $k \in \omega$ and hence by virtue of Lemma 4.4(c) there exists an $m \in \omega$ s.t. $D(m) \vdash_L A(p_1, p_2, \dots)$ so by Lemma 4.9 $D(g(m)) \vdash_L A(p_1, p_2, \dots)$ whence by Lemmas 4.3(b), 4.8(a) and 4.10

$$\text{IS}_1 + \Pr^N(\bar{1}) \vdash \text{IS}_1 + \ell \neq 0 \vdash (A(p_1, p_2, \dots))^\circ \quad \blacksquare$$

Lemma 4.11 licenses us to define a mapping $*$: $\text{rng } \nu \rightarrow \mathfrak{D}_T$ by putting

$$(15) \quad (\nu i)^* \equiv p_i^\circ \equiv \ell(\bar{i}) * \overline{p_i}$$

because if $\nu i = \nu j$ then this lemma guarantees that

$$\top \vdash \text{IS}_1 + \Pr^N(\bar{1}) \vdash p_i^\circ \leftrightarrow p_j^\circ$$

We shall show that $*$ gives rise to an embedding of \mathfrak{D} into \mathfrak{D}_T .

4.12. Proof of Theorem 4.1 concluded.

Clearly $\text{rng } *$ is r.e.

Let $A(\nu_1, \dots, \nu_i)$ hit τ in \mathfrak{D} . Then by Lemma 4.11 we have

$$\begin{aligned} T \vdash \text{I}\Sigma_1 + \text{Pr}^N(\bar{1}) \vdash (A(p_1, \dots, p_i))^\circ \\ \vdash A((\nu_1)^*, \dots, (\nu_i)^*) \end{aligned}$$

Conversely, let $T \vdash A((\nu_1)^*, \dots, (\nu_i)^*)$. If it were not the case that $A(\nu_1, \dots, \nu_i) = \tau$ then by Lemma 4.4(d) we would have $D(g(m))$ non $\vdash_{\mathbf{L}} A(p_1, \dots, p_i)$ for every $m \in \omega$. Hence for each m there would exist a node $a \in M_i$ s.t. $a \# \square^N_{\perp}$, $D(g(m))$ non $\vdash_{\mathbf{L}} \neg\Psi_a$ and $a \# \neg A(p_1, \dots, p_i)$. Since there are only finitely many nodes in M_i forcing \square^N_{\perp} we can using Lemma 4.7(a) choose a single a for all $m \in \omega$. By Lemma 4.10

$$T \vdash (A(p_1, \dots, p_i))^\circ$$

implies

$$T \vdash \ell(\bar{1}) \neq \bar{a}$$

whence by Lemma 4.8(c) $\ell \neq 0$ contrary to Lemma 4.3(d). The contradiction completes the proof of Theorem 4.1. \blacksquare

§ 5. The strong disjunction property and apt formulae

5.1. Definition. A formula A is apt if $A \text{ non } \vdash_{\mathbf{L}} \perp$ and for each pair B, C of formulas

$$A \vdash_{\mathbf{L}} \Box B \vee \Box C \Rightarrow A \vdash_{\mathbf{L}} B \text{ or } A \vdash_{\mathbf{L}} C.$$

The definition of an apt formula bears a strong semblance to the strong disjunction property in diagonalizable algebras. A tighter connection between these will be brought out in Lemma 5.16.

5.2. Lemma. If A is apt and $A \vdash_{\mathbf{L}} \Box B_0 \vee \dots \vee \Box B_n$ then for some i s.t. $0 \leq i \leq n$ we have $A \vdash_{\mathbf{L}} B_i$.

Proof. Use induction on n . For $n = 0$ and $n = 1$ the claim holds by the definition of apt formulae. Let

$$A \vdash_{\mathbf{L}} \Box B_0 \vee \dots \vee \Box B_n \vee \Box B_{n+1}$$

Then

$$A \vdash_{\mathbf{L}} \Box (\Box B_0 \vee \dots \vee \Box B_n) \vee \Box B_{n+1}$$

whence by the aptness of A

$$A \vdash_{\mathbf{L}} \Box B_0 \vee \dots \vee \Box B_n \text{ or } A \vdash_{\mathbf{L}} B_{n+1}.$$

Now apply the induction hypothesis to the former case. ■

We work now towards an effective description of apt formulae. To this end it will be convenient to think of each formula A in $F^n(\vec{p})$ as the disjunction of a set γ of atoms in $A^n(\vec{p})$. We shall introduce a preorder $Q^n(\vec{p})$ on $A^n(\vec{p})$. By analyzing the preordered set obtained by restriction of $Q^n(\vec{p})$ to a certain subset of γ it will be decided whether A is apt.

5.3. Definition. The binary relation $Q^n(\vec{p})$ on $A^n(\vec{p})$ is defined by putting

$$B Q^n(\vec{p}) C \Leftrightarrow \text{non } \vdash_{\mathbf{L}} B \rightarrow \Box C$$

5.4. Lemma. Let \mathcal{K} be a \vec{p} -model, let b and c be nodes of \mathcal{K} s.t. c is accesible from b and give the (n, \vec{p}) -characters of b and c the names B and C respectively. Then $B \mathcal{Q}^n(\vec{p}) C$.

Proof. Obvious. ■

5.5. Definition. For A a formula, we define $[A]_{n, \vec{p}}$, the (n, \vec{p}) -shadow of A , to be the conjunction of all formulas in $\mathbf{F}^n(\vec{p})$ implied by A .

5.6. Lemma. If $n \leq m$ and $\vec{p} \subseteq \vec{q}$ then for each formula A one has

$$\vdash_{\mathbf{L}} [A]_{n, \vec{p}} \leftrightarrow [[A]_{m, \vec{q}}]_{n, \vec{p}}$$

Proof. Straightforward. ■

5.7. Lemma. If $A \in \mathbf{A}^n(\vec{q})$, $m \leq n$ and $\vec{p} \subseteq \vec{q}$ then the (m, \vec{p}) -shadow of A is an element of $\mathbf{A}^m(\vec{p})$.

Proof. Trivial. ■

5.8. Lemma. For $B, C \in \mathbf{A}^{n+1}(\vec{p})$ one has $B \mathcal{Q}^{n+1}(\vec{p}) C$ iff
 (i) the (n, \vec{p}) -shadow of C is an (n, \vec{p}) -possible world of B
 and

(ii) the (n, \vec{p}) -possible worlds of C are among those of B .

Proof. (only if) is established by considering a rooted model that forces $B \wedge \diamond C$.

(if) Let $\mathcal{K}_1 \vDash B$ and $\mathcal{K}_2 \vDash C$. Graft \mathcal{K}_2 above the root of \mathcal{K}_1 . Use (i) and (ii) to see that the resulting model forces $B \wedge \diamond C$. ■

5.9. Corollary. $\mathcal{Q}^n(\vec{p})$ is transitive.

Proof. Follows immediately from Lemma 5.8. ■

5.10. Definition. If a formula A is in $\mathbf{F}^n(\vec{p})$ and $\vdash_{\mathbf{L}} A \leftrightarrow \bigvee \gamma$ with $\gamma \subseteq \mathbf{A}^n(\vec{p})$ then $\bigvee \gamma$ is called the (n, \vec{p}) -normal form

of A . A formula A in $F^n(\vec{p})$ is called (n, \vec{p}) -trimmed if letting $\mathbb{W} \gamma$ be its (n, \vec{p}) -normal form one has $A \vdash_L \neg G$ for no $G \in \gamma$. Equivalently, A is (n, \vec{p}) -trimmed if and only if $\vdash_L A \leftrightarrow [A]_{n, \vec{p}}$.

5.11. Lemma. *A formula of the form $\Box^+ A$ is (n, \vec{p}) -trimmed whenever $\Box^+ A \in F^n(\vec{p})$.*

Proof. This is trivial since $\Box^+ A \vdash_L \neg G$ implies $\vdash_L \Box^+ A \rightarrow \neg G$. ■

5.12. Lemma. *To each formula A in $F^n(\vec{p})$ there corresponds an (n, \vec{p}) -trimmed formula B s.t. $A \vdash_L B$ and $B \vdash_L A$. The formula B with these properties is unique up to L -equivalence.*

Proof. Take B to be the conjunction of all such formulas C in $F^n(\vec{p})$ that $A \vdash_L C$. Let $\mathbb{W} \gamma$ be the (n, \vec{p}) -normal form of B . If $G \in \mathbb{A}^n(\vec{p})$ and $B \vdash_L \neg G$ then $\neg G$ is a conjunct of B and hence $G \text{ non } \in \gamma$.

We leave uniqueness to the reader. ■

5.13. Definition. Let $A \in F^n(\vec{p})$ be (n, \vec{p}) -trimmed and let $\mathbb{W} \gamma$ be the (n, \vec{p}) -normal form of A . A formula $E \in \gamma$ is called an (n, \vec{p}) -bottom of A if $E \mathcal{Q}^n(\vec{p}) C$ for each $C \in \gamma$. In this case A is said to be (n, \vec{p}) -bottomed.

The following lemma gives us a convenient characterization of apt formulae along with an algorithm for deciding aptness.

5.14. Lemma. *Let A be a formula in $F^n(\vec{p})$ and let B be the (n, \vec{p}) -trimmed formula which corresponds to A by Lemma 5.12. Then the following are equivalent:*

- (i) A is apt;
- (ii) B is (n, \vec{p}) -bottomed;
- (iii) A is irrefutable in L and for each pair $\mathcal{K}_1, \mathcal{K}_2$ of finite rooted models in which A holds there exists a model \mathcal{H} s.t. $\mathcal{H} \models A$ and \mathcal{K}_1 and \mathcal{K}_2 are isomorphic to proper cones of \mathcal{H} .

Proof. (i) \Rightarrow (ii) Let $\mathbb{W} \gamma$ be the (n, \vec{p}) -normal form of B . An (n, \vec{p}) -bottom E of B exists for otherwise we would have $A \vdash_L$

$\bigvee \{ \Box G \mid G \in \gamma \}$ which by the aptness of A and by Lemma 5.2 implies $A \vdash_{\mathbf{L}} \Box G$ for some $G \in \gamma$ contradicting the (n, \vec{p}) -trimmedness of B .

(ii) \Rightarrow (iii) Let $E \in \mathbf{F}^n(\vec{p})$ be an (n, \vec{p}) -bottom of B and let \mathcal{H} be a rooted \vec{p} -model forcing $E \wedge \Box^+ A$ (and therefore $\Box^+ B$). For $\vec{q} \supseteq \vec{p}$ take a pair of \vec{q} -models in which A holds and extend to \vec{q} the forcing relation at the nodes of \mathcal{H} in an arbitrary way. Next graft the chosen pair of \vec{q} -models above the root of \mathcal{H} . The resulting model will still force $E \wedge \Box^+ B$ (this can be seen through Lemma 5.8) and hence will also force $\Box^+ A$.

(iii) \Rightarrow (i) Let $A \vdash_{\mathbf{L}} \Box C \vee \Box D$. If it were the case that neither $A \vdash_{\mathbf{L}} C$ nor $A \vdash_{\mathbf{L}} D$ then there would exist two finite rooted models \mathcal{K}_1 and \mathcal{K}_2 in which A holds s.t. $\mathcal{K}_1 \text{ non } * C$ and $\mathcal{K}_2 \text{ non } * D$. Taking the model \mathcal{H} which corresponds to \mathcal{K}_1 and \mathcal{K}_2 by the assumption we would have $\mathcal{H} \text{ non } * \Box C \vee \Box D$ and yet $\mathcal{H} \vDash A$. A contradiction. ■

5.15. Corollary. *Aptness is decidable.*

Proof. Follows from (i) \Leftrightarrow (ii) of Lemma 5.14. ■

Consider the following property of modal schemes A :

for each pair B, C of modal formulas
 $\vdash_{\mathbf{L}+A} \Box B \vee \Box C$ implies $\vdash_{\mathbf{L}+A} B$ or $\vdash_{\mathbf{L}+A} C$

Chagrov [15] shows it to be undecidable in contrast to Corollary 5.15.

5.16. Lemma. *Let \mathfrak{D} be a diagonalizable algebra generated by x_0, x_1, \dots . Then \mathfrak{D} enjoys the strong disjunction property iff for each modal formula (= diagonalizable polynomial) B s.t. $B(x_0, x_1, \dots) = \top$ there exists an apt formula A s.t. $A(x_0, x_1, \dots) = \top$ and $\vdash_{\mathbf{L}} A \rightarrow B$.*

Proof. (if) Suppose that for elements c and d of \mathfrak{D} we have $\Box c \vee \Box d = \top$. Let $c = C(x_0, x_1, \dots)$ and $d = D(x_0, x_1, \dots)$ so that

$$\Box C(x_0, x_1, \dots) \vee \Box D(x_0, x_1, \dots) = \top$$

By the assumptions on \mathfrak{D} there exists an apt formula A s.t. $A \vdash_{\mathbf{L}} \Box C \vee \Box D$ and $A(x_0, x_1, \dots) = \tau$. We therefore have that $A \vdash_{\mathbf{L}} C$ or $A \vdash_{\mathbf{L}} D$ whence $c = \tau$ or $d = \tau$. This proves the strong disjunction property.

(only if) Let $B \in \mathbf{F}^n(\vec{p})$ be s.t. $B(x_0, x_1, \dots) = \tau$ and let A be the conjunction of all such formulas C in $\mathbf{F}^n(\vec{p})$ that $C(x_0, x_1, \dots) = \tau$ (there are, up to \mathbf{L} -equivalence, only finitely many). Clearly $\vdash_{\mathbf{L}} A \rightarrow B$. Suppose $\mathbb{W} \gamma$ is the (n, \vec{p}) -normal form of A . Note that A is (n, \vec{p}) -trimmed because if $A \vdash_{\mathbf{L}} \neg C$ for some $C \in \mathbf{A}^n(\vec{p})$ then $\neg C(x_0, x_1, \dots) = \tau$ whence $\vdash_{\mathbf{L}} A \rightarrow \neg C$ and so $C \text{ non} \in \gamma$. If there were no (n, \vec{p}) -bottom to A then we would have $A \vdash_{\mathbf{L}} \mathbb{W} \{ \Box G \mid G \in \gamma \}$ and hence

$$\mathbb{W} \{ \Box G(x_0, x_1, \dots) \mid G \in \gamma \} = \tau$$

whence by the strong disjunction property we have $\neg G(x_0, x_1, \dots) = \tau$ for some $G \in \gamma$. Therefore $\vdash_{\mathbf{L}} A \rightarrow \neg G$ by the choice of A and so $G \text{ non} \in \gamma$. The contradiction shows that A is (n, \vec{p}) -bottomed and therefore apt by Lemma 5.14. \blacksquare

5.17. Definition. An element a of a diagonalizable algebra \mathfrak{D} with the strong disjunction property is *admissible* if $a \rightarrow \Box b = \tau$ implies $b = \tau$ for each element b of \mathfrak{D} .

One of the intended uses of the notion of an admissible element will be based on the fact that if T is a sound enough theory then the elements of \mathfrak{D}_T that correspond to true sentences of low arithmetical complexity have to be admissible.

In fact, if T is a Σ_1 -sound theory then a sentence φ is an admissible element of \mathfrak{D}_T if and only if $T + \varphi$ is Σ_1 -sound.

5.18. Lemma. Let \mathfrak{D} be a diagonalizable algebra with the strong disjunction property.

- (a) The element τ of \mathfrak{D} is admissible.
- (b) If an element a of \mathfrak{D} is admissible, $b \in \mathfrak{D}$ and $a \rightarrow b = \tau$ then b is admissible.
- (c) If an element a of \mathfrak{D} is admissible and b is an arbitra-

ry element of \mathfrak{D} then at least one of the elements $a \wedge b$ and $a \wedge \neg b$ is admissible.

Proof. (a) and (b) are straightforward.

(c) Suppose neither $a \wedge b$ nor $a \wedge \neg b$ is admissible. Then there exist elements c and d of \mathfrak{D} distinct from τ s.t.

$$a \wedge b \rightarrow \Box c = a \wedge \neg b \rightarrow \Box d = \tau$$

But then $a \rightarrow \Box c \vee \Box d = \tau$ whence $a \rightarrow \Box(\Box c \vee \Box d) = \tau$. Since a is admissible we have that $\Box c \vee \Box d = \tau$ so by the strong disjunction property $c = \tau$ or $d = \tau$ which contradicts the assumptions and therefore proves the lemma. ■

5.19. Definition. Let A be an apt formula. A formula B is said to *support* a formula A if for each formula C one has

$$A \vdash_{\mathbf{L}} B \rightarrow \Box C \Rightarrow A \vdash_{\mathbf{L}} C.$$

The relation "to support" parallels the notion of an admissible element of a diagonalizable algebra with the strong disjunction property in the same way that aptness parallels the strong disjunction property itself.

5.20. Lemma. If B supports an apt formula A and $A \vdash_{\mathbf{L}} B \rightarrow (\Box C_0 \vee \dots \vee \Box C_n)$ then for some i s.t. $0 \leq i \leq n$ we have $A \vdash_{\mathbf{L}} C_i$.

Proof. Analogous to that of Lemma 5.2. ■

5.21. Lemma. Let A be an apt formula.

(a) τ supports A .

(b) If B supports A and $\vdash_{\mathbf{L}} B \rightarrow C$ then C supports A .

(c) Let B support A . Then for each formula C at least one of the formulas $B \wedge C$ and $B \wedge \neg C$ supports A .

Proof. Similiar to the proof of Lemma 5.18. ■

To decide the relation " A supports B " we have to take a little bit closer look at the (n, \vec{p}) -bottoms of B than we did

when deciding aptness.

Our investigation of the relation "A supports B" will henceforth be restricted to \square -free formulas A. This involves no loss of generality because A supports B if and only if q supports $B \wedge (q \leftrightarrow A)$ where q is a new propositional letter.

5.22. Lemma. Let $A \in F^n(\vec{p})$, $\alpha \in F^0$ and let B be the (n, \vec{p}) -trimmed formula which corresponds to A by Lemma 5.12. Suppose A is apt. Then the following are equivalent:

- (i) α supports A;
- (ii) For some (n, \vec{p}) -bottom E of B the formula $E \wedge \alpha$ is irrefutable in L (which is the same as to say that α is consistent with the real world part of E in propositional logic);
- (iii) For each finite rooted model K in which A holds there exists a rooted model \mathcal{K} s.t. $\mathcal{K} \vDash \alpha \wedge \square^+ A$ and K is isomorphic to a proper cone of \mathcal{K} .

Proof. (i) \Rightarrow (ii) Let $\mathbb{W} \gamma$ be the (n, \vec{p}) -normal form of B. Since B is (n, \vec{p}) -trimmed $A \vDash_L \neg G$ holds for no $G \in \gamma$. Therefore by Lemma 5.20

$$A \text{ non } \vDash_L \alpha \rightarrow \mathbb{W} \{ \square \neg G \mid G \in \gamma \}$$

This implies that there exists a rooted model K forcing $\square^+ A \wedge \alpha \wedge \mathbb{M} \diamond \gamma$. For E the (n, \vec{p}) -character of K we have that $E \wedge \alpha$ is irrefutable in L and $E \vDash Q^n(\vec{p}) G$, all $G \in \gamma$.

(ii) \Rightarrow (iii) Consider a \vec{p} -model \mathcal{K} of $E \wedge \square^+ A$. Since $E \wedge \alpha$ is irrefutable in L there exists such an extension of the forcing relation at the nodes of \mathcal{K} that the resulting model \mathcal{K} forces $\alpha \wedge E \wedge \square^+ A$. If one grafts a model in which A (and hence $\mathbb{W} \gamma$) holds above the root of K and extends the forcing relation to new propositional letters if necessary then Lemma 5.8 guarantees that the result forces $\alpha \wedge E \wedge \square^+ A$.

(iii) \Rightarrow (i) Suppose $A \text{ non } \vDash_L D$. Then there is a finite rooted model K forcing $\neg D \wedge \square^+ A$. From (iii) we obtain a model \mathcal{K} forcing $\alpha \wedge \square^+ A$ of which K is a proper cone. The model \mathcal{K} testifies $A \text{ non } \vDash_L \alpha \rightarrow \square D$. Thus we have shown that α supports A. ■

5.23. Corollary. *The binary relation "A supports B" is decidable.*

Proof. Immediate from Corollary 5.15 and (i) \Leftrightarrow (ii) of Lemma 5.22. ■

5.24. Lemma. *Let \mathfrak{D} be a diagonalizable algebra with the strong disjunction property generated by x_0, x_1, \dots . Suppose $\alpha \in \mathbf{F}^0$. An element $a = \alpha(x_0, x_1, \dots)$ of \mathfrak{D} is admissible iff for each modal formula (= diagonalizable polynomial) B s.t. $B(x_0, x_1, \dots) = \top$ there exists an apt formula C supported by α s.t. $C(x_0, x_1, \dots) = \top$ and $\vdash_{\mathbf{L}} C \rightarrow B$.*

Proof. (if) Suppose $a \rightarrow \Box b = \top$ for b an element of \mathfrak{D} . Let $b = B(x_0, x_1, \dots)$. There exists an apt formula C s.t. $C(x_0, x_1, \dots) = \top$, α supports C and $C \vdash_{\mathbf{L}} \alpha \rightarrow \Box B$. By the definition of the relation "to support" we have $C \vdash_{\mathbf{L}} B$ therefore $b = \top$ and hence a is admissible.

(only if) Let $B(x_0, x_1, \dots) = \top$ and let n and \vec{p} be s.t. $B \in \mathbf{F}^n(\vec{p})$. Call C the conjunction of all such formulas D in $\mathbf{F}^n(\vec{p})$ that $D(x_0, x_1, \dots) = \top$. Clearly $\vdash_{\mathbf{L}} C \rightarrow B$. As in Lemma 5.16 we can show that C is apt and (n, \vec{p}) -trimmed. Let $\mathbb{W} \gamma$ be the (n, \vec{p}) -normal form of C . If C had no (n, \vec{p}) -bottom E s.t. $\text{non } \vdash_{\mathbf{L}} \neg(E \wedge \alpha)$ then we would have

$$\begin{aligned} C \vdash_{\mathbf{L}} \alpha &\rightarrow \mathbb{W} \{ \Box G \mid G \in \gamma \} \\ &\rightarrow \Box \mathbb{W} \{ \Box G \mid G \in \gamma \} \end{aligned}$$

and hence

$$a \rightarrow \Box \mathbb{W} \{ \Box G(x_0, x_1, \dots) \mid G \in \gamma \} = \top$$

Since a is admissible and \mathfrak{D} possesses the strong disjunction property, on the strength of Lemma 5.20 there exists a $G \in \gamma$ s.t. $\neg G(x_0, x_1, \dots) = \top$ thus contradicting the choice of C . Therefore an (n, \vec{p}) -bottom E of C s.t. $\text{non } \vdash_{\mathbf{L}} \neg(E \wedge \alpha)$ exists and hence by Lemma 5.22 a supports C q.e.d. ■

The next goal is to show that the taking of shadows of certain formulas preserves aptness and does not diminish the number

of supporting formulas in F^0 .

5.25. Lemma. Suppose $m \leq n$ and $\vec{q} \subseteq \vec{p}$.

(a) Let $A \in F^n(\vec{p})$. If A is (n, \vec{p}) -trimmed then $[A]_{m, \vec{q}}$ is (m, \vec{q}) -trimmed.

(b) Suppose $A \in A^{n+1}(\vec{p})$ and let $A \equiv \alpha \wedge \square \mathbb{W} \gamma \wedge \mathbb{M} \diamond \gamma$ where $\alpha \in A^0(\vec{p})$ and $\gamma \subseteq A^n(\vec{p})$. Then

$$\vdash_L [A]_{m+1, \vec{q}} \leftrightarrow [\alpha]_{0, \vec{q}} \wedge \square \mathbb{W} [\gamma]_{m, \vec{q}} \wedge \mathbb{M} \diamond [\gamma]_{m, \vec{q}}$$

where $[\gamma]_{m, \vec{q}}$ stands for $\{ [G]_{m, \vec{q}} \mid G \in \gamma \}$.

(c) Let $A, B \in A^n(\vec{p})$. Then $A \mathcal{Q}^n(\vec{p}) B$ implies $[A]_{m, \vec{q}} \mathcal{Q}^m(\vec{q}) [B]_{m, \vec{q}}$.

(d) Let $\mathbb{W} \gamma$ be the (n, \vec{p}) -normal form of $A \in F^n(\vec{p})$. Then

$$\vdash_L [\mathbb{W} \gamma]_{m, \vec{q}} \leftrightarrow \mathbb{W} [\gamma]_{m, \vec{q}}$$

(e) Suppose $A \in F^n(\vec{p})$. If A is (n, \vec{p}) -trimmed and apt then $[A]_{m, \vec{q}}$ is apt and if in addition a formula $\alpha \in F^0$ supports A then it also supports $[A]_{m, \vec{q}}$.

Proof. (a) Let $\mathbb{W} \delta$ and $\mathbb{W} \gamma$ be the (n, \vec{p}) - and the (m, \vec{q}) -normal forms of A and of $[A]_{m, \vec{q}}$ respectively. If $[A]_{m, \vec{q}}$ were not (m, \vec{q}) -trimmed then $[A]_{m, \vec{q}} \vdash_L \neg G$ for some $G \in \gamma$. There clearly exists a formula $D \in \delta$ s.t. $\vdash_L D \rightarrow G$ and so

$$A \vdash_L [A]_{m, \vec{q}} \vdash_L \neg G \vdash_L \neg D$$

contrary to the (n, \vec{p}) -trimmedness of A .

(b) Clearly the r.h.s. is an element of $F^{m+1}(\vec{q})$. By Lemmas 2.2 and 5.7 we only have to show

$$\vdash_L A \rightarrow [\alpha]_{0, \vec{q}} \wedge \square \mathbb{W} [\gamma]_{m, \vec{q}} \wedge \mathbb{M} \diamond [\gamma]_{m, \vec{q}}$$

But this is clear since

$$\vdash_L \alpha \rightarrow [\alpha]_{0, \vec{q}}$$

and for each $G \in \gamma$

$$\vdash_{\mathbf{L}} G \rightarrow [G]_{m, \vec{q}}$$

hence

$$\vdash_{\mathbf{L}} \Box W \gamma \rightarrow \Box W [\gamma]_{m, \vec{q}} \quad \text{and} \quad \vdash_{\mathbf{L}} \mathbb{M} \diamond \gamma \rightarrow \mathbb{M} \diamond [\gamma]_{m, \vec{q}}$$

This completes the proof.

(c) By Lemma 5.7 both $[A]_{m, \vec{q}}$ and $[B]_{m, \vec{q}}$ are in $\mathbf{A}^m(\vec{q})$. If $m = 0$ the claim is obvious so let $n > 0$ and $m > 0$.

From Lemma 5.8 we have that all the $(n-1, \vec{p})$ -possible worlds of B as well as $[B]_{n-1, \vec{p}}$ are among the $(n-1, \vec{p})$ -possible worlds of A . By (b) the $(m-1, \vec{q})$ -possible worlds of $[A]_{m, \vec{q}}$ and $[B]_{m, \vec{q}}$ are precisely the $(m-1, \vec{q})$ -shadows of the $(n-1, \vec{p})$ -possible worlds of A and B respectively and $[B]_{m-1, \vec{q}}$ is an $(m-1, \vec{q})$ -possible world of $[A]_{m, \vec{q}}$. Therefore by Lemma 5.8 $[A]_{m, \vec{q}} \mathbf{Q}^m(\vec{q}) [B]_{m, \vec{q}}$ q.e.d.

(d) Straightforward.

(e) By (a) $[A]_{m, \vec{q}}$ is (m, \vec{q}) -trimmed. With (c) and (d) it is seen that $[A]_{m, \vec{q}}$ is (m, \vec{q}) -bottomed. If α supports A then by Lemma 5.22 there exists an (n, \vec{p}) -bottom E of A s.t. $\text{non } \vdash_{\mathbf{L}} \neg(E \wedge \alpha)$. By (c) and (d) $[E]_{m, \vec{q}}$ is an (m, \vec{q}) -bottom of $[A]_{m, \vec{q}}$ and clearly $\text{non } \vdash_{\mathbf{L}} \neg([E]_{m, \vec{q}} \wedge \alpha)$. Hence by Lemma 5.22 $[A]_{m, \vec{q}}$ is supported by α . ■

5.26. Lemma. *A formula B is apt iff $\Box^+ B$ is apt. A supports B iff A supports $\Box^+ B$.*

Proof. This is clear because for any formula C we have $\vdash_{\mathbf{L}} C$ if and only if $\Box^+ B \vdash_{\mathbf{L}} C$. ■

5.27. Corollary. *Let A be apt. Then $[\Box^+ A]_{m, \vec{q}}$ is apt and if a formula $\alpha \in \mathbf{F}^0$ supports A then it also supports $[\Box^+ A]_{m, \vec{q}}$.*

Proof. Immediate from Lemmas 5.11, 5.25(e) and 5.26. ■

Finally we prove a lemma which will enable us to construct apt formulae with a number of meritorious properties.

5.28. Lemma. Let $\vec{p} \subseteq \vec{q}$, $A \in F(\vec{q})$ and $B \in F(\vec{p})$. Suppose that B is apt and that $\Box^+ A$ is \vec{p} -conservative over $\Box^+ B$. Then there exists an apt formula $C \in F(\vec{q})$ s.t. $C \vdash_{\mathbf{L}} A$ and $\Box^+ C$ is \vec{p} -conservative over $\Box^+ B$. Moreover C can be chosen to be supported by every formula in $F^0(\vec{p})$ that supports B .

Proof. To keep notation at bay we shall only consider the case when B supports just two elements of $F^0(\vec{p})$, namely α and β . It will easily be seen how to generalize the proof to any larger number of formulas.

Let $A \in F^N(\vec{q})$ and let $\mathbb{W} \delta$ be the (n, \vec{q}) -normal form of the (n, \vec{q}) -trimmed formula corresponding to A by Lemma 5.12. Put

$$\gamma_{\alpha, \beta} = \{ D \in \delta \mid \text{there exist formulas } E_{\alpha}, E_{\beta} \in \delta \text{ s.t.} \\ E_{\alpha} Q^n(\vec{q}) E_{\beta} Q^n(\vec{q}) E_{\alpha} Q^n(\vec{q}) D \\ \text{and neither } \alpha \wedge E_{\alpha} \text{ nor } \beta \wedge E_{\beta} \text{ is refutable in } \mathbf{L} \}$$

Let $\Box_D G$ denote the formula $\Box(D \rightarrow G)$. Dually, $\Diamond_D G$ denotes $\Diamond(D \wedge G)$. Note that since B is apt and supported by α and β we have $B \vdash_{\mathbf{L}} F$ whenever $B \vdash_{\mathbf{L}} \Box_{\alpha} \Box F$ or $B \vdash_{\mathbf{L}} \Box_{\beta} \Box F$.

Claim 1. For each formula $G \in \delta - \gamma_{\alpha, \beta}$ there exists an $N \in \omega$ s.t. $\mathbb{W} \delta \vdash_{\mathbf{L}} (\Box_{\alpha} \Box_{\beta})^N \Box G$.

For if this were not so then there would exist models of $\Box^+ \mathbb{W} \delta \wedge (\Diamond_{\alpha} \Diamond_{\beta})^N \Diamond G$ for arbitrarily large $N \in \omega$. Hence by Lemma 5.4 there would exist arbitrarily long sequences $D_1, E_1, \dots, D_N, E_N$ of elements of δ s.t. $D_i Q^n(\vec{q}) E_i Q^n(\vec{q}) D_{i+1} Q^n(\vec{q}) G$, $\vdash_{\mathbf{L}} D_i \rightarrow \alpha$ and $\vdash_{\mathbf{L}} E_i \rightarrow \beta$ for each i satisfying $1 \leq i < N$. But since $Q^n(\vec{q})$ is transitive and δ is finite this would imply the existence of formulas $D, E \in \delta$ s.t. $D Q^n(\vec{q}) E Q^n(\vec{q}) D Q^n(\vec{q}) G$, $\vdash_{\mathbf{L}} D \rightarrow \alpha$ and $\vdash_{\mathbf{L}} E \rightarrow \beta$ and this puts G in $\gamma_{\alpha, \beta}$ and therefore contradicts the assumption on G . Thus Claim 1 is proved.

Claim 2. $\Box^+ \mathbb{W} \gamma_{\alpha, \beta}$ is \vec{p} -conservative over $\Box^+ B$.

Suppose $F \in F(\vec{p})$ and $\mathbb{W} \gamma_{\alpha, \beta} \vdash_{\mathbf{L}} F$. Since $\delta - \gamma_{\alpha, \beta}$ is

finite Claim 1 ensures the existence of a single $N \in \omega$ s.t.
 $\mathbb{W} \delta \vdash_{\mathbf{L}} (\Box_{\alpha} \Box_{\beta})^N \Box G$ whenever $G \in \delta - \gamma_{\alpha, \beta}$.

We have

$$\begin{aligned} \mathbb{W} \delta \vdash_{\mathbf{L}} (\Box_{\alpha} \Box_{\beta})^N \Box \bigwedge \{ \neg G \mid G \in \delta - \gamma_{\alpha, \beta} \} \\ \vdash_{\mathbf{L}} (\Box_{\alpha} \Box_{\beta})^N \Box \mathbb{W} \delta \\ \vdash_{\mathbf{L}} (\Box_{\alpha} \Box_{\beta})^N \Box \mathbb{W} \gamma_{\alpha, \beta} \\ \vdash_{\mathbf{L}} (\Box_{\alpha} \Box_{\beta})^N \Box F \end{aligned}$$

But since $\Box^+ \mathbb{W} \delta$ is \vec{p} -conservative over $\Box^+ B$ we also have $B \vdash_{\mathbf{L}} (\Box_{\alpha} \Box_{\beta})^N \Box F$ whence $B \vdash_{\mathbf{L}} F$ for B is apt and α and β support B . This proves Claim 2.

Next we let

$$\begin{aligned} \varepsilon_{\alpha, \beta} = \{ D \in \delta \mid \text{there exist formulas } E_{\alpha}, E_{\beta} \in \delta \text{ s.t.} \\ D \ Q^n(\vec{q}) \ E_{\alpha} \ Q^n(\vec{q}) \ E_{\beta} \ Q^n(\vec{q}) \ D \\ \text{and neither } \alpha \wedge E_{\alpha} \text{ nor } \beta \wedge E_{\beta} \text{ is refutable in } \mathbf{L} \} \end{aligned}$$

and for each $E \in \varepsilon_{\alpha, \beta}$

$$\gamma_E = \{ D \in \delta \mid E \ Q^n(\vec{q}) \ D \}$$

Claim 3. $\mathbb{W} \gamma_{\alpha, \beta} \vdash_{\mathbf{L}} \mathbb{W} \{ \Box^+ \mathbb{W} \gamma_E \mid E \in \varepsilon_{\alpha, \beta} \}$.

Let a rooted \vec{q} -model \mathcal{K} be s.t. $\mathcal{K} \models \mathbb{W} \gamma_{\alpha, \beta}$ and let D be the (n, \vec{q}) -character of \mathcal{K} . Since $D \in \gamma_{\alpha, \beta}$ there exists a formula $E \in \varepsilon_{\alpha, \beta}$ s.t. $E \ Q^n(\vec{q}) \ D$. Consider an arbitrary node a of \mathcal{K} . For G the (n, \vec{q}) -character of $\mathcal{K}[a]$ one has $D \ Q^n(\vec{q}) \ G$ or $D = G$ by Lemma 5.4. Therefore by Corollary 5.9 $E \ Q^n(\vec{q}) \ G$ hence $G \in \gamma_E$. So $\mathcal{K} \models \mathbb{W} \gamma_E$ and $\mathcal{K} \models \mathbb{W} \{ \Box^+ \mathbb{W} \gamma_E \mid E \in \varepsilon_{\alpha, \beta} \}$ q.e.d.

Claim 4. For each $E \in \varepsilon_{\alpha, \beta}$ the formula $\mathbb{W} \gamma_E$ is (n, \vec{q}) -trimmed.

Take a formula $G \in \gamma_E$. Let \mathcal{H} be a rooted \vec{q} -model forcing $E \wedge \Box^+ \mathbb{W} \delta$ (such a model exists because $\mathbb{W} \delta$ is (n, \vec{q}) -trimmed). By Lemma 5.8 $\mathbb{W} \gamma_E$ holds in \mathcal{H} . Next pick a rooted \vec{q} -model \mathcal{K} s.t. $\mathcal{K} \models G \wedge \Box^+ \mathbb{W} \delta$. Since $E \ Q^n(\vec{q}) \ G$ we have that $\mathcal{K} \models \mathbb{W} \gamma_E$. Now graft \mathcal{K} above the root of \mathcal{H} . By Lemma 5.8 it is easily seen that the resulting model forces $E \wedge \Box^+ \mathbb{W} \gamma_E$ and it clearly forces $\diamond G$. So $\mathbb{W} \gamma_E$ non $\vdash_{\mathbf{L}} \neg G$. Thus $\mathbb{W} \gamma_E$ is (n, \vec{q}) -trimmed.

Claim 5. There exists an $E \in \varepsilon_{\alpha, \beta}$ s.t. $\Box^+ \mathbb{W} \gamma_E$ is \vec{p} -conservative over $\Box^+ B$.

Deny this. For each $E \in \varepsilon_{\alpha, \beta}$ let $G_E \in F(\vec{p})$ be a formula s.t. $\mathbb{W} \gamma_E \vdash_L G_E$ and B non $\vdash_L G_E$. Now use Claim 3:

$$\begin{aligned} \mathbb{W} \gamma_{\alpha, \beta} &\vdash_L \mathbb{W} \{ \Box^+ \mathbb{W} \gamma_E \mid E \in \varepsilon_{\alpha, \beta} \} \\ &\vdash_L \mathbb{W} \{ \Box^+ G_E \mid E \in \varepsilon_{\alpha, \beta} \} \\ &\vdash_L \mathbb{W} \{ \Box G_E \mid E \in \varepsilon_{\alpha, \beta} \} \end{aligned}$$

By Claim 2 this gives

$$B \vdash_L \mathbb{W} \{ \Box G_E \mid E \in \varepsilon_{\alpha, \beta} \}$$

whence for some $E \in \varepsilon_{\alpha, \beta}$ we have $B \vdash_L G_E$ by Lemma 5.2 for B is apt. This contradiction settles Claim 5.

Now note that we have actually proved that at least one of the formulas $\mathbb{W} \gamma_E$ meets the requirements on C in the statement of the present lemma. For $\mathbb{W} \gamma_E \vdash_L \mathbb{W} \gamma_{\alpha, \beta} \vdash_L \mathbb{W} \delta \vdash_L A$ because $\gamma_E \subseteq \gamma_{\alpha, \beta} \subseteq \delta$; for each $E \in \varepsilon_{\alpha, \beta}$ the formula $\mathbb{W} \gamma_E$ is clearly (n, \vec{q}) -bottomed by E and hence apt by Claim 4 and Lemma 5.14 and at least one of the formulas $\Box^+ \mathbb{W} \gamma_E$ is \vec{p} -conservative over $\Box^+ B$. Finally, $\mathbb{W} \gamma_E$ is clearly supported by both α and β for each $E \in \varepsilon_{\alpha, \beta}$ ■

As usual we shall not hesitate to use appropriate versions of certain results of this § formalized in $\text{I}\Sigma_1$. Thus Lemma 5.28 is meant to be applied to nonstandard formulae.

§ 6. Σ_1 -ill theories of infinite credibility extent

6.1. Theorem. *Let T be a Σ_1 -ill theory of infinite credibility extent. A denumerable diagonalizable algebra \mathfrak{D} is isomorphic to an r.e. subalgebra of \mathfrak{D}_T iff*

- (i) \mathfrak{D} is positive and
- (ii) the height of \mathfrak{D} is infinite.

Many details of the proof of this theorem are similiar to those of that of Theorem 4.1. Therefore the proofs of certain lemmas are omitted whenever they exhibit no considerable deviation from the proofs of corresponding lemmas in § 4. Also the conventions of § 4 on formalized modal logic and Kripke models are still valid.

First we define the Solovay function for M_0 along with its limit value:

- (1) $h_0(0) = 0$
 - (2)
- $$h_0(x+1) = a \quad \text{if (i) } a \in M_0 ;$$
- $$\quad \quad \quad \text{(ii) } h_0(x) R a \quad \text{and}$$
- $$\quad \quad \quad \text{(iii) } \overline{\text{Prf}(x, \ell_0 = \bar{a} \rightarrow \exists y (h_0(x) R h_0(y) R \bar{a}))}$$
- $$= h_0(x) \quad \text{if no } a \text{ satisfying (i) - (iii) exists}$$
- (3) $\ell_0 = \lim_{x \rightarrow \infty} h_0(x) \quad \text{if } h_0 \text{ reaches a limit}$
 $= 0 \quad \text{otherwise}$

6.2. Lemma ($I\Sigma_1$).

- (a) $\forall x \forall y (x \leq y \rightarrow h_0(x) = h_0(y) \vee h_0(x) R h_0(y))$
- (b) $\ell_0 = \lim_{x \rightarrow \infty} h_0(x)$
- (c) $\forall x (h_0(x) = \ell_0 \vee h_0(x) R \ell_0)$
- (d) $\forall a \in M_0 (\ell_0 R a \rightarrow \neg \overline{\text{Pr}(\ell_0 = \bar{a} \rightarrow \exists y (\ell_0 R h_0(y) R \bar{a}))})$
- (e) $\forall a \in M_0 (\ell_0 R a \rightarrow \neg \overline{\text{Pr}(\ell_0 \neq \bar{a})})$
- (f) $\ell_0 \neq 0 \rightarrow \exists a \in M_0 \cup \{0\} \exists x (h_0(x) = a \wedge$

$$\wedge \text{Prf} (x, \overline{\ell_0 = \bar{\ell}_0} \rightarrow \exists y (\bar{a} R h_0(y) R \bar{\ell}_0))$$

$$(g) \ell_0 \neq 0 \rightarrow \text{Pr} (\overline{\ell_0 \neq \bar{\ell}_0})$$

$$(h) \ell_0 \neq 0 \rightarrow \text{Pr} (\overline{\bar{\ell}_0 R \ell_0})$$

Proof. The only twist new to § 4 occurs in (b). In the present situation we have to apply the least number principle on n to the formula $\exists x h_0(x) \neq \square^n 1$. ■

6.3. Lemma.

$$(a) I\Sigma_1 \vdash \forall n (n \neq 0 \rightarrow \text{Pr}^n (\bar{1}) \leftrightarrow \ell_0 \neq \square^n 1)$$

$$(b) \text{For no } a \in M_0 \text{ do we have } T \vdash \ell_0 \neq \bar{a}.$$

$$(c) \mathbb{N} \neq \ell_0 = 0$$

The theory T will prove $\ell_0 \neq 0$ as likely as not. In the former case the constructions below could be considerably simplified along the lines of § 4.

As in § 4, let $\nu: \omega - \{0\} \rightarrow \mathcal{D}$ be a positive numeration of \mathcal{D} and let $\{A(m)\}_{m \in \omega}$ be a Δ_0 enumeration of diagonalizable polynomials in propositional letters $\{p_i\}_{i \in \omega - \{0\}}$ that turn to τ of \mathcal{D} on substituting νi for p_i . We construct a better behaved and a slightly longer Δ_0 sequence $\{D(m)\}_{m \in \omega \cdot 2}$. The domain of the Δ_0 function $k(\cdot)$ is however just ω .

$$(4) D(0) = \tau$$

$$(5) k(0) = 0$$

$$(6) D(x+1) = A(x) \quad \text{if (i) } A(x) \vdash_{\mathbf{L}} D(x) ;$$

$$\quad \quad \quad \text{(ii) } A(x) \vdash_{\mathbf{L}} A(k(x)) \quad \text{and}$$

$$\quad \quad \quad \text{(iii) } \square^+ A(x) \text{ is conservative}$$

$$= D(x) \quad \text{otherwise}$$

$$(7) k(x+1) = k(x) + 1 \quad \text{if } D(x+1) \vdash_{\mathbf{L}} A(k(x))$$

$$= k(x) \quad \text{otherwise}$$

$$(8) \text{Let } D(\omega+x) \text{ be the formula manufactured by Lemma 5.28 s.t.}$$

$$(i) D(\omega+x) \vdash_{\mathbf{L}} D(x) ;$$

$$(ii) D(\omega+x) \text{ is apt and}$$

(iii) $\Box^+ D(\omega+x)$ is conservative

6.4. Lemma.

(a) $I\Sigma_1 \vdash \forall x \in \omega \forall y \in \omega (x \leq y \rightarrow D(y) \vdash_{\mathbf{L}} D(x))$

(b) $I\Sigma_1 \vdash \forall x \in \omega D(\omega+x) \vdash_{\mathbf{L}} D(x)$

(c) $I\Sigma_1 \vdash \forall x \in \omega \cdot 2 \text{ "}\Box^+ D(x) \text{ is conservative"}$

(d) $I\Sigma_1 \vdash \forall x \in \omega \text{ "}D(\omega+x) \text{ is apt"}$

(e) For each $y \in \omega$ there exists an $x \in \omega$ s.t. $D(x) \vdash_{\mathbf{L}} A(y)$.

(f) For each $x \in \omega$ there exists a $y \in \omega$ s.t. $D(x) = A(y)$. ■

The definition of the Solovay functions $h(\cdot, \cdot)$ is typographically identical with (8) - (12) of § 4. Of course the provability predicate employed here is that of the theory T of Theorem 6.1 and the Δ_0 function symbol $g(\cdot)$ will be defined later in a way different from that of § 4.

(9) $h(0, x) = h_0(x)$

(10) $h(i+1, 0) = 0$

(11)

$h(i+1, x+1) = a$ if (i) $a \in M_{i+1}$;

(ii) $h(i, x) \neq h(i, x+1)$;

(iii) $h(i+1, x) R a$;

(iv) $a \triangleleft h(i, x+1)$;

(v) if $h(i+1, x) = 0$

then $D(g(x))$ non $\vdash_{\mathbf{L}} \neg \Psi_a$;

(vi) for each b satisfying (i) - (v) in place of a one has

$\forall z \leq x \text{ (Prf } (z, \ell(\bar{i+1}) = \bar{b} \rightarrow \exists y (\bar{h}(i+1, x) R h(\bar{i+1}, y) R \bar{b})) \rightarrow$

$\rightarrow \exists w \leq z \text{ Prf } (w, \ell(\bar{i+1}) = \bar{a} \rightarrow \exists y (\bar{h}(i+1, x) R h(\bar{i+1}, y) R \bar{a})))$

(vii) a is minimal among those c that satisfy

(i) - (vi) in place of a

= $h(i+1, x)$ if no a satisfying (i) - (vi) exists

$$(12) \quad \ell(0) = \ell_0$$

$$(13) \quad \ell(i+1) = \lim_{x \rightarrow \infty} h(i+1, x) \quad \text{if } h(i+1, \cdot) \text{ reaches a limit} \\ = 0 \quad \text{otherwise}$$

6.5. Lemma ($I\Sigma_1$).

$$(a) \quad \forall i \forall x \forall y (x \leq y \rightarrow h(i, x) = h(i, y) \vee h(i, x) R h(i, y))$$

$$(b) \quad \forall i \forall x h(i+1, x) \triangleleft h(i, x)$$

$$(c) \quad \forall j \forall i < j \forall x h(j, x) \triangleleft h(i, x)$$

$$(d) \quad \forall i \ell(i) = \lim_{x \rightarrow \infty} h(i, x)$$

$$(e) \quad \forall i \forall x (h(i, x) R \ell(i) \vee h(i, x) = \ell(i))$$

$$(f) \quad \forall j \forall i < j \ell(j) \triangleleft \ell(i) \quad \blacksquare$$

As in § 4 let $\ell = 0$ abbreviate $\ell(\bar{i}) = 0$ for each i .

6.6. Lemma ($I\Sigma_1 + \ell \neq 0$).

$$(a) \quad \forall i \text{ Pr } (\overline{\ell(\bar{i}) R \ell(\bar{i})})$$

$$(b) \quad \forall i \forall a \in M_i (\ell(i) R a \rightarrow \\ \rightarrow \neg \text{Pr } (\ell(\bar{i}) = \bar{a} \rightarrow \exists y (\overline{\ell(\bar{i}) R h(\bar{i}, y) R \bar{a}})))$$

$$(c) \quad \forall i \forall a \in M_i (\ell(i) R a \rightarrow \neg \text{Pr } (\overline{\ell(\bar{i}) \neq \bar{a}}))$$

Proof is essentially the same as that of Lemma 4.6. The only trouble happens with (b) and it is that the quantifier $\forall a \in M_i$ in the formula

$$\forall a \in M_i (\ell(i) R a \rightarrow \neg \text{Pr } (\overline{\ell(\bar{i}) = \bar{a} \rightarrow \exists y (\overline{\ell(\bar{i}) R h(\bar{i}, y) R \bar{a}})}))$$

is no longer bounded and so we can not claim that this formula is $\Delta_0(\Sigma_1)$. However this quantifier does not, in a sense, "mind" being bounded. That is, we apply induction on i just as in the proof of Lemma 4.6(b) to the formula

$$\forall a \in M_i (a * \square^N \perp \wedge \ell(i) R a \rightarrow \\ \rightarrow \neg \text{Pr } (\overline{\ell(\bar{i}) = \bar{a} \rightarrow \exists y (\overline{\ell(\bar{i}) R h(\bar{i}, y) R \bar{a}})}))$$

which is $\Delta_0(\Sigma_1)$ with n a free variable to obtain

$$\forall i \forall a \in M_i (a \neq 0 \wedge \ell(i) R a \rightarrow$$

$$\rightarrow \neg \text{Pr} (\ell(\bar{i}) = \bar{a} \rightarrow \exists y (\ell(\bar{i}) R h(\bar{i}, y) R \bar{a}))$$

and this formula after being prefixed by $\forall n$ turns equivalent to

$$\forall i \forall a \in M_i (\ell(i) R a \rightarrow \neg \text{Pr} (\ell(\bar{i}) = \bar{a} \rightarrow \exists y (\ell(\bar{i}) R h(\bar{i}, y) R \bar{a})))$$

q.e.d. ■

Let σ be a false Σ_1 sentence proved by T. We assume that σ is of the form $\exists x \sigma_0(x)$ where $\sigma_0(x)$ is Δ_0 and introduce in $I\Sigma_1 + \sigma$ a closed ε -term ε s.t.

$$(14) \quad I\Sigma_1 + \sigma \vdash \sigma_0(\varepsilon) \wedge \forall y < \varepsilon \neg \sigma_0(y)$$

(This is clearly possible by the (Δ_0) least number principle).
When working in $I\Sigma_1$ we can treat expressions

$$x \leq \varepsilon, x = \varepsilon \quad \text{and} \quad x \geq \varepsilon$$

as abbreviations for the Δ_0 expressions

$$\forall y < x \neg \sigma_0(y), \quad \sigma_0(x) \wedge \forall y < x \neg \sigma_0(y) \quad \text{and} \quad \exists y < x \sigma_0(y)$$

respectively.

Here is the definition of g :

(15)

$g(x) = z$ if (i) $z < x$;

(ii) $z < \varepsilon$;

(iii) there exists an $i \in \omega$ and a node $a \in M_i$
s.t.

$$\text{Prf} (z, \ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a}))$$

and $D(g(z))$ non $\vdash_{\mathbf{L}} \neg \Psi_a$,

or there exists an $i \in \omega$ and a formula A s.t.

$$\text{Prf} (z, \ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A})$$

and $D(g(z))$ non $\vdash_{\mathbf{L}} A$;

(iv) z is minimal among those satisfying (i) - (iii)

$= x$ if $x < \varepsilon$ and no z satisfying (i) - (iii) exists

= $\omega + \delta$ if $x \geq \delta$ and no z satisfying (i) - (iii) exists

The reason why we introduce the second disjunct in (iii) of (15) is that when finishing the proof of Theorem 6.1 we shall need to know that T does not prove sentences of the form $\ell(i) \neq A$ unless $\exists x A(x) \vdash_{\mathbf{L}} A$. (iii) is a way to make sure that once an unwanted sentence $\ell(i) \neq A$ is proved it results in the proof of an equally unwanted sentence of the form $\ell(i) \neq a$ which we are able to bring to a contradiction. In § 4 an analogous situation was handled by a compactness argument which depended crucially on the number of nodes accessible to each of $h(i, \cdot)$ being finite. In the present § this is not the case and so the construction has to be more alert. This twist also comes from Jumelet [27].

6.7. Lemma ($\text{I}\Sigma_1$).

(a) $\forall x \forall y (x \leq y \rightarrow D(g(y)) \vdash_{\mathbf{L}} D(g(x)))$

(b) $\forall x \geq \delta \forall y D(\omega+x) \vdash_{\mathbf{L}} D(g(y))$

(c) $\forall i \forall x \forall a \in M_i \left(\text{Prf}(x, \ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \wedge \right.$
 $\left. \wedge D(g(x)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a \rightarrow \forall y \geq x g(y) = g(x) \right)$

(d) $\forall i \forall x \forall A \in \mathbf{F}(i) \left(\text{Prf}(x, \ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A}) \wedge \right.$
 $\left. \wedge D(g(x)) \text{ non } \vdash_{\mathbf{L}} A \rightarrow \forall y \geq x g(y) = g(x) \right)$

(e) $\forall x < \delta (g(x) \neq x \rightarrow$

$\rightarrow \exists i (\exists a \in M_i (\text{Pr}(\ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \wedge$
 $\wedge \forall z D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a) \vee$

$\vee \exists A \in \mathbf{F}(i) (\text{Pr}(\ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A}) \wedge \forall z D(g(z)) \text{ non } \vdash_{\mathbf{L}} A))$)

(f) $\forall x \geq \delta (g(x) \neq \omega + x \rightarrow$

$\rightarrow \exists i (\exists a \in M_i (\text{Pr}(\ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \wedge$
 $\wedge \forall z D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a) \vee$

$\vee \exists A \in \mathbf{F}(i) (\text{Pr}(\ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A}) \wedge \forall z D(g(z)) \text{ non } \vdash_{\mathbf{L}} A))$)

Proof. By inspection of (15) we see that either for some $z < \delta$ the conditions (ii) - (iv) of (15) hold in which case for all $x \geq z$ there holds $g(x) = z$, or δ exists and for all $y <$

Δ we have $g(y) = y$ and for all $x \geq \Delta$, $g(x) = \omega + \Delta$.

In the former case (a) is a direct consequence of Lemma 6.4(a) and in the latter case (a) follows from Lemma 6.4(a) and (b).

Clause (b) is also immediate.

(c) and (d) are proved in perfect analogy with Lemma 4.7(b).

(e) and (f) enjoy proofs similiar to that of Lemma 4.7(c). ■

6.8. Lemma ($I\Sigma_1$).

(a) $l \neq 0 \rightarrow$

$\rightarrow \forall i \forall x \forall A \in \mathbf{F}(i) (h(i, x) = 0 \wedge D(g(x)) \vdash_{\mathbf{L}} A \rightarrow l(i) \# A)$

(b) $l = 0 \rightarrow$

$\rightarrow \forall i \forall a \in M_i \overline{\text{Pr}(l(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a}))} \rightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} \neg \Psi_a$

(c) $l = 0 \rightarrow \forall i \forall a \in M_i \overline{\text{Pr}(l(\bar{i}) \neq \bar{a})} \rightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} \neg \Psi_a$

(d) $l = 0 \rightarrow$

$\rightarrow A \in \mathbf{F} \overline{\text{Pr}(l \neq 0 \rightarrow l(\bar{i}) \# \bar{A})} \rightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} A$

Proof. Clauses (a) - (c) are proved in the same way as those of Lemma 4.8. When handling (b) however we have to execute a trick similiar to that in the proof of Lemma 6.6(b), that is, before applying induction we impose on $\forall a \in M_i$ the dummy bound $a \# \square^{\mathbf{n}} \perp$.

(d) Assume $l = 0$ and $\overline{\text{Pr}(x, l \neq 0 \rightarrow l(\bar{i}) \# \bar{A})}$. If $D(g(x)) \vdash_{\mathbf{L}} A$ then we are done.

If $D(g(x))$ non $\vdash_{\mathbf{L}} A$ then by Lemma 6.7(d) $\forall y \geq x g(y) = g(x)$ and so Lemma 3.5(g) provides a node $a \in M_i$ s.t.

$\forall y D(g(y))$ non $\vdash_{\mathbf{L}} \neg \Psi_a$ and $a \# \neg A$

We have $\overline{\text{Pr}(l \neq 0 \rightarrow l(\bar{i}) \neq \bar{a})}$ whence $\overline{\text{Pr}(l(\bar{i}) \neq \bar{a})}$. Now (c) brings us to a contradiction. ■

In order to construct the desired embedding we have to define an analogon of forcing relation at 0. This analogon will be denoted by $T(\cdot)$. And even before we construct $T(\cdot)$ we have to introduce some notation.

In $I\Sigma_1$, we shall think of lower case Greek letters from the beginning of the alphabet as variables ranging over finite strings of ι 's and τ 's. Λ is the empty string. The Δ_0 function $lh(\cdot)$ tells the number of "digits" in a string. The i th digit in α is written as $(\alpha)_i$. We shall always be careful enough not to use $(\alpha)_i$ when $lh(\alpha) > i$. Stipulate also that each string begins with its first digit so that the expression $(\alpha)_0$ is meaningless. We write $\alpha < \beta$ if α and β are strings of equal length and α lexicographically precedes β . If one adopts the first coding of strings that comes to mind then $<$ can be taken to coincide with the usual ordering of integers. Finally, $\alpha \subseteq \beta$ means that α is an initial segment of β .

In fact we shall identify strings of length i with elements of $A^0(i)$ so that when we say " α supports A " where $lh(\alpha) = i$ we actually mean that the formula $\bigwedge \{ p_j \leftrightarrow (\alpha)_j \mid 1 \leq j \leq i \}$ supports A .

Define in $I\Sigma_1 + \sigma$:

$$(16) \quad \text{Adm}(\alpha) \equiv \text{"}\alpha \text{ supports } D(\omega+\Delta)\text{"}$$

6.9. Lemma ($I\Sigma_1 + \sigma$).

$$(a) \quad \text{Adm}(\bar{\Lambda})$$

$$(b) \quad \forall \alpha \forall \beta (\text{Adm}(\beta) \wedge \alpha \subseteq \beta \rightarrow \text{Adm}(\alpha))$$

$$(c) \quad \forall i \forall \alpha (lh(\alpha) \leq i \wedge \text{Adm}(\alpha) \rightarrow$$

$$\rightarrow \exists \beta (lh(\beta) = i \wedge \alpha \subseteq \beta \wedge \text{Adm}(\beta)))$$

$$(d) \quad \forall i \exists \alpha (lh(\alpha) = i \wedge \text{Adm}(\alpha))$$

Proof. (a) follows at once from Lemmas 5.21(a) and 6.4(d).

(b) follows from Lemma 5.21(b).

(c) By Lemma 5.21(c) if β is a string of length i suppor-

ting $D(\omega+\delta)$ then there exists a string of length $i + 1$ which also supports $D(\omega+\delta)$ and of which β is an initial segment. Applying induction on i we establish the claim. Induction is applicable because the formula

$$\forall \alpha (\text{lh}(\alpha) \leq i \wedge \text{Adm}(\alpha) \rightarrow \exists \beta (\text{lh}(\beta) = i \wedge \alpha \subseteq \beta \wedge \text{Adm}(\beta)))$$

is Π_1 over $\text{IS}_1 + \sigma$. Indeed $\text{Adm}(\beta)$ is Δ_1 over $\text{IS}_1 + \sigma$ and the condition $\text{lh}(\beta) = i$ is a primitive recursive bound on the quantifier $\exists \beta$ so taking $\text{Adm}(\beta)$ in its Π_1 form we can by an instance of the Σ_1 collection schema available in IS_1 also bring the formula $\exists \beta (\text{lh}(\beta) = i \wedge \alpha \subseteq \beta \wedge \text{Adm}(\beta))$ into Π_1 form. $\text{Adm}(\alpha)$ is to be rewritten as a Σ_1 formula.

(d) follows from (a) and (c). ■

Still in $\text{IS}_1 + \sigma$, put

$$(17) \quad \text{Adm}^+(\alpha) \equiv \text{Adm}(\alpha) \wedge \forall \beta < \alpha (\text{lh}(\alpha) = \text{lh}(\beta) \rightarrow \neg \text{Adm}(\beta))$$

6.10. Lemma ($\text{IS}_1 + \sigma$).

$$(a) \quad \forall i \exists! \alpha (\text{lh}(\alpha) = i \wedge \text{Adm}^+(\alpha))$$

$$(b) \quad \forall i \forall \alpha (\text{lh}(\alpha) \leq i \wedge \text{Adm}^+(\alpha) \rightarrow \rightarrow \exists \beta (\text{lh}(\beta) = i \wedge \alpha \subseteq \beta \wedge \text{Adm}^+(\beta)))$$

$$(c) \quad \forall \alpha \forall \beta (\text{Adm}^+(\alpha) \wedge \text{Adm}^+(\beta) \wedge \text{lh}(\alpha) \leq \text{lh}(\beta) \rightarrow \alpha \subseteq \beta)$$

In other words, $\text{Adm}^+(\cdot)$ singles out an infinite branch in the tree of finite \perp - τ -strings.

Proof. (a) follows from Lemma 6.9(d) by the (Δ_1) least number principle.

(b) Suppose $\text{Adm}^+(\alpha)$ for a string α of length $\leq i$. By (a) there exists a string β of length i s.t. $\text{Adm}^+(\beta)$. Consider the initial segment $\gamma \subseteq \beta$ of length equal to that of α . We claim $\alpha = \gamma$. For if $\alpha < \gamma$ then by Lemma 6.9(c) there exists a string δ of length i s.t. $\alpha \subseteq \delta$ and $\text{Adm}(\delta)$. Since $\alpha < \gamma$ implies $\delta < \beta$ this contradicts $\text{Adm}^+(\beta)$. Finally, it can not be the case that $\gamma < \alpha$ because then $\text{Adm}^+(\alpha)$ would not hold.

(c) Let α and β satisfy $\text{Adm}^+(\cdot)$ and let the length of β be greater than the length of α . By (b) there is a string γ of

length i prolonging α and s.t. $\text{Adm}^+(\gamma)$. Conclude by (a) $\beta = \gamma$. ■

At last we can define $T(\cdot)$:

$$(18) \quad T(A) \equiv \exists i \exists k \exists \lambda \in F^0(i+k) \exists B_1, \dots, B_k \in F(i) \exists \alpha \\
\text{lh}(\alpha) = i + k \wedge \lambda((\alpha)_1, \dots, (\alpha)_{i+k}) = \tau \wedge \\
\wedge A = \lambda(p_1, \dots, p_i, \Box B_1, \dots, \Box B_k) \wedge \forall j \leq i (j \neq 0 \rightarrow \\
\rightarrow (\alpha)_j = \tau \leftrightarrow \exists \beta (\text{lh}(\beta) = i \wedge \text{Adm}^+(\beta) \wedge (\beta)_j = \tau)) \wedge \\
\wedge \forall j \leq k (j \neq 0 \rightarrow (\alpha)_{i+j} = \tau \leftrightarrow D(\omega+\Delta) \vdash_L B_j))$$

6.11. Lemma ($I\Sigma_1 + \sigma$).

$$(a) \quad \forall i (T(p_i) \leftrightarrow \exists \beta (\text{lh}(\beta) = i \wedge \text{Adm}^+(\beta) \wedge (\beta)_i = \tau))$$

$$(b) \quad \forall A \in F (T(\Box A) \leftrightarrow D(\omega+\Delta) \vdash_L A)$$

$$(c) \quad \forall i \forall \lambda \in F^0(i) \forall B_1, \dots, B_i \in F (T(\lambda(B_1, \dots, B_i)) \leftrightarrow \\
\leftrightarrow \exists \alpha (\text{lh}(\alpha) = i \wedge \lambda((\alpha)_1, \dots, (\alpha)_i) = \tau \wedge \\
\wedge \forall j \leq i (j \neq 0 \rightarrow (\alpha)_j = \tau \leftrightarrow T(B_j)))$$

that is, $T(\cdot)$ distributes over Boolean connectives.

Proof. A routine inspection of (18). ■

6.12. Lemma ($I\Sigma_1 + \sigma$).

$$(a) \quad \forall \alpha (T(\bigwedge_{1 \leq j \leq \text{lh}(\alpha)} (p_j \leftrightarrow (\alpha)_j)) \leftrightarrow \text{Adm}^+(\alpha))$$

$$(b) \quad \forall A \in F \forall B \in F (T(\Diamond A \wedge \Box B) \rightarrow T(\Diamond(A \wedge \Box B)))$$

$$(c) \quad \forall A \in F (T(\Box A) \rightarrow T(A))$$

Proof. (a) Let $i = \text{lh}(\alpha)$.

$$T(\bigwedge_{1 \leq j \leq i} (p_j \leftrightarrow (\alpha)_j)) \leftrightarrow \forall j \leq i (j \neq 0 \rightarrow T(p_j) \leftrightarrow (\alpha)_j = \tau) \\
\text{(by Lemma 6.11(c))} \\
\leftrightarrow \forall j \leq i (j \neq 0 \rightarrow \exists \beta (\text{lh}(\beta) = j \wedge \\
\text{Adm}^+(\beta) \wedge (\beta)_j = \tau) \leftrightarrow (\alpha)_j = \tau) \\
\text{(by Lemma 6.11(a))} \\
\leftrightarrow \forall j \leq i (j \neq 0 \rightarrow \exists \gamma (\text{lh}(\gamma) = i \wedge \\
\text{Adm}^+(\gamma) \wedge (\gamma)_j = \tau) \leftrightarrow (\alpha)_j = \tau) \\
\text{(by Lemma 6.10(b) and (c))} \\
\leftrightarrow \forall \gamma (\text{lh}(\gamma) = i \wedge \text{Adm}^+(\gamma) \rightarrow \\
\rightarrow \forall j \leq i (j \neq 0 \rightarrow (\gamma)_j = (\alpha)_j))$$

(by Lemma 6.10(a))

$$\leftrightarrow \forall \gamma \text{ (lh}(\gamma) = i \wedge \text{Adm}^+(\gamma) \rightarrow \gamma = \alpha)$$

$$\leftrightarrow \text{Adm}^+(\alpha) \quad (\text{by Lemma 6.10(a)}) \quad \text{q.e.d.}$$

(b) If $T(\diamond A \wedge \Box B)$ then by Lemma 6.11(b) and (c) we have that $D(\omega+\Delta) \vdash_{\mathbf{L}} B$ and hence $D(\omega+\Delta) \vdash_{\mathbf{L}} \Box B$. Suppose $T(\diamond(A \wedge \Box B))$ is not the case, that is, by Lemma 6.11(b) $D(\omega+\Delta) \vdash_{\mathbf{L}} \Box B \rightarrow \neg A$. But then $D(\omega+\Delta) \vdash_{\mathbf{L}} \neg A$ which contradicts $T(\diamond A)$.

(c) We shall prove that $T(A)$ implies $T(\diamond A)$.

Let A be $\lambda(p_1, \dots, p_i, \Box B_1, \dots, \Box B_k)$ for λ a Boolean formula. Bring λ into the disjunctive normal form:

$$\bigvee_j \left(\bigwedge_n (p_n \leftrightarrow (\alpha)_n^j) \wedge \bigwedge_m (\Box B_m \leftrightarrow (\beta)_m^j) \right)$$

for α and β appropriate matrices of \perp 's and \top 's. Since we have $T(A)$ by Lemma 6.11(c) this implies the existence of a j_0 s.t.

$$T\left(\bigwedge_n (p_n \leftrightarrow (\alpha)_n^{j_0}) \wedge \bigwedge_m (\Box B_m \leftrightarrow (\beta)_m^{j_0})\right)$$

Clearly the above is equivalent to

$$T\left(\bigwedge_n (p_n \leftrightarrow (\alpha)_n) \wedge \bigwedge_m \Box C_m \wedge \bigwedge_l \diamond E_l\right)$$

for the obvious choice of C_m 's and E_l 's (we let $(\cdot)_i$ stand for $(\cdot)_i^{j_0}$). From this it follows on the strength of (a) that α supports $D(\omega+\Delta)$ and so by Lemma 5.20

$$D(\omega+\Delta) \text{ non } \vdash_{\mathbf{L}} \bigwedge_n (p_n \leftrightarrow (\alpha)_n) \rightarrow \bigvee_l \Box \neg E_l$$

therefore

$$T(\diamond(\bigwedge_n (p_n \leftrightarrow (\alpha)_n) \wedge \bigwedge_l \diamond E_l))$$

With the help of (b) this yields

$$T(\diamond(\bigwedge_n (p_n \leftrightarrow (\alpha)_n) \wedge \bigwedge_m \Box C_m \wedge \bigwedge_l \diamond E_l))$$

that is, $T(\diamond A)$. ■

As in § 4 we define a mapping $^\circ : \{p_i\}_{i \in \omega - \{0\}} \rightarrow \mathfrak{D}_{\mathbf{T}}$:

$$(19) \quad p_i^\circ \equiv \ell \neq 0 \wedge \ell(\bar{I}) * \overline{p_i} \vee \ell = 0 \wedge T(\overline{p_i})$$

$^\circ$ is prolonged to all modal formulae.

6.13. Lemma.

(a) For each $i \in \omega$ and for each modal formula $A(p_1, \dots, p_i)$ there holds

$$I\Sigma_1 \vdash \ell \neq 0 \rightarrow (A(p_1, \dots, p_i))^\circ \leftrightarrow \ell(\bar{i}) * \overline{A(p_1, \dots, p_i)}$$

$$(b) I\Sigma_1 \vdash \forall i \forall A \in F(i) \Pr \overline{(\ell \neq 0 \rightarrow A^\circ \leftrightarrow \ell(\bar{i}) * \bar{A})}$$

$$(c) I\Sigma_1 \vdash \ell = 0 \rightarrow \forall A \in F (\Pr \overline{A^\circ} \rightarrow \exists x D(g(x)) \vdash_L A)$$

$$(d) I\Sigma_1 + \sigma \vdash \ell = 0 \rightarrow g(\Delta) = \omega + \Delta$$

(e) For each $i \in \omega$ and for each modal formula $A(p_1, \dots, p_i)$ there holds

$$I\Sigma_1 + \sigma \vdash (A(p_1, \dots, p_i))^\circ \leftrightarrow \\ \leftrightarrow (\ell \neq 0 \wedge \ell(\bar{i}) * \overline{A(p_1, \dots, p_i)} \vee \ell = 0 \wedge T(\overline{A(p_1, \dots, p_i)}))$$

Proof. (a) Analogous to the proof of Lemma 4.10. The assumption $\ell \neq 0$ being a Σ_1 sentence finds its way inside the provability predicate and therefore validates the induction step for \square .

(b) is proved by (Σ_1) induction on the structure of A . For the induction step, formalize that of (a).

(c) follows at once from (b) and Lemma 6.8(d).

(d) Suppose $g(\Delta) \neq \omega + \Delta$. By Lemma 6.7(f) this means that either

(i) there exists an $i \in \omega$ and a node $a \in M_i$ s.t.

$$\Pr \overline{(\ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a}))} \text{ and} \\ \forall z D(g(z)) \text{ non } \vdash_L \neg \Psi_a$$

or

(ii) there exists an $i \in \omega$ and a formula $A \in F(i)$ s.t.

$$\Pr \overline{(\ell \neq 0 \rightarrow \ell(\bar{i}) * \bar{A})} \text{ and } \forall z D(g(z)) \text{ non } \vdash_L A$$

But (i) contradicts Lemma 6.8(b) and (ii) contradicts Lemma 6.8(d). Thus $g(\Delta) = \omega + \Delta$.

(e) We execute induction on A . Suppose $A \in F(i)$. The only

interesting case is \Box . So assume A is $\Box B$ and go inside $\text{I}\Sigma_1 + \sigma$. By (a) we may also assume $\ell = 0$.

(\rightarrow) We have $\text{Pr}(\overline{B^\circ})$ whence by (b) $\text{Pr}(\overline{\ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A}})$ and so from Lemma 6.8(d) we have $\exists x D(g(z)) \vdash_{\mathbf{L}} B$. Hence by Lemma 6.7(b) $D(\omega+\delta) \vdash_{\mathbf{L}} B$. Lemma 6.11(b) yields then $\text{T}(\overline{\Box B})$.

(\leftarrow) Since $\text{T}(\Box B)$ is Σ_1 over $\text{I}\Sigma_1 + \sigma$ it implies $\text{Pr}(\overline{\text{T}(\Box B)})$ and hence $\text{Pr}(\overline{\text{T}(\bar{B})})$ by Lemma 6.12(c) formalized. Therefore $\text{Pr}(\overline{\ell = 0 \rightarrow \text{T}(\bar{B})})$.

On the other hand, $\text{T}(\Box B)$ is equivalent to $D(\omega+\delta) \vdash_{\mathbf{L}} B$ and hence by (d) to $D(g(\delta)) \vdash_{\mathbf{L}} B$. Since clearly $h(i, \delta) = 0$ for each $i \in \omega$ it is seen through Lemma 6.8(a) formalized that $\text{Pr}(\overline{\ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{B}})$.

Thus $\text{T}(\Box B)$ implies $\text{Pr}(\overline{B^\circ})$. ■

For proofs of lemmata of the kind represented by Lemmas 4.10 and 6.13(e) (i.e. lemmas of the form $\vdash A^\circ \leftrightarrow \ell \neq A$) it is typical to use some property like $\text{Pr}(\overline{\bar{\ell} R \ell})$ which is usually enjoyed by all nodes of the model but the root 0. Therefore these lemmas usually need the assumption that the function h leaves 0 unless the node 0 is *reflexive*, that is $0 \neq \Box A$ implies $0 \neq A$ for all the relevant formulas A . In the latter case the proof goes through equally well (cf. Solovay [47]). Another way one can use this observation is to let h jump to a reflexive node the moment some Δ_0 event happens. A clever choice of this Δ_0 event can help to obtain an h with some extra desirable properties. This idea flowered in Beklemishev [5] and [6].

In most applications the number of formulas for which it is important that $0 \neq \Box A \rightarrow A$ is finite. Our construction, on the contrary, purports to take care of all the infinite collection of modal formulae. Moreover, our Kripke models do not stay the same and since the diagonalizable algebras we deal with are not generally ω -consistent we can not generally do with models whose root comes close to being reflexive. Recall however that the

theory T of the present § believes that there exists a nonstandardly large recursive number Δ . So the way out is to fool T into thinking that after the moment Δ the model stops changing and the root of this frozen model is reflexive. This is the content of Lemma 6.12(c). In fact to achieve reflexivity of 0 at the moment Δ we have to delete a nonstandardly sophisticated collection of nodes which is specified by Lemma 5.28.

It should be noted that the construction in § 6 of Beklemishev [5] also may be thought of as chopping off certain parts of the Kripke model at a nonstandard moment so as to make the root eventually reflexive, and that that construction led us to the one presented in this §.

6.14. Lemma. $\mathbb{N} \models "g \text{ is the identity function}"$

Proof. If $g(x) \neq x$ for some $x \in \omega$ then it must be for one of the two reasons given in Lemma 6.7(e). Lemma 6.8(b) and (d) shows that either of the two reasons implies $\ell \neq 0$. Quod non. ■

6.15. Lemma. *If there holds $A(v_1, v_2, \dots) = \top$ for $A(x_1, x_2, \dots)$ a diagonalizable polynomial then*

$$I\Sigma_1 + \sigma \vdash (A(p_1, p_2, \dots))^\circ$$

Proof. Let $A(p_1, p_2, \dots) \in F(i)$. As in Lemma 4.11 we have that $D(g(m)) \vdash_L D(m) \vdash_L A(p_1, \dots, p_i)$ for some $m \in \omega$. Note that $h(i, m) = 0$.

Reason in $I\Sigma_1 + \sigma$. By Lemmas 6.11(b) and 6.7(b) we have $T(\overline{\Box A(p_1, \dots, p_i)})$ hence by Lemma 6.12(c) $T(\overline{A(p_1, \dots, p_i)})$. If $\ell \neq 0$ then by Lemma 6.8(a) $\ell(i)$ forces $A(p_1, \dots, p_i)$.

In view of Lemma 6.13(e) this amounts to $(A(p_1, \dots, p_i))^\circ$ q.e.d. ■

In full analogy with § 4 we define $^* : \text{rng } \nu \rightarrow \mathfrak{D}_T :$

$$(20) \quad (\nu i)^* \equiv p_i^\circ$$

and show that * embeds \mathfrak{D} into $\mathfrak{D}_T :$

6.16. Proof of Theorem 6.1

is concluded in nearly the same manner as the proof of Theorem 4.1 (see 4.12). The only difference is that instead of the compactness argument in 4.12 we use Lemma 6.8(d) to see that $T \vdash (A(\nu_1, \nu_2, \dots))^*$ implies $A(\nu_1, \nu_2, \dots) = \tau$. ■

The reasons why the proof of Theorem 6.1 requires the use of an infinite sequence of increasingly restrictive conditions on the range of the Solovay function h (cf. (v) of (11)) to carry out the embedding of \mathfrak{D} into \mathfrak{D}_T are somewhat deeper than those of Theorem 4.1. Even if one is going to model in \mathfrak{D}_T a finitely generated diagonalizable algebra it will not generally do to impose on h the constant condition

$$\text{if } h(x) = 0, \text{ then } h(x+1) = 0 \text{ or } \forall m h(x+1) \neq D(m)$$

even in the case when this condition is recursive. To see that, think of the diagonalizable algebra of infinite height on just one generator a which satisfies the relation

$$a \rightarrow \diamond^n_T = \tau$$

for each $n \in \omega$ and $a \neq 1$.

$$(g) \ell_0 \neq 0 \rightarrow \Pr (\overline{\ell_0 \neq \bar{\ell}_0})$$

$$(h) \ell_0 \neq 0 \rightarrow \Pr (\overline{\bar{\ell}_0 R \ell_0})$$

7.3. Lemma.

$$(a) \text{I}\Sigma_1 \vdash \forall n (n \neq 0 \rightarrow \Pr^n (\bar{1}) \leftrightarrow \ell_0 * \Box^n \perp)$$

$$(b) \text{For no } a \in M_0 \text{ do we have } \text{T} \vdash \ell_0 \neq \bar{a}.$$

$$(c) \text{N} \vDash \ell_0 = 0$$

As usual we fix a positive numeration $\nu : \omega - \{0\} \rightarrow \mathcal{D}$ and a Δ_0 enumeration $\{A(m)\}_{m \in \omega}$ of the set of modal formulae that ν brings to τ . $\{A(m)\}_{m \in \omega}$ gives rise to a better manageable sequence $\{D(m)\}_{m \in \omega}$. As in § 6 our main concern is to guarantee that 0 is reflexive, that is $0 * \Box A$ implies $0 * A$ for each formula A . This turns out to be possible once we secure that each of the formulas in $\{D(m)\}_{m \in \omega}$ is apt.

$$(4) D(0) = \tau$$

$$(5) k(0) = 0$$

$$(6) D(x+1) = A(x) \quad \text{if (i) } A(x) \vdash_{\text{T}} D(x) ;$$

$$(ii) A(x) \vdash_{\text{L}} A(k(x)) ;$$

$$(iii) \Box^+ A(x) \text{ is conservative and}$$

$$(iv) A(x) \text{ is apt}$$

$$= D(x) \quad \text{otherwise}$$

$$(7) k(x+1) = k(x) + 1 \quad \text{if } D(x+1) \vdash_{\text{L}} A(k(x))$$

$$= k(x) \quad \text{otherwise}$$

Recall that $[A]_{i,i}$ is the (i, p_1, \dots, p_i) -shadow of the formula A (see Definition 5.5). All notation concerning \perp - τ -strings not explained here comes from § 6.

7.4. Lemma.

$$(a) \text{I}\Sigma_1 \vdash \forall x \forall y (x \leq y \rightarrow D(y) \vdash_{\text{L}} D(x))$$

$$(b) \text{I}\Sigma_1 \vdash \forall x \text{"}\Box^+ D(x) \text{ is conservative"}$$

$$(c) \text{I}\Sigma_1 \vdash \forall i \forall x \text{"}[}\Box^+ D(x)]_{i,i} \text{ is apt"}$$

(d) $\text{I}\Sigma_1 \vdash \forall i \forall j \forall x \forall \alpha (i \leq j \rightarrow$

$\rightarrow \text{"}\alpha \text{ supports } [\square^+ D(x)]_{j,j} \text{"} \rightarrow \text{"}\alpha \text{ supports } [\square^+ D(x)]_{i,i} \text{"}$)

(e) For each $y \in \omega$ there exists an $x \in \omega$ s.t. $D(x) \vdash_{\mathbf{L}} A(y)$.

(f) For each $x \in \omega$ there exists a $y \in \omega$ s.t. $D(x) = A(y)$.

Proof. (a), (b) and (f) are unproblematic.

(c) This is easy for $x = 0$. For $x > 0$, let $D(x) = A(y)$. From (iv) of (6) and by Lemma 5.26 we have that $\square^+ A(y)$ is apt and therefore by Corollary 5.27 $[\square^+ D(x)]_{i,i}$ is also apt.

(d) By Lemma 5.6

$$\vdash_{\mathbf{L}} [\square^+ D(x)]_{i,i} \leftrightarrow [[\square^+ D(x)]_{j,j}]_{i,i}$$

Since α supports $[\square^+ D(x)]_{j,j}$, $[\square^+ D(x)]_{j,j}$ is (j,j) -trimmed by Lemmas 5.11 and 5.25(a) and apt by (c) we have by Lemma 5.25(e) that α supports $[\square^+ D(x)]_{i,i}$.

(e) Suppose y is the minimal s.t. $D(x) \vdash_{\mathbf{L}} A(y)$ for no $x \in \omega$. Then $\lim_{x \rightarrow \infty} k(x) = y$ and $\lim_{x \rightarrow \infty} D(x) = A(z)$ for some $z < y$. By Lemma 5.16 there exists an apt formula B s.t. $B(\nu_1, \nu_2, \dots) = \tau$ and $B \vdash_{\mathbf{L}} A(z) \wedge A(y)$. Since the height of \mathcal{D} is infinite $\square^+ B$ is conservative. Note that the number of formulas \mathbf{L} -equivalent to B is infinite and therefore for some $w > y$ one has $\vdash_{\mathbf{L}} A(w) \leftrightarrow B$. But then clearly $D(w+1) = A(w) \vdash_{\mathbf{L}} B \vdash_{\mathbf{L}} A(y)$. ■

The following definitions are cited verbatim from § 6.

(8) $h(0, x) = h_0(x)$

(9) $h(i+1, 0) = 0$

(10)

$$h(i+1, x+1) = a \text{ if } \begin{array}{l} \text{(i) } a \in M_{i+1} ; \\ \text{(ii) } h(i, x) \neq h(i, x+1) ; \\ \text{(iii) } h(i+1, x) R a ; \\ \text{(iv) } a \triangleleft h(i, x+1) ; \\ \text{(v) if } h(i+1, x) = 0 \end{array}$$

then $D(g(x))$ non $\vdash_L \neg \Psi_a$;

(vi) for each b satisfying (i) - (v) in place of a one has

$$\forall z \leq x \text{ (Prf } (z, \overline{\ell(i+1)} = \bar{b} \rightarrow \exists y (\overline{h(i+1,x)} R h(i+1,y) R \bar{b})) \rightarrow \\ \rightarrow \exists w \leq z \text{ Prf } (w, \overline{\ell(i+1)} = \bar{a} \rightarrow \exists y (\overline{h(i+1,x)} R h(i+1,y) R \bar{a})))$$

(vii) a is minimal among those c that satisfy (i) - (vi) in place of a

$= h(i+1,x)$ if no a satisfying (i) - (vi) exists

(11) $\ell(0) = \ell_0$

(12) $\ell(i+1) = \lim_{x \rightarrow \infty} h(i+1,x)$ if $h(i+1, \cdot)$ reaches a limit
 $= 0$ otherwise

7.5. Lemma ($I\Sigma_1$).

(a) $\forall i \forall x \forall y (x \leq y \rightarrow h(i,x) = h(i,y) \vee h(i,x) R h(i,y))$

(b) $\forall i \forall x h(i+1,x) \triangleleft h(i,x)$

(c) $\forall j \forall i < j \forall x h(j,x) \triangleleft h(i,x)$

(d) $\forall i \ell(i) = \lim_{x \rightarrow \infty} h(i,x)$

(e) $\forall i \forall x (h(i,x) R \ell(i) \vee h(i,x) = \ell(i))$

(f) $\forall j \forall i < j \ell(j) \triangleleft \ell(i)$ ■

As usual $\ell = 0$ means $\ell(\bar{i}) = 0$ for each i .

7.6. Lemma ($I\Sigma_1 + \ell \neq 0$).

(a) $\forall i \text{ Pr } (\overline{\ell(i)} R \ell(\bar{i}))$

(b) $\forall i \forall a \in M_i (\ell(i) R a \rightarrow \\ \rightarrow \neg \text{Pr } (\overline{\ell(\bar{i})} = \bar{a} \rightarrow \exists y (\overline{\ell(i)} R h(\bar{i},y) R \bar{a})))$

(c) $\forall i \forall a \in M_i (\ell(i) R a \rightarrow \neg \text{Pr } (\overline{\ell(\bar{i})} \neq \bar{a}))$ ■

(13)

$g(x) = z$ if (i) $z < x$;

(ii) there exists an $i \in \omega$ and a node $a \in M_i$

s.t.

$$\text{Prf } (z, \ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a}))$$

and $D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a$,

or there exists an $i \in \omega$ and a formula A s.t.

$$\text{Prf } (z, \ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A})$$

and $D(g(z)) \text{ non } \vdash_{\mathbf{L}} A$;

(iii) z is minimal among those satisfying (i) and (ii)

= x if no z satisfying (i) and (ii) exists

7.7. Lemma ($\text{I}\Sigma_1$).

$$(a) \forall x \forall y (x \leq y \rightarrow D(g(y)) \vdash_{\mathbf{L}} D(g(x)))$$

$$(b) \forall i \forall x \forall a \in M_i \left(\text{Prf } (x, \ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \wedge \right. \\ \left. \wedge D(g(x)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a \rightarrow \forall y \geq x g(y) = g(x) \right)$$

$$(c) \forall i \forall x \forall A \in F(i) \left(\text{Prf } (x, \ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A}) \wedge \right. \\ \left. \wedge D(g(x)) \text{ non } \vdash_{\mathbf{L}} A \rightarrow \forall y \geq x g(y) = g(x) \right)$$

$$(d) \forall x (g(x) \neq x \rightarrow$$

$$\rightarrow \exists i (\exists a \in M_i \left(\text{Pr } (\ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \wedge \right. \\ \left. \wedge \forall z D(g(z)) \text{ non } \vdash_{\mathbf{L}} \neg \Psi_a \right) \vee$$

$$\vee \exists A \in F(i) \left(\text{Pr } (\ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A}) \wedge \forall z D(g(z)) \text{ non } \vdash_{\mathbf{L}} A \right)) \blacksquare$$

7.8. Lemma ($\text{I}\Sigma_1$).

$$(a) \ell \neq 0 \rightarrow$$

$$\rightarrow \forall i \forall x \forall A \in F(i) (h(i, x) = 0 \wedge D(g(x)) \vdash_{\mathbf{L}} A \rightarrow \ell(i) \neq A)$$

$$(b) \ell = 0 \rightarrow$$

$$\rightarrow \forall i \forall a \in M_i \left(\text{Pr } (\ell(\bar{i}) = \bar{a} \rightarrow \exists y (0 R h(\bar{i}, y) R \bar{a})) \rightarrow \right. \\ \left. \rightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} \neg \Psi_a \right)$$

$$(c) \ell = 0 \rightarrow \forall i \forall a \in M_i \left(\text{Pr } (\ell(\bar{i}) \neq \bar{a}) \rightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} \neg \Psi_a \right)$$

$$(d) \ell = 0 \rightarrow$$

$$\rightarrow \forall A \in \mathcal{F} \quad (\text{Pr } (\ell \neq 0 \rightarrow \ell(\bar{i}) \neq \bar{A}) \rightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} A) \quad \blacksquare$$

Consider the (i, i) -shadow $[\square^+ D(g(x))]_{i, i}$ of $\square^+ D(g(x))$. Since the number of formulas in $\mathcal{F}^i(i)$ is finite modulo \mathbf{L} -equivalence and the sequence $\{\square^+ D(g(x))\}_{x \in \omega}$ grows in $\vdash_{\mathbf{L}}$ -strength we have that the sequence $\{[\square^+ D(g(x))]_{i, i}\}_{x \in \omega}$ reaches a limit. By an appropriate instance of the Σ_1 least number principle this observation is formalizable in $\mathbf{I}\Sigma_1$ and so we can introduce a new term $[\square^+ D(g)]_{i, i}$ in this theory so that $\mathbf{I}\Sigma_1$ proves

$$(14) \quad [\square^+ D(g)]_{i, i} = \lim_{x \rightarrow \infty} [\square^+ D(g(x))]_{i, i}$$

The definition of the formula $\text{Adm}(\cdot)$ is slightly different from the one in § 6.

$$(15) \quad \text{Adm}(\alpha) \equiv \forall i \text{ "}\alpha \text{ supports } [\square^+ D(g)]_{i, i}\text{"}$$

7.9. Lemma.

$$(a) \quad \mathbf{I}\Sigma_1 \vdash \text{Adm}(\bar{\Lambda})$$

$$(b) \quad \mathbf{I}\Sigma_1 \vdash \forall \alpha \forall \beta (\text{Adm}(\beta) \wedge \alpha \subseteq \beta \rightarrow \text{Adm}(\alpha))$$

(c) For each $i \in \omega$

$$\mathbf{I}\Sigma_1 \vdash \forall \alpha (\text{lh}(\alpha) \leq \bar{i} \wedge \text{Adm}(\alpha) \rightarrow$$

$$\rightarrow \exists \beta (\text{lh}(\beta) = \bar{i} \wedge \alpha \subseteq \beta \wedge \text{Adm}(\beta)))$$

(d) For each $i \in \omega$

$$\mathbf{I}\Sigma_1 \vdash \exists \alpha (\text{lh}(\alpha) = \bar{i} \wedge \text{Adm}(\alpha))$$

Proof. (a) By Lemma 7.4(c) the formula $[\square^+ D(g)]_{i, i}$ is apt for each i and the claim follows by Lemma 5.21(a).

(b) follows from Lemma 5.21(b).

(c) The proof is analogous to that of Lemma 6.9(c). In the present situation however we have to do with external induction because the formula we induct on is too arithmetically complex. To carry out the induction step we have to show that if for all i a string α of length i supports $[\square^+ D(g)]_{j, j}$ then there exists a string β of length $i + 1$ extending α which supports

$[\Box^+D(g)]_{j,j}$ for all j . Now if this were true of neither of the two candidates β_1 and β_2 for the role of β then there would exist j_1 and j_2 s.t. α_i does not support $[\Box^+D(g)]_{j_i,j_i}$, $i = 1, 2$. Let j be the maximum of j_1 and j_2 . By Lemma 7.4(d) we have that neither β_1 nor β_2 supports $[\Box^+D(g)]_{j,j}$. Hence by Lemma 5.21(c) α does not support $[\Box^+D(g)]_{j,j}$. This proves the claim.

(d) follows from (a) and (c). ■

In Lemma 7.9(c) and (d) we can not retain the majestic uniformity of Lemma 6.9. In fact by using results of Adamowicz [1] combined with methods of § 9 it can be shown that there exists a positive numeration μ of a diagonalizable algebra with the strong disjunction property s.t. the statement

$$\forall i \exists \alpha (\text{lh}(\alpha) = i \wedge \text{Adm}(\alpha))$$

for the formula $\text{Adm}(\cdot)$ built up starting from μ is not provable in $\text{I}\Sigma_1$ (nor even in the theory of all the Π_3 truths). Hence the loss of uniformity in many of the succeeding lemmas, most notably in Lemma 7.12(c). I do not know whether a cleverer definition of $\text{Adm}(\cdot)$ might help.

$$(16) \quad \text{Adm}^+(\alpha) \equiv \text{Adm}(\alpha) \wedge \forall \beta < \alpha (\text{lh}(\alpha) = \text{lh}(\beta) \rightarrow \neg \text{Adm}(\beta))$$

7.10. Lemma. For each $i \in \omega$ $\text{I}\Sigma_1$ proves

$$(a) \quad \exists ! \alpha (\text{lh}(\alpha) = \bar{i} \wedge \text{Adm}^+(\alpha))$$

$$(b) \quad \forall \alpha (\text{lh}(\alpha) \leq \bar{i} \wedge \text{Adm}^+(\alpha) \rightarrow$$

$$\rightarrow \exists \beta (\text{lh}(\beta) = \bar{i} \wedge \alpha \subseteq \beta \wedge \text{Adm}^+(\beta)))$$

$$(c) \quad \forall \alpha \forall \beta (\text{Adm}^+(\alpha) \wedge \text{Adm}^+(\beta) \wedge \text{lh}(\alpha) \leq \text{lh}(\beta) \leq \bar{i} \rightarrow \alpha \subseteq \beta)$$

■

$$(17) \quad \text{T}(A) \equiv \exists i \exists k \exists \lambda \in \mathbf{F}^0(i+k) \exists B_1, \dots, B_k \in \mathbf{F}(i) \exists \alpha$$

$$(\text{lh}(\alpha) = i + k \wedge \lambda((\alpha)_1, \dots, (\alpha)_{i+k}) = \tau \wedge$$

$$\wedge A = \lambda(p_1, \dots, p_i, \Box B_1, \dots, \Box B_k) \wedge \forall j \leq i (j \neq 0 \rightarrow$$

$$\rightarrow (\alpha)_j = \tau \leftrightarrow \exists \beta (\text{lh}(\beta) = i \wedge \text{Adm}^+(\beta) \wedge (\beta)_j = \tau)) \wedge$$

$$\wedge \forall j \leq k (j \neq 0 \rightarrow (\alpha)_{i+j} = \tau \leftrightarrow \exists x D(g(x)) \Vdash_{\mathbf{L}} B_j))$$

7.11. Lemma ($I\Sigma_1$).

- (a) $\forall i (T(p_i) \leftrightarrow \exists \beta (\text{lh}(\beta) = i \wedge \text{Adm}^+(\beta) \wedge (\beta)_i = \tau))$
 (b) $\forall A \in F (T(\Box A) \leftrightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} A)$
 (c) $\forall i \forall \lambda \in F^0(i) \forall B_1, \dots, B_i \in F (T(\lambda(B_1, \dots, B_i)) \leftrightarrow$
 $\leftrightarrow \exists \alpha (\text{lh}(\alpha) = i \wedge \lambda((\alpha)_1, \dots, (\alpha)_i) = \tau \wedge$
 $\wedge \forall j \leq i (j \neq 0 \rightarrow (\alpha)_j = \tau \leftrightarrow T(B_j))) \blacksquare$

7.12. Lemma. For each $i \in \omega$ $I\Sigma_1$ proves

- (a) $\forall \alpha (\text{lh}(\alpha) \leq \bar{i} \rightarrow T(\bigwedge_{1 \leq j \leq \text{lh}(\alpha)} (p_j \leftrightarrow (\alpha)_j)) \leftrightarrow \text{Adm}^+(\alpha))$
 (b) $\forall A \in F \forall B \in F (T(\Diamond A \wedge \Box B) \rightarrow T(\Diamond(A \wedge \Box B)))$
 (c) $\forall A \in F(\bar{i}) (T(\Box A) \rightarrow T(A))$

Proof is essentially the same as that of Lemma 6.12. The only new detail is that in (c) we have to use the fact that

$$I\Sigma_1 \vdash \forall A \in F (\exists i [\Box^+ D(g)]_{i,i} \vdash_{\mathbf{L}} A \leftrightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} A) \quad \blacksquare$$

We define the mapping $^\circ : \{p_i\}_{i \in \omega - \{0\}} \rightarrow \mathfrak{D}_{\mathbf{T}}$:

$$(18) \quad p_i^\circ \equiv \ell \neq 0 \wedge \ell(\bar{i}) * \overline{p_i} \vee \ell = 0 \wedge T(\overline{p_i})$$

and prolong it to all modal formulae.

7.13. Lemma.

(a) For each $i \in \omega$ and for each modal formula $A(p_1, \dots, p_i)$ there holds

$$I\Sigma_1 \vdash \ell \neq 0 \rightarrow (A(p_1, \dots, p_i))^\circ \leftrightarrow \ell(\bar{i}) * \overline{A(p_1, \dots, p_i)}$$

$$(b) \quad I\Sigma_1 \vdash \forall i \forall A \in F(i) \Pr(\overline{\ell \neq 0 \rightarrow A^\circ \leftrightarrow \ell(\bar{i}) * \overline{A}})$$

$$(c) \quad I\Sigma_1 \vdash \ell = 0 \rightarrow \forall A \in F (\Pr(\overline{A^\circ} \rightarrow \exists x D(g(x)) \vdash_{\mathbf{L}} A))$$

(d) For each $i \in \omega$ and for each modal formula $A(p_1, \dots, p_i)$ there holds

$$I\Sigma_1 \vdash (A(p_1, \dots, p_i))^\circ \leftrightarrow$$

$$\leftrightarrow (\ell \neq 0 \wedge \ell(\bar{i}) * \overline{A(p_1, \dots, p_i)} \vee \ell = 0 \wedge T(\overline{A(p_1, \dots, p_i)})) \quad \blacksquare$$

7.14. Lemma. $\mathbb{N} \models "g \text{ is the identity function}"$ ■

7.15. Lemma. *If there holds $A(v_1, v_2, \dots) = \top$ for $A(x_1, x_2, \dots)$ a diagonalizable polynomial then*

$$\text{IS}_1 \vdash (A(p_1, p_2, \dots))^\circ$$
 ■

Now having defined

$$(19) \quad (v_i)^* \equiv p_i^\circ$$

we can finish the proof of Theorem 7.1 just as we did in 6.16 with Theorem 6.1. ■

§ 8. An application

In this § we shall apply Theorem 7.1 to give an alternative proof of a proposition which was used in Simmons [41] to obtain some interesting information on the structure of the E -tree.

8.1. Proposition (Simmons [41]). *Let T be a Σ_1 -sound theory and τ a false Σ_1 sentence. Then there exists a family $\{\sigma_\alpha\}_{\alpha \in \mathbb{Q}}$ of Σ_1 sentences s.t.*

$$T \vdash \tau \rightarrow \sigma_\alpha$$

and

$$T \vdash \text{Pr}_T(\overline{\sigma_\alpha}) \rightarrow \sigma_\beta$$

whenever $\alpha, \beta \in \mathbb{Q}$ and $\alpha < \beta$ (Here \mathbb{Q} is the set of rationals under the natural ordering).

Proof. Consider the following set of modal formulae in propositional letters $\{p_\alpha\}_{\alpha \in \mathbb{Q}}$:

$$\mathcal{F} = \{ \Box \Box p_\alpha \rightarrow \Box p_\beta \mid \alpha, \beta \in \mathbb{Q} \text{ and } \alpha < \beta \}$$

We shall show that the quotient algebra \mathbf{F}/\mathcal{F} of the free diagonalizable algebra \mathbf{F} on the generators $\{p_\alpha\}_{\alpha \in \mathbb{Q}}$ modulo the τ -filter generated by \mathcal{F} enjoys the strong disjunction property. We fix an effective repetition-free enumeration $\{\alpha_i\}_{i \in \omega}$ of \mathbb{Q} . Then if some diagonalizable polynomial $A(p_{\alpha_0}, p_{\alpha_1}, \dots)$ hits τ in \mathbf{F}/\mathcal{F} there exists a finite subset \mathcal{F} of \mathcal{F} s.t.


$$\bigwedge \mathcal{F} \vdash_{\mathbf{L}} A(p_{\alpha_0}, p_{\alpha_1}, \dots)$$

W.l.o.g. we may assume that for some $N \in \omega$ the set \mathcal{F} looks like this:

$$\mathcal{F}_N = \{ \Box \Box p_{\alpha_i} \rightarrow \Box p_{\alpha_j} \mid i, j \leq N \text{ and } \alpha_i < \alpha_j \}$$

Claim. For each $N \in \omega$ the formula $\bigwedge \mathcal{F}_N$ is apt.

Assume for simplicity that for $i, j \leq N$ we have $\alpha_i < \alpha_j$ if and only if $i < j$. Consider the following Kripke model:



(Only the letters forced are shown at each node).

Imagine two models in which $\mathbb{M} \varphi_N$ holds grafted above the root of the model depicted. We want to get convinced that the resulting model models $\square^+ \mathbb{M} \varphi_N$. One clearly has that $\mathbb{M} \varphi_N$ is forced at every node which is not the root of the new model (for the "old" nodes this is verified by inspection of the picture). As for the root itself, note that it forces $\square \square p_{\alpha_i}$ for no $i \leq N$ and hence forces $\mathbb{M} \varphi_N$. By Lemma 5.14 conclude that $\mathbb{M} \varphi_N$ is apt and that the Claim is proved.

From the Claim it follows by Lemma 5.16 that \mathbb{F}/φ possesses the strong disjunction property. Note also that \mathbb{F}/φ is clearly positive and that the theory $T + \neg\tau$ is Σ_1 -sound. Hence by Theorem 7.1 there exists an embedding $\sigma^* : \mathbb{F}/\varphi \rightarrow \mathfrak{D}_{T+\neg\tau}$.

Now let

$$\sigma_\alpha \equiv (\square p_\alpha)^*$$

We trivially have $T + \tau \rightarrow \sigma_\alpha$ for each $\alpha \in \mathbb{Q}$ because

$$T + \tau \rightarrow \text{Pr}_T(\bar{\tau})$$

$$\begin{aligned}
&\rightarrow \Pr_{T+\neg\tau}(\bar{1}) \\
&\rightarrow \Pr_{T+\neg\tau}^*((p_\alpha)^*) \\
&\rightarrow (\Box p_\alpha) \\
&\rightarrow \sigma_\alpha
\end{aligned}$$

Let $\alpha, \beta \in \mathbb{Q}$ and $\alpha < \beta$. Then since $\Box p_\alpha \rightarrow \Box p_\beta$ is in \mathcal{S} and $*$ is an embedding we have

$$\begin{aligned}
T + \neg\tau \vdash \Pr_T(\overline{\sigma_\alpha}) &\rightarrow \Pr_{T+\neg\tau}^*(\overline{\sigma_\alpha}) \\
&\rightarrow (\Box p_\alpha)^* \\
&\rightarrow (\Box p_\beta)^* \\
&\rightarrow \sigma_\beta
\end{aligned}$$

Combining this with $T + \tau \vdash \sigma_\beta$ we get

$$T \vdash \Pr_T(\overline{\sigma_\alpha}) \rightarrow \sigma_\beta$$

q.e.d. ■

Mark that a proof of Proposition 8.1 similiar to ours is obtainable by the methods of Jumelet [27].

§ 9. A question of arithmetical complexity

In §§ 4, 6 and 7 we have constructed embeddings $*$: $\mathcal{D} \rightarrow \mathcal{D}_T$ of positive diagonalizable algebras into diagonalizable algebras of various theories T. This § takes a look at the arithmetical complexity of sentences in $\text{rng } *$. We register the arithmetical complexity of sentences in $\text{rng } *$ of the $*$'s constructed and give a lower bound on this complexity for the case of Σ_2 -sound theories T under reasonable assumptions on $*$ to the effect that our constructions were fairly optimal in this respect.

9.A. Finite credibility extent

Recall that in § 4 we had $(\nu i)^* \equiv \ell(\bar{i}) * \overline{p_i}$ where $\ell(i)$ denoted the limit of a primitive recursive function $h(i, \cdot)$ climbing up the Kripke model M_i . Moreover, the part of M_i a priori accessible to $h(i, \cdot)$ was finite since it was specified by the condition $a * \square^n \perp$, n being the credibility extent of T. Therefore

$$I\Sigma_1 \vdash \ell(\bar{i}) * \overline{p_i} \leftrightarrow \bigvee \{ \ell(\bar{i}) = \bar{a} \mid a \in M_i \text{ and } a * \square^n \perp \wedge p_i \}$$

Now $\ell(\bar{i}) = \bar{a}$ is equivalent to the statement

$$\forall x (h(i, x) R \bar{a} \vee h(i, x) = \bar{a}) \wedge \exists x h(i, x) = \bar{a}$$

which is a Boolean combination of Σ_1 sentences and therefore each sentence in $\text{rng } *$ is a Boolean combination of Σ_1 sentences over $I\Sigma_1$. (These considerations come from Solovay [47].)

9.B. Σ_1 -ill theories of infinite credibility extent

From § 6 we have

$$(\nu i)^* \equiv \ell \neq 0 \wedge \ell(\bar{i}) * \overline{p_i} \vee \ell = 0 \wedge T(\overline{p_i})$$

Note that $\ell \neq 0$ is Σ_1 , $\ell = 0$ is Π_1 and $\ell(\bar{i}) * \overline{p_i}$ is Δ_2 over $I\Sigma_1$. The sentence $T(\overline{p_i})$ is built from formulas of the form " α supports $D(\omega + \alpha)$ " and some Δ_0 formulas with the help of Boolean connectives and primitive recursively bounded quantification. Now formulas of this sort involving $D(\omega + \alpha)$ can be either written in Σ_1 or in Π_1 form for

$$\begin{aligned} T \vdash \dots D(\omega+\delta) \dots &\leftrightarrow \exists x (x = \delta \wedge \dots D(\omega+x) \dots) \\ &\leftrightarrow \forall x (x = \delta \rightarrow \dots D(\omega+x) \dots) \end{aligned}$$

and therefore $T(\overline{p_i})$ is a $\Delta_0(\Sigma_1)$ (and hence Δ_2) sentence over $I\Sigma_1$.

All this amounts to $(\forall i)^*$ being Δ_2 over $I\Sigma_1$ and it was shown by Gaifman and Smoryński (cf. Remark 3.5.iii in Chapter 3 of Smoryński [46]) that one can not generally do with Boolean combinations of Σ_1 sentences.

9.C. Σ_1 -sound theories

In § 7 $(\forall i)^*$ was defined just as in § 6:

$$(\forall i)^* \equiv \ell \neq 0 \wedge \ell(\bar{i}) * \overline{p_i} \vee \ell = 0 \wedge T(\overline{p_i})$$

The $T(\cdot)$ of § 7 was however considerably different from that of § 6. The most complex parts of it are subformulas of the form

$$\forall i \text{ "}\alpha \text{ supports } [\Box^+ D(g)]_{i,i}\text{"}$$

where $[\Box^+ D(g)]_{i,i} = \lim_{x \rightarrow \infty} [\Box^+ D(g(x))]_{i,i}$. These formulas are clearly Π_2 over $I\Sigma_1$ and so $(\forall i)^*$ is a $\Delta_0(\Sigma_2)$ sentence and hence a Boolean combination of Σ_2 sentences.

An important particular case when one can do better than $\Delta_0(\Sigma_2)$ is the case of finitely generated algebras. These algebras only need sentences Δ_2 over $I\Sigma_1$. That this is so can be seen as follows.

Let $\{x_1, \dots, x_n\}$ be the generators of such an algebra. When enumerating the relations holding in this algebra we can restrict our attention to formulas in $F(n)$. Let $\alpha \in \mathbf{A}^0(n)$ be s.t. $\alpha(x_1, \dots, x_n)$ is admissible. When designing the sequence $\{D(m)\}_{m \in \omega}$ we can then impose the additional requirement that α support D_x for all x (that this requirement is harmless can be seen through Lemma 5.24) and define $\text{Adm}^+(\beta)$ to be the Δ_0 formula $\beta \subseteq \bar{\alpha}$. The formula $T(\cdot)$ and $*$ are then built from $\{D(x)\}_{x \in \omega}$ in the same manner as in § 7 so that $(\forall i)^*$ is clearly Δ_2 over $I\Sigma_1$.

Another particular case admitting an improvement is that of a Σ_2 -ill theory T . It suffices then to use sentences that are Δ_2 over T . The Σ_2 -ill theory T proves that some actually total re-

cursive function f is not total. The following changes are to be introduced in § 7:


We inhibit slightly the process of gradually strengthening the formulas $\{D(m)\}_{m \in \omega}$ by the requirement that the n th change of the value of D be only allowed to take place after $f(n)$ has converged. Then the formula

$$\text{Adm}(\alpha) \equiv \forall i \text{ "}\alpha \text{ supports } [\Box^+ D(g)]_{i,i}\text{"}$$

is Δ_2 over T for T proves that the sequence $\{D(g(x))\}_{x \in \omega}$ freezes after some $\Delta_0(\Sigma_1)$ -definable moment.

The rest of this § will be devoted to an explanation why the use of highly complex sentences is generally unavoidable when embedding positive algebras in diagonalizable algebras of Σ_2 -sound theories.

Note that the embeddings $^* : \mathfrak{D} \rightarrow \mathfrak{D}_T$ constructed in preceding §§ employed an arbitrary positive numeration ν of \mathfrak{D} and the * we obtained was in each case recursive w.r.t. ν , that is, there always existed a recursive function $^\circ$ s.t. the following diagram commuted:



We are going to show that if T is a Σ_2 -sound theory then there exists a positive numeration of a diagonalizable algebra with the strong disjunction property s.t. the above diagram commutes for no pair $(^*, \circ)$ with $\text{rng } ^* \subseteq \Delta_2$ and $^\circ$ recursive.

To this end we need some definitions and lemmas.

9.1. Definition. Consider the set $2^{<\omega}$ (which we will think of as consisting of \perp - τ -strings) ordered by the usual "initial segment" relation \subseteq . A tree is a downwards closed subset of $2^{<\omega}$. ($2^{<\omega}$ grows upwards.) A tree T is *efflorescent* if it is not empty

and for each $\alpha \in T$ there exists a $\beta \in T$ s.t. α is a proper initial segment of β .

9.2. Definition. Let ν be a numeration of a diagonalizable algebra \mathfrak{D} with the strong disjunction property. The *admissibility tree* of ν is then the set

$$\{ \alpha \in 2^{<\omega} \mid \bigwedge_{1 \leq i \leq \text{lh}(\alpha)} (\nu i \leftrightarrow (\alpha)_i) \}$$

is an admissible element of \mathfrak{D} }


9.3. Lemma. *The admissibility tree of a (positive) numeration ν of a diagonalizable algebra \mathfrak{D} with the strong disjunction property is a (Π_2^0) efflorescent tree.*

Proof. That the admissibility tree is a tree follows from Lemma 5.18(b). Efflorescence is an immediate consequence of Lemma 5.18(a) and (c). If ν is positive then the equalities of diagonalizable polynomials with elements of rng^* as variables that hold in \mathfrak{D} are recursively enumerable and hence admissibility is Π_2^0 by inspection of Definition 5.17. ■

9.4. Lemma. *Let T be an arbitrary Π_2^0 efflorescent tree. There exists a positive numeration μ of a diagonalizable algebra with the strong disjunction property whose admissibility tree is T .*

Proof. Let \mathbf{F} be the free diagonalizable algebra on the generators $\{p_i\}_{i \in \omega - \{0\}}$. We shall compile an r.e. set \mathcal{C} of formulas in $\{p_i\}_{i \in \omega - \{0\}}$ and let μ be the numeration of \mathbf{F}/\mathcal{C} induced by the natural numeration of \mathbf{F} .

Define a family of Kripke models $\{g_n^\varepsilon\}_{n \in \omega}^{\varepsilon \in 2^{<\omega}}$ by letting g_n^ε be the p_1 -model shown in the picture (we assume $i = \text{lh}(\varepsilon)$):



For typographical reasons the name of the model $\mathcal{J}_n^\varepsilon$ will also stand for the formula $\Psi_{\mathcal{J}_n^\varepsilon}(p_1)$.

Claim 1. $\vdash_{\mathbf{L}} \mathcal{J}_n^\varepsilon \rightarrow \neg \mathcal{J}_m^\delta$ unless $\varepsilon = \delta$ and $n = m$.

This is an easy consequence of Lemma 3.3(c) and (d) since the models $\mathcal{J}_n^\varepsilon$ are clearly differentiated and $\mathcal{J}_n^\varepsilon$ is not isomorphic to \mathcal{J}_m^δ unless $\varepsilon = \delta$ and $n = m$.

Claim 2. $\vdash_{\mathbf{L}} \mathcal{J}_n^\varepsilon \rightarrow \Box \neg \mathcal{J}_m^\delta$, all $\varepsilon, \delta \in 2^{<\omega}$, all $n, m \in \omega$.

This follows from Lemma 3.5(b) because \mathcal{J}_m^δ is not isomorphic to a proper cone of $\mathcal{J}_n^\varepsilon$.

Now let

$$T = \{ \alpha \in 2^{<\omega} \mid \forall x \exists y R(\alpha, x, y) \}$$

with $R(\alpha, x, y)$ decidable. Consider a particular element ε of $2^{<\omega}$. Let ε be of length i . We describe the construction of a set \mathcal{C}_ε by recursive stages. One begins with Stage 0 at the beginning of

which \mathcal{E}_ε is empty. In what follows * denotes concatenation of strings.

Stage n .

Look for j_1 and j_2 in ω s.t. $R(\varepsilon * \langle 1 \rangle, n, j_1)$ and $R(\varepsilon * \langle \tau \rangle, n, j_2)$. On finding j_1 add the formula

$$\left(\bigwedge_{1 \leq j \leq i} (p_j \leftrightarrow (\varepsilon)_j) \right) \wedge p_{i+1} \rightarrow \cdot \Box \neg \mathcal{J}_n^\varepsilon$$

to \mathcal{E}_ε . Once j_2 is found one adds to \mathcal{E}_ε the formula

$$\left(\bigwedge_{1 \leq j \leq i} (p_j \leftrightarrow (\varepsilon)_j) \right) \wedge \neg p_{i+1} \rightarrow \cdot \Box \neg \mathcal{J}_n^\varepsilon$$

If one has found both j_1 and j_2 then

$$\neg \mathcal{J}_n^\varepsilon$$

is appended to \mathcal{E}_ε and we go to Stage $n + 1$.

Note that it very well may happen that the number of stages is finite. In this case the last stage takes an infinite amount of "time".

Let $\mathcal{E}_\varepsilon^n$ be the part of \mathcal{E}_ε compiled during the first n stages. Put

$$C_N = \bigwedge \{ \bigwedge \mathcal{E}_\alpha^N \mid \alpha \in 2^{\leq N} \}$$

$$\mathcal{E} = \bigcup \{ \mathcal{E}_\alpha \mid \alpha \in 2^{<\omega} \} = \{ C_N \mid N \in \omega \}$$

Clearly if $A(p_1, p_2, \dots) = \tau$ in \mathbf{F}/\mathcal{E} then $C_N \vdash_L A$ for some $N \in \omega$.

Consider the (p_1, \dots, p_N) -model \mathcal{H}_N^δ (we assume $\delta \neq \Lambda$):

$$2N \text{ nodes } \left\{ \begin{array}{l} * \quad p_1, \dots, p_N \\ \cdot \\ \cdot \\ \cdot \\ * \quad p_1, \dots, p_N \\ \vdots \\ * \quad p_1 \leftrightarrow (\delta)_1, \dots, p_N \leftrightarrow (\delta)_N \end{array} \right.$$

Let α^0 denote the string α with the last digit deleted and

let α^- stand for α with the last digit replaced by its negation. Neither 0 nor $^-$ is meant to be applied to Λ .

Suppose $\delta \in T$ and picture two models in which C_{N-1} holds grafted above the root of \mathcal{K}_N^δ . We are going to show that C_{N-1} holds in the resulting model \mathcal{K} .

Clearly C_{N-1} can not fail at any of the new nodes of \mathcal{K} . As for the non-bottom nodes of \mathcal{K}_N^δ , note that these can force $\mathcal{J}_n^\varepsilon$ for no $\alpha \in 2^{<\omega}$, $n \in \omega$ because the root of $\mathcal{J}_n^\varepsilon$ forces $\neg p_1$. Therefore we only have to consider the root of \mathcal{K} .

Let us take a closer look at C_{N-1} . The conjuncts of this formula are formulas of either of the two forms

$$\neg \mathcal{J}_n^\varepsilon \quad \text{and} \quad \left(\bigwedge_{1 \leq j \leq \text{lh}(\alpha)} (p_j \leftrightarrow (\alpha)_j) \right) \rightarrow \Box \neg \mathcal{J}_n^{\alpha^0}$$

with $\alpha \in 2^{\leq N}$ and $n < N$.

All the formulas of the first form are forced at the root of \mathcal{K} because \mathcal{K} is very high. Suppose that the antecedent of the formula

$$\left(\bigwedge_{1 \leq j \leq \text{lh}(\alpha)} (p_j \leftrightarrow (\alpha)_j) \right) \rightarrow \Box \neg \mathcal{J}_n^{\alpha^0}$$

is forced at the root of \mathcal{K} (we then have $\alpha \subseteq \delta$ and hence $\alpha \in T$ because T is efflorescent) and that this formula is in $C_{\alpha^0}^N$. This indicates that there exists a $j \in \omega$ s.t. $R(\alpha^-, n, j)$. Since $\alpha \in T$ we have that there exists an $i \in \omega$ s.t. $R(\alpha, n, i)$ and so the formula $\neg \mathcal{J}_n^{\alpha^0}$ is also in $C_{\alpha^0}^N$. As was shown above $\neg \mathcal{J}_n^{\alpha^0}$ holds in \mathcal{K} and therefore

$$\left(\bigwedge_{1 \leq j \leq \text{lh}(\alpha)} (p_j \leftrightarrow (\alpha)_j) \right) \rightarrow \Box \neg \mathcal{J}_n^{\alpha^0}$$

also does.

In view of Lemmas 5.14, 5.16, 5.22 and 5.24 these arguments prove the strong disjunction property of F/ε and that each element of T is in the admissibility tree of μ .

Next pick a string ε not in T . Let us see that the formula $\bigwedge \{ p_j \leftrightarrow (\varepsilon)_j \mid 1 \leq j \leq \text{lh}(\varepsilon) \}$ is not an admissible element

of F/\mathcal{C} . Since T is efflorescent there exists a string $\delta \subseteq \varepsilon$ s.t. $\delta^- \in T$ and $\delta \text{ non } \in T$ and we shall be done once we show $\bigwedge \{ p_j \leftrightarrow (\delta)_j \mid 1 \leq j \leq \text{lh}(\delta) \}$ not to be admissible.

By the choice of $R(\alpha, x, y)$ there exists an $N \in \omega$ s.t. $\exists j R(\delta^-, N, j)$ and $\forall i \neg R(\delta, N, i)$. By the construction of \mathcal{C}_{δ^0} we then have that the formula $\neg \mathcal{F}_N^{\delta^0}$ will never get in \mathcal{C} and in F/\mathcal{C} one has

$$\left(\bigwedge_{1 \leq j \leq \text{lh}(\delta)} (p_j \leftrightarrow (\delta)_j) \right) \rightarrow \Box \neg \mathcal{F}_N^{\delta^0} = \top$$

Fix an arbitrary $M \in \omega$. Take the model $\mathcal{F}_N^{\delta^0}$ and extend the forcing relation at the nodes of this model to the tuple (p_1, \dots, p_M) in an arbitrary way. C_{M-1} holds in this model because by Claim 1 it forces $\neg \mathcal{F}_n^\alpha$ unless $\alpha = \delta^0$ and $n = N$, the formula $\neg \mathcal{F}_N^{\delta^0}$ is not a conjunct of C_M and by Claim 2 the model forces $\Box \mathcal{F}_n^\alpha$, all $\alpha \in 2^{<\omega}$, $n \in \omega$.

Thus we have shown that $C_M \not\vdash_L \neg \mathcal{F}_N^{\delta^0}$ for no $M \in \omega$ and therefore

$$\neg \mathcal{F}_N^{\delta^0} \neq \top$$

in F/\mathcal{C} . Thus δ is not in the admissibility tree of μ q.e.d. \blacksquare

9.5. Lemma. *Let T be a Σ_2 -sound theory. Then the admissibility tree of the gödelnumbering of $\mathcal{D}_T^{\Delta_2}$ enjoys an infinite Δ_2^0 branch.*

Proof. Each sentence which is Δ_2 over T may be thought of as a pair (σ, π) of sentences σ in Σ_2 and π in Π_2 s.t. $T \vdash \sigma \leftrightarrow \pi$. By our assumption on T we have that σ is true if and only if π is true. Also note that if a Δ_2 sentence is true in the above sense then it corresponds to an admissible element of $\mathcal{D}_T^{\Delta_2}$ because if $T \vdash \sigma \rightarrow \text{Pr}(\bar{\varphi})$ then $\text{Pr}(\bar{\varphi})$ is true and hence $T \vdash \varphi$. Now the truth of Σ_2 sentences as well as falsity of Π_2 sentences is verifiable by a Turing machine with an oracle for $0'$ and hence the set of true sentences Δ_2 over T is Δ_2^0 . It clearly gives

rise to an infinite Δ_2^0 branch in the admissibility tree of $\mathfrak{D}_T^{\Delta_2}$. ■

9.6. Lemma. *There exists an efflorescent Π_2^0 tree with no infinite Σ_2^0 branch.*

Proof. Left as an exercise for the reader. Alternatively, the reader may check that if $\exists x \forall y D(\alpha, x, y, z)$ is a Σ_2^0 predicate on \perp - \top -strings α universal in z with $D(\alpha, x, y, z)$ decidable then the set of strings δ satisfying

$$\forall z \forall \alpha (\text{lh}(\alpha) = z + 1 \wedge \alpha \subseteq \delta \rightarrow$$

$$\rightarrow. \forall x (\exists y \neg D(\alpha, x, y, z) \vee \exists \beta (\text{lh}(\beta) = \text{lh}(\alpha) \wedge$$

$$\wedge. \alpha < \beta \rightarrow \exists w \leq x \forall v D(\beta, w, v, z) \wedge. \beta < \alpha \rightarrow \exists w < x \forall v D(\beta, w, v, z)))$$

is such a tree. ■

To give an example of a positive numeration μ of a diagonalizable algebra with the strong disjunction property which is not embeddable into $\mathfrak{D}_T^{\Delta_2}$ for a Σ_2 -sound T recursively w.r.t. μ it suffices to take a positive numeration $\mu : \omega \rightarrow \mathfrak{D}$ whose admissibility tree is the tree T constructed in Lemma 9.6. For if there existed an embedding of \mathfrak{D} into $\mathfrak{D}_T^{\Delta_2}$ recursive w.r.t. μ then by Lemma 9.5 we could single out an infinite Δ_2^0 branch in T which is not there.

9.7. Corollary. *Let T be a Σ_2 -sound theory. There exists no recursive (nor even recursive in $0'$) embedding of $\mathfrak{D}_T^{\Delta_0(\Sigma_2)}$ into $\mathfrak{D}_T^{\Delta_2}$.* ■

In connection with Corollary 9.7 we should mention a result of Pour-El & Kripke [37] which shows that if the diagonalizable structures of $\mathfrak{D}_T^{\Delta_2}$ and $\mathfrak{D}_T^{\Delta_0(\Sigma_2)}$ are forgotten then the underlying Boolean algebras are recursively isomorphic.

be the formula provided by Lemma 5.28 s.t.

- (i) $D_{i+1}(\omega+x) \vdash_{\mathbf{L}} D_{i+1}(x)$;
- (ii) $D_{i+1}(\omega+x) \vdash_{\mathbf{L}} D_i(\omega+x)$;
- (iii) $D_{i+1}(\omega+x)$ is apt ;
- (iv) $\Box^+ D_{i+1}(\omega+x)$ is i -conservative over $\Box^+ D_i(\omega+x)$ and
- (v) $D_{i+1}(\omega+x)$ supports every formula in $\mathbf{F}^0(i)$ that $D_i(\omega+x)$ does.

(Note that, unlike x , the i in $D_i(x)$ is an index rather than a free variable.)

10.2. Lemma. For each $i \in \omega$ s.t. $i > 0$ there holds

- (a) $\mathbf{I}\Sigma_1 \vdash \forall x \in \omega \forall y \in \omega (x \leq y \rightarrow D_i(y) \vdash_{\mathbf{L}} D_i(x))$
- (b) $\mathbf{I}\Sigma_1 \vdash \forall x \in \omega D_i(\omega+x) \vdash_{\mathbf{L}} D_i(x)$
- (c) $\mathbf{I}\Sigma_1 \vdash \forall x \in \omega D_{i+1}(\omega+x) \vdash_{\mathbf{L}} D_i(\omega+x)$
- (d) $\mathbf{I}\Sigma_1 \vdash \forall x \in \omega \cdot 2$ " $\Box^+ D_{i+1}(x)$ is i -conservative over $\Box^+ D_i(x)$ "
- (e) $\mathbf{I}\Sigma_1 \vdash \forall x \in \omega$ " $D_i(\omega+x)$ is apt"
- (f) $\mathbf{I}\Sigma_1 \vdash \forall x \in \omega \forall \alpha \in \mathbf{F}^0(\bar{i})$ (" α supports $D_i(\omega+x)$ " \rightarrow
 \rightarrow " α supports $D_{i+1}(\omega+x)$ ")
- (g) For each $y \in \omega$ there exists an $x \in \omega$ s.t. $D_i(x) \vdash_{\mathbf{L}} A_i(y)$.
- (h) For each $x \in \omega$ there exists a $y \in \omega$ s.t. $D_i(x) = A_i(y)$.

Proof. (a) and (h) are easy.

To prove clause (d) for the case $x \in \omega$ it suffices to notice that if a formula A is i -conservative over a formula B then it also is i -conservative over $B \wedge C$ for any formula C .

Claim. For all $i \in \omega$ $\mathbf{I}\Sigma_1$ proves that for each $x \in \omega$

- (i) " $D_i(\omega+x)$ is apt"
- (ii) $D_i(\omega+x) \vdash_{\mathbf{L}} D_i(x)$
- (iii) " $\Box^+ D_{i+1}(x)$ is i -conservative over $\Box^+ D_i(\omega+x)$ "
- (iv) "The premises of (7) are satisfied for i "

The Claim is proved by induction on i . For $i = 0$ this is

clear since $\Box^+ D_1(x)$ has to be conservative by (iii) of (4) whence (iii) and (iv) of the Claim follow.

Suppose that $i > 0$ and that the Claim holds for $i - 1$. Then by (iv) of the induction hypothesis we have that $D_i(\omega+x)$ is apt, $D_i(\omega+x) \vdash_{\mathbf{L}} D_i(x)$ and since by the ω -part of (d) one has that $\Box^+ D_{i+1}(x)$ is i -conservative over $\Box^+ D_i(x)$ this implies that $\Box^+ D_{i+1}(x)$ is also i -conservative over $\Box^+ D_i(\omega+x)$. Thus the Claim is proved.

(b) and (e) are direct consequences of the Claim.

Clauses (c) and (f) and clause (d) for $x = \omega + y$ follow from (iv) of the Claim combined with (iii), (v) and (iv) of (7) respectively.

(g) We proceed by induction on i . Suppose there exists an $y \in \omega$ s.t. $D_{i+1}(x) \vdash_{\mathbf{L}} A_{i+1}(y)$ for no $x \in \omega$. Pick the minimal such y . Then $\lim_{x \rightarrow \infty} k_{i+1}(x) = y$ and $\lim_{x \rightarrow \infty} D_{i+1}(x) = A_{i+1}(z)$ for some $z < y$. Consider the formula $A_{i+1}(z) \wedge A_{i+1}(y)$. Let C be the formula in $\mathbf{F}(i)$ s.t. $\vdash_{\mathbf{L}} \Box^+(A_{i+1}(z) \wedge A_{i+1}(y)) \rightarrow C$ and $\vdash_{\mathbf{L}} C \rightarrow D$ whenever $\vdash_{\mathbf{L}} \Box^+(A_{i+1}(z) \wedge A_{i+1}(y)) \rightarrow D$ and $D \in \mathbf{F}(i)$ (here we use the Uniform Craig Interpolation Lemma). Clearly $\vdash_{\mathbf{L}} C \leftrightarrow \Box^+ C$. Since $\{A_i(m)\}_{m \in \omega}$ and $\{A_{i+1}(m)\}_{m \in \omega}$ enumerate the relations of one and the same diagonalizable algebra we have that for some $w \in \omega$ $A_i(w) \vdash_{\mathbf{L}} C$. Therefore by the induction hypothesis there exists a $v \in \omega$ s.t. $D_i(v) \vdash_{\mathbf{L}} C$. By the choice of C the formula $\Box^+(A_{i+1}(z) \wedge A_{i+1}(y))$ is i -conservative over $\Box^+ C$ and hence over $\Box^+ D_i(w)$ for all $w \geq v$. Now formulas \mathbf{L} -equivalent to $A_{i+1}(z) \wedge A_{i+1}(y)$ occur unboundedly often in $\{A_{i+1}(m)\}_{m \in \omega}$ and, as we have shown, infinitely many such occurrences satisfy (i) - (iii) of (4). Thus we have reached a contradiction and therefore proved (f). ■

We now define a sequence of Δ_0 functions $\{g_i(\cdot)\}_{i \in \omega - \{0\}}$:

(8)

$g_{i+1}(x) = z$ if (i) $z < x$;

(ii) $z < \Delta$;

(iii) there exists a formula $A \in \mathbf{F}(i+1)$ s.t.

Prf $(z, \ell_{i+1} * \overline{\neg A} \rightarrow \exists y (0 R h_{i+1}(y) R \ell_{i+1}))$

and $(\bigwedge_{0 < j \leq i+1} D_j(g_j(z)))$ non $\vdash_L A$,

or there exists a j s.t. $0 < j \leq i+1$ and $g_j(x) = z$;

(iv) z is minimal among those satisfying (i) - (iii)

= x if $x < \delta$ and no z satisfying (i) - (iii) exists

= $\omega + \delta$ if $x \geq \delta$ and no z satisfying (i) - (iii) exists

10.3. Lemma. For each $i \in \omega$ s.t $i > 0$ $I\Sigma_1$ proves

(a) $\forall x g_i(x) \geq g_{i+1}(x)$

(b) $\forall x$ " $\square^+ D_{i+1}(g_{i+1}(x))$ is \bar{i} -conservative over $\square^+ D_i(g_i(x))$ "

(c) $\forall x \forall y (x \leq y \rightarrow D_i(g_i(y)) \vdash_L D_i(g_i(x)))$

(d) $\forall x = \delta \forall y D_i(\omega+x) \vdash_L D_i(g_i(y))$

(e) $\forall x \forall A \in F(\bar{i})$ $(\text{Prf } (x, \ell_i * \overline{\neg A} \rightarrow \exists y (0 R h_i(y) R \ell_i)) \wedge$
 $\wedge (\bigwedge_{0 < j \leq i} D_j(g_j(x)))$ non $\vdash_L A \rightarrow \forall y \geq x g_i(y) = g_i(x)$)

(f) $\forall x < \delta (g_i(x) \neq x \rightarrow$

$\rightarrow \exists A \in F(\bar{i})$ $(\text{Pr } (\ell_i * \overline{\neg A} \rightarrow \exists y (0 R h_i(y) R \ell_i)) \wedge$
 $\wedge \forall z D_i(g_i(z))$ non $\vdash_L A) \vee \bigwedge_{0 < j < i} g_j(x) \neq x$)

(g) $\forall x = \delta (g_i(x) \neq \omega + x \rightarrow$

$\rightarrow \exists A \in F(\bar{i})$ $(\text{Pr } (\ell_i * \overline{\neg A} \rightarrow \exists y (0 R h_i(y) R \ell_i)) \wedge$
 $\wedge \forall z D_i(g_i(z))$ non $\vdash_L A) \vee \bigwedge_{0 < j < i} g_j(x) \neq \omega + x$)

Proof. (a) is obvious.

(b) is proved by induction on x using the fact that if a formula A is i -conservative over a formula B then A is i -conservative over any formula that is stronger than B .

(c) - (g) are proved in the same way as corresponding clauses of Lemma 6.7. ■

Next we write down a sequence of Solovay functions.

$$(9) \quad h_{i+1}(0) = 0$$

(10)

$$h_{i+1}(x+1) = a \quad \text{if (i) } a \in M_{i+1} ;$$

$$\text{(ii) } h_i(x) \neq h_i(x+1) ;$$

$$\text{(iii) } h_{i+1}(x) R a ;$$

$$\text{(iv) } a \triangleleft h_i(x+1) ;$$

$$\text{(v) if } h_{i+1}(x) = 0 \text{ then } a \neq D_{i+1}(g_{i+1}(x)) ;$$

$$\text{(vi) if } h_{i+1}(x) = 0 \text{ then for each } b \text{ satisfying (i) - (v) in place of } a \text{ one has}$$

$$\forall z \leq x \quad (\exists A \in F(\overline{i+1})) \quad (\text{Prf}(z, \ell_{i+1} * \overline{\neg A} \rightarrow \exists y (0 R h_{i+1}(y) R \ell_{i+1})) \wedge$$

$$\wedge \left(\bigwedge_{0 < j \leq i+1} D_j(g_j(z)) \right) \text{ non } \vdash_L A \wedge b \text{ non } * A) \rightarrow \exists w \leq z \exists A \in F(\overline{i+1})$$

$$(\text{Prf}(w, \ell_{i+1} * \overline{\neg A} \rightarrow \exists y (0 R h_{i+1}(y) R \ell_{i+1})) \wedge$$

$$\wedge \left(\bigwedge_{0 < j \leq i+1} D_j(g_j(w)) \right) \text{ non } \vdash_L A \wedge a \text{ non } * A)$$

$$\text{(vii) if } h_{i+1}(x) \neq 0 \text{ then for each } b \text{ satisfying (i) - (v) in place of } a \text{ one has}$$

$$\forall z \leq x \quad (\text{Prf}(z, \ell_{i+1} = \overline{b} \rightarrow \exists y (\overline{h_{i+1}(x)} R h_{i+1}(y) R \overline{b})) \rightarrow$$

$$\rightarrow \exists w \leq z \text{ Prf}(w, \ell_{i+1} = \overline{a} \rightarrow \exists y (\overline{h_{i+1}(x)} R h_{i+1}(y) R \overline{a})))$$

$$\text{(viii) } a \text{ is minimal among those } c \text{ that satisfy}$$

$$\text{(i) - (vii) in place of } a$$

$$= h_{i+1}(x) \quad \text{if no } a \text{ satisfying (i) - (vii) exists}$$

$$(11) \quad \ell_{i+1} = \lim_{x \rightarrow \infty} h_{i+1}(x) \quad \text{if } h_{i+1} \text{ reaches a limit}$$

$$= 0 \quad \text{otherwise}$$

10.4. Lemma. For each $i, j \in \omega$ s.t. $i < j$ $I\Sigma_1$ proves

- (a) $\forall x \forall y (x \leq y \rightarrow h_i(x) = h_i(y) \vee h_i(x) R h_i(y))$
- (b) $\forall x h_{i+1}(x) \triangleleft h_i(x)$
- (c) $\forall x h_j(x) \triangleleft h_i(x)$
- (d) $\ell_i = \lim_{x \rightarrow \infty} h_i(x)$
- (e) $\forall x (h_i(x) R \ell_i \vee h_i(x) = \ell_i)$

(f) $l_j \triangleleft l_i$

Proof. We only prove (b) and that, as usual, by induction on x .

Assume $h_{i+1}(x) \triangleleft h_i(x) \neq h_i(x+1)$. The case when $h_i(x) \neq 0$ is treated as in § 6. So let $h_i(x) = h_{i+1}(x) = 0$.

We have to show that there exists a node $a \in M_{i+1}$ s.t. $a \triangleleft h_i(x+1)$ and $a \vDash D_{i+1}(g_{i+1}(x))$. Assuming there is no such a we have by Lemmas 3.3(d) and 3.7(d) that

$$\vdash_{\mathbf{L}} \Box^+ D_{i+1}(g_{i+1}(x)) \rightarrow \neg \Psi_{h_i(x+1)}$$

Now from Lemma 10.3(b) we know that $\Box^+ D_{i+1}(g_{i+1}(x))$ is i -conservative over $\Box^+ D_i(g_i(x))$ and hence

$$\vdash_{\mathbf{L}} \Box^+ D_i(g_i(x)) \rightarrow \neg \Psi_{h_i(x+1)}$$

But note that since $0 = h_i(x) \neq h_i(x+1)$ we have by (v) of (9) that $h_i(x+1) \vDash D_i(g_i(x))$ ergo

$$\vdash_{\mathbf{L}} \Psi_{h_i(x+1)} \rightarrow \Box^+ D_i(g_i(x))$$

by Lemma 3.3(d). The contradiction proves (b). ■

10.5. Lemma. For each $i \in \omega$ $\text{I}\Sigma_1 + \ell \neq 0$ proves

(a) $\Pr(\overline{l_i} R l_i)$

(b) $\forall a \in M_{\overline{i}} (l_i R a \rightarrow \neg \Pr(\overline{l_i} = \overline{a} \rightarrow \exists y (\overline{l_i} R h_i(y) R \overline{a})))$

(c) $\forall a \in M_{\overline{i}} (l_i R a \rightarrow \neg \Pr(\overline{l_i} \neq \overline{a}))$

Proof. Confer Lemma 6.6. ■

10.6. Lemma. For each $i \in \omega$ s.t. $i > 0$ $\text{I}\Sigma_1$ proves

(a) $l \neq 0 \rightarrow \forall x (h_i(x) = 0 \wedge \rightarrow l_i \vDash \bigwedge_{0 < j \leq i} D_j(g_j(x)))$

(b) $l \neq 0 \rightarrow$

$\rightarrow \forall x \forall A \in \mathbf{F}(\overline{i}) (h_i(x) = 0 \wedge (\bigwedge_{0 < j \leq i} D_j(g_j(x))) \vdash_{\mathbf{L}} A \rightarrow \cdot l_i \vDash A)$

(c) $l = 0 \rightarrow$

$$\rightarrow \forall A \in F(\bar{i}) \quad \overline{\text{Pr} (\ell_i \neq \bar{A} \rightarrow \exists y (0 R h_i(y) R \ell_i))} \rightarrow \\ \rightarrow \exists x \left(\bigwedge_{0 < j \leq i} D_j(g_j(x)) \right) \vdash_{\mathbf{L}} A$$

(d) $\ell = 0 \rightarrow$

$$\rightarrow \forall A \in F(\bar{i}) \quad \overline{\text{Pr} (\ell \neq 0 \rightarrow \ell_i \neq \bar{A})} \rightarrow \exists x \left(\bigwedge_{0 < j \leq i} D_j(g_j(x)) \right) \vdash_{\mathbf{L}} A$$

Proof. (a) We prove this by induction on i . For $i = 0$ the claim is trivial. If $h_{i+1}(x) = 0$ then by (v) of (10) and by Lemma 10.3(c) h_{i+1} can only jump to a node at which $D_{i+1}(g_{i+1}(x))$ holds and therefore $\ell_{i+1} \neq D_{i+1}(g_{i+1}(x))$. On the other hand by Lemma 10.4(f) we have $\ell_{i+1} \triangleleft \ell_i$ and $\ell_i \neq \bigwedge_{0 < j \leq i} D_j(g_j(x))$ by the induction hypothesis, ergo $\ell_{i+1} \neq \bigwedge_{0 < j \leq i} D_j(g_j(x))$. Thus $\ell_{i+1} \neq \bigwedge_{0 < j \leq i+1} D_j(g_j(x))$ and we have carried out the induction step.

(b) follows from (a).

(c) Once again, induction on x . Let $i = 0$ and let A be a formula in $F(0)$ s.t. $\text{non} \vdash_{\mathbf{L}} A$. Suppose

$$\overline{\text{Pr} (\ell_0 \neq \bar{A} \rightarrow \exists y (0 R h_0(y) R \ell_0))}$$

Then there exists a node $a \in M_0$ s.t. $a \text{ non} \neq A$ and we have by Lemma 6.2(c)

$$\overline{\text{Pr} (\ell_0 = \bar{a} \rightarrow \exists y (0 R h_0(y) R \bar{a}))}$$

But this combined with $\ell = 0$ contradicts Lemma 6.2(d).

Now we let $i > 0$, assume that the claim holds for i , deny it for $i + 1$ and look for a contradiction. Let $x \in \omega$ be s.t.

$$\overline{\text{Prf} (x, \ell_{i+1} \neq \bar{A} \rightarrow \exists y (0 R h_{i+1}(y) R \ell_{i+1}))}$$

and $\left(\bigwedge_{0 < j \leq i+1} D_j(g_j(x)) \right) \text{ non} \vdash_{\mathbf{L}} A$ for some $A \in F(i)$. By the Δ_0 least number principle we can assume that x is minimal among the proofs of this kind. We then have by (8) that $\forall y \geq x \ g_{i+1}(y) = g_{i+1}(x)$ so $D_{i+1}(g_{i+1}(z)) \vdash_{\mathbf{L}} A$ for no z .

Consider the weakest formula B in $F(i)$ s.t.

$$\vdash_{\mathbf{L}} B \rightarrow \square^+ D_{i+1}(g_{i+1}(x)) \rightarrow A$$

which exists by the Uniform Craig Interpolation Lemma. Let a be an arbitrary node of M_i forcing $\neg B$. We show that there is a node $b \in M_{i+1}$ s.t. $b \triangleleft a$ and $b \Vdash \neg A \wedge \Box^+ D_{i+1}(g_{i+1}(x))$. For if this were not the case then $e \Vdash \Box^+ D_{i+1}(g_{i+1}(x)) \rightarrow A$ for each $e \in M_{i+1}$ s.t. $e \triangleleft a$ and therefore by Lemmas 3.3(d) and 3.7(d)

$$\Vdash_{\mathbf{L}} \Psi_a \rightarrow \Box^+ D_{i+1}(g_{i+1}(x)) \rightarrow A$$

whence $\Vdash_{\mathbf{L}} \Psi_a \rightarrow B$ because B is the weakest formula in $\mathbf{F}(i)$ implying $\Box^+ D_{i+1}(g_{i+1}(x)) \rightarrow A$. But this contradicts the assumption $a \Vdash \neg B$.

Reason in $\text{I}\Sigma_1$. If h_i jumps from 0 to a node forcing $\neg B$ then by (vi) of (10) h_{i+1} will have to jump directly to a node forcing $\neg A$ (this is, because conditions (i) - (v) of (10) will obviously be satisfied). So outside $\text{I}\Sigma_1$ we have

$$\Pr (\ell_i \Vdash \neg B \rightarrow \exists y (0 R h_i(y) R \ell_i) \vee \ell_{i+1} \Vdash \neg A)$$

which together with Lemma 10.4(b) and the assumption on A gives

$$\Pr (\ell_i \Vdash \neg B \rightarrow \exists y (0 R h_i(y) R \ell_i))$$

Now note that we have $(\bigwedge_{0 < j \leq i} D_j(g_j(x))) \Vdash_{\mathbf{L}} B$ for no $z \in \omega$ because then we would have

$$(\bigwedge_{0 < j \leq i} D_j(g_j(x))) \Vdash_{\mathbf{L}} \Box^+ D_{i+1}(g_{i+1}(x)) \rightarrow A$$

and hence $(\bigwedge_{0 < j \leq i+1} D_j(g_j(x))) \Vdash_{\mathbf{L}} A$ contrary to assumptions. The situation in which B has found itself can now easily be seen to contradict the induction hypothesis and so (c) is proved.

(d) follows from (c). ■

Next we deal with 0 (in $\text{I}\Sigma_1 + \sigma$). The formulas $\text{Adm}_i(\cdot)$ are defined for $i \in \omega - \{0\}$:

$$(12) \quad \text{Adm}_i(\alpha) \equiv \text{lh}(\alpha) \leq \bar{i} \wedge \text{"}\alpha \text{ supports } D_i(\omega + \delta)\text{"}$$

10.7. Lemma. For each $i \in \omega - \{0\}$ $\text{I}\Sigma_1 + \sigma$ proves

$$(a) \quad \text{Adm}_i(\bar{\Lambda})$$

$$(b) \quad \forall \alpha \forall \beta (\text{Adm}_i(\beta) \wedge \alpha \subseteq \beta \rightarrow \text{Adm}_i(\alpha))$$

- (c) $\forall \alpha (\text{Adm}_i(\alpha) \rightarrow \exists \beta (\text{lh}(\beta) = \bar{i} \wedge \alpha \subseteq \beta \wedge \text{Adm}_i(\beta)))$
(d) $\exists \alpha (\text{lh}(\alpha) = \bar{i} \wedge \text{Adm}_i(\alpha))$
(e) $\forall \alpha (\text{lh}(\alpha) \leq \bar{i} \rightarrow \text{Adm}_i(\alpha) \leftrightarrow \text{Adm}_{i+1}(\alpha))$

Proof. (a) - (d) are proved as in Lemma 6.9.

(e) (\rightarrow) is a straightforward consequence of Lemma 10.2(f).

(\leftarrow) Since by Lemma 10.2(c) and (d) $D_{i+1}(\omega+\delta) \vdash_{\mathbf{L}} D_i(\omega+\delta)$ and $\square^+ D_{i+1}(\omega+\delta)$ is i -conservative over $\square^+ D_i(\omega+\delta)$ we have that $\square^+ D_i(\omega+\delta)$ is the (N, i) -shadow of $\square^+ D_{i+1}(\omega+\delta)$ for some large enough $N \in \omega$. Now by Lemma 5.26 and Corollary 5.27 we have that $D_i(\omega+\delta)$ is supported by each string that supports $D_{i+1}(\omega+\delta)$. ■

$$(13) \quad \text{Adm}_i^+(\alpha) \equiv \text{Adm}_i(\alpha) \wedge \forall \beta < \alpha (\text{lh}(\alpha) = \text{lh}(\beta) \rightarrow \neg \text{Adm}_i(\beta))$$

10.8. Lemma. For each $i \in \omega$ s.t. $i > 0$ $\text{I}\Sigma_1 + \sigma$ proves

- (a) $\exists! \alpha (\text{lh}(\alpha) = \bar{i} \wedge \text{Adm}_i^+(\alpha))$
(b) $\forall \alpha (\text{Adm}_i^+(\alpha) \rightarrow \exists \beta (\text{lh}(\beta) = i \wedge \alpha \subseteq \beta \wedge \text{Adm}_i^+(\beta)))$
(c) $\forall \alpha \forall \beta (\text{Adm}_i^+(\alpha) \wedge \text{Adm}_i^+(\beta) \wedge \text{lh}(\alpha) \leq \text{lh}(\beta) \rightarrow \alpha \subseteq \beta)$
(d) $\forall \alpha (\text{lh}(\alpha) \leq \bar{i} \rightarrow \text{Adm}_i^+(\alpha) \leftrightarrow \text{Adm}_{i+1}^+(\alpha))$
(e) $\forall \alpha \forall \beta (\text{Adm}_i^+(\alpha) \wedge \text{Adm}_{i+1}^+(\beta) \wedge \text{lh}(\alpha) \leq \text{lh}(\beta) \rightarrow \alpha \subseteq \beta)$

Proof. Clauses (a) - (c) are proved in full analogy with those of Lemma 6.10.

(d) follows from Lemma 10.7(e).

(e) follows from (c) and (d). ■

$$(14) \quad \text{T}_i(A) \equiv \exists k \exists \lambda \in \mathbf{F}^0(\bar{i}+k) \exists B_1, \dots, B_k \in \mathbf{F}(\bar{i}) \exists \alpha$$

$$(\text{lh}(\alpha) = \bar{i} + k \wedge \lambda((\alpha)_{\bar{i}}, \dots, (\alpha)_{\bar{i}+k}) = \tau \wedge$$

$$\wedge A = \lambda(p_1, \dots, p_i, \square B_1, \dots, \square B_k) \wedge \forall j \leq \bar{i} (j \neq 0 \rightarrow$$

$$\rightarrow (\alpha)_j = \tau \leftrightarrow \exists \beta (\text{lh}(\beta) = \bar{i} \wedge \text{Adm}_i^+(\beta) \wedge (\beta)_j = \tau)) \wedge$$

$$\wedge \forall j \leq k (j \neq 0 \rightarrow (\alpha)_{\bar{i}+j} = \tau \leftrightarrow D_i(\omega+\delta) \vdash_{\mathbf{L}} B_j))$$

10.9. Lemma. For each $i, j \in \omega$ s.t. $0 < i \leq j$ $\text{I}\Sigma_1 + \sigma$ proves

- (a) $T_j(\overline{p_i}) \leftrightarrow \exists \beta \text{ (lh}(\beta) = \overline{j} \wedge \text{Adm}_j^+(\beta) \wedge (\beta)_{\overline{i}} = \tau)$
- (b) $\forall A \in F(\overline{i}) \text{ (} T_i(\Box A) \leftrightarrow D_i(\omega + \Delta) \vdash_{\mathbf{L}} A \text{)}$
- (c) $\forall k \forall \lambda \in F^0(k) \forall B_1, \dots, B_k \in F \text{ (} T_i(\lambda(B_1, \dots, B_k)) \leftrightarrow$
 $\leftrightarrow \exists \alpha \text{ (lh}(\alpha) = k \wedge \lambda((\alpha)_1, \dots, (\alpha)_k) = \tau \wedge$
 $\wedge \forall l \leq k \text{ (} l \neq 0 \rightarrow (\alpha)_l = \tau \leftrightarrow T_i(B_l) \text{))}$
- (d) $\forall A \in F(\overline{i}) \text{ (} T_i(A) \leftrightarrow T_{i+1}(A) \text{)}$
- (e) $\forall A \in F(\overline{i}) \text{ (} T_i(A) \leftrightarrow T_j(A) \text{)}$

Proof. (a) - (c) are proved as in Lemma 6.11.

(d) For A a propositional letter in $F(i)$ we have this by Lemma 10.8(d) and (e) and (a) of the present lemma. For A of the form $\Box B$ this follows from (b) combined with Lemma 10.2(c) and (d). Finally, one executes induction on the structure of A with the help of (c).

(e) follows from (d). ■

10.10. Lemma. For each $i \in \omega$ s.t. $i > 0$ $\text{I}\Sigma_1 + \sigma$ proves

- (a) $\forall \alpha \text{ (lh}(\alpha) \leq \overline{i} \rightarrow T_i(\bigwedge_{1 \leq j \leq \text{lh}(\alpha)} (p_j \leftrightarrow (\alpha)_j)) \leftrightarrow \text{Adm}_i^+(\alpha) \text{)}$
- (b) $\forall A \in F(\overline{i}) \forall B \in F(\overline{i}) \text{ (} T_i(\Diamond A \wedge \Box B) \rightarrow T_i(\Diamond(A \wedge \Box B)) \text{)}$
- (c) $\forall A \in F(\overline{i}) \text{ (} T_i(\Box A) \rightarrow T_i(A) \text{)}$

Proof. Analogous to Lemma 6.12. ■

Now we map $\{p_i\}_{i \in \omega - \{0\}}$ (and hence the whole of F) into \mathfrak{D}_T :

$$(15) \quad p_i^\circ \equiv l \neq 0 \wedge l_i * \overline{p_i} \vee l = 0 \wedge T_i(\overline{p_i})$$

10.11. Lemma. For each $i \in \omega$ s.t. $i > 0$ there holds

- (a) For each modal formula $A(p_1, \dots, p_i)$ one has

$$\text{I}\Sigma_1 \vdash l \neq 0 \rightarrow (A(p_1, \dots, p_i))^\circ \leftrightarrow l_i * \overline{A(p_1, \dots, p_i)}$$

- (b) $\text{I}\Sigma_1 \vdash \forall A \in F(\overline{i}) \text{ Pr } (l \neq 0 \rightarrow \overline{A^\circ \leftrightarrow l_i * \overline{A}})$

(note that $^\circ$ restricted to $F(i)$ is representable)

- (c) $\text{I}\Sigma_1 \vdash l = 0 \rightarrow$

$$\rightarrow \forall A \in \mathcal{F}(\bar{i}) \quad (\text{Pr } \overline{A^\circ}) \rightarrow \exists x \left(\bigwedge_{0 < j \leq i} D_j(g_j(x)) \right) \vdash_{\mathbf{L}} A)$$

$$(d) \quad \text{I}\Sigma_1 + \sigma \vdash \ell = 0 \rightarrow g_i(\Delta) = \omega + \Delta$$

(e) For each modal formula $A(p_1, \dots, p_i)$ one has

$$\begin{aligned} \text{I}\Sigma_1 + \sigma \vdash (A(p_1, \dots, p_i))^\circ &\leftrightarrow \\ &\leftrightarrow (\ell \neq 0 \wedge \ell_i \neq \overline{A(p_1, \dots, p_i)} \vee \ell = 0 \wedge T_i(\overline{A(p_1, \dots, p_i)})) \end{aligned}$$

Proof. Fairly similar to the proof of Lemma 6.13. When proving (d) use induction on i and Lemmas 10.3(g) and 10.6(c). When carrying out the induction step in (e) corresponding to a Boolean connective use Lemma 10.9(e). ■

10.12. Lemma. For each $i \in \omega$ s.t. $i > 0$

$$\mathbb{N} \models "g_i \text{ is the identity function}"$$

Proof. This is proved by induction on i with the help of Lemmas 10.3(f) and 10.6(c). ■

10.13. Lemma. If there holds $A(v_1, v_2, \dots) = \tau$ for $A(x_1, x_2, \dots)$ a diagonalizable polynomial then

$$\text{I}\Sigma_1 + \sigma \vdash (A(p_1, p_2, \dots))^\circ$$

Proof. Confer Lemma 6.15. ■

Finally, set

$$(16) \quad (v_i)^* \equiv p_i^\circ$$

and use Lemmas 10.6(d), 10.11(e) and 10.13 to finish the proof of Theorem 10.1. ■

§ 11. Arbitrary subalgebras. Σ_1 -sound theories

In this § we show that the straightforward attempt to generalize Theorem 7.1 to arbitrary subalgebras of diagonalizable algebras of Σ_1 -sound theories the way Theorem 10.1 generalizes Theorems 4.1 and 6.1 fails. We shall present an example of a locally positive diagonalizable algebra with the strong disjunction property which is not isomorphic to a subalgebra of \mathcal{D}_T for a Σ_1 -sound theory T . To do so we first need to list some definitions and facts from recursion theory.

11.1. Definition. Consider Turing machines with two different halting states 0 and 1. Such machines are said to be *0-1-Turing*. A 0-1-valued partial recursive function f is *computed* by a 0-1-Turing machine T if

$$f(n) = i \Leftrightarrow T \text{ halts in state } i \text{ on input } n$$

A mapping φ of ω onto the set of 0-1-valued partial recursive functions s.t. $\varphi_i(n)$ is a binary recursive function is called a *numbering* (of 0-1-valued partial recursive functions). A numbering is *acceptable* (cf. Rogers [38]) if there exists a total recursive function τ s.t. for each $i \in \omega$ $\tau(i)$ is a φ -index for the function computed by the i th 0-1-Turing machine (that is, $\varphi_{\tau(i)}$ is this function).

11.2. Definition. Let φ be an acceptable numbering. A binary partial recursive function $\Phi_i(n)$ is called a *Blum complexity measure* (for φ) (cf. Blum [10]) if

- (i) $\Phi_i(n)$ converges if and only if $\varphi_i(n)$ converges and
- (ii) the ternary relation $\Phi_i(n) \leq m$ is decidable.

In 11.3 - 11.5 we shall assume that Φ is a Blum complexity measure for an acceptable numbering φ .

We quote a theorem due to Blum which will help us to construct the desired diagonalizable algebra. Although in Blum [10] the theorem was claimed for acceptable numberings of all the partial recursive functions, an easy inspection of Blum's proofs

shows it to be also valid for acceptable numberings of 0-1-valued partial recursive functions.

11.3. Proposition (Blum [10]). *To each partial recursive function f there corresponds a 0-1-valued partial recursive function g s.t.*

- (i) $\text{dom } g = \text{dom } f$ and
- (ii) if i is a φ -index for g then $\Phi_i(n) \geq f(n)$ for all but finitely many $n \in \text{dom } f$. ■

From this theorem we infer an easy corollary.

11.4. Corollary. *Let X be an r.e. subset of ω . There exists a sequence of 0-1-valued partial recursive functions $\{g_i\}_{i \in \omega}$ s.t.*

- (i) for each $i \in \omega$ $\text{dom } g_i = X$ and
- (ii) for each partial recursive function f with $\text{dom } f = X$ there exists an $i \in \omega$ s.t. for each φ -index j for g_i one has $f(n) < \Phi_j(n)$ for all but finitely many $n \in X$.
(Clearly g_i can not be recursive in i). ■

11.5. Lemma. *Let X be a nonrecursive r.e. subset of ω and let the domain of φ_i be X . Then for each total recursive function f one has $f(n) \leq \Phi_i(n)$ for infinitely many $n \in X$.*

Proof. If this were not so then X could easily be shown decidable. ■

Define the modal formula $\#_n$ to be $\Box^{n+1} \perp \wedge \Diamond^n \top$. In this § we shall give way to an unmerciful intrusion of modal logic on arithmetic.

11.6. Definition. Let T be a theory of infinite credibility extent. Consider a particular numbering δ^T of 0-1-valued partial recursive functions which will take (gödelnumbers of) arithmetic sentences for indices:

$$\delta_\alpha^T(n) = 0 \quad \text{if} \quad T \vdash \#_{\frac{n}{\alpha}} \rightarrow \alpha$$

$$= 1 \quad \text{if } T \vdash \#_{\bar{n}} \rightarrow \neg \alpha$$

$$\text{divergent} \quad \text{if } T + \#_{\bar{n}} \text{ does not decide } \alpha$$

Further define

$$\Delta_{\alpha}^T(n) = p \quad \text{if } p \text{ is the least proof in } T \text{ of either } \#_{\bar{n}} \rightarrow \alpha \quad \text{or}$$

$$\#_{\bar{n}} \rightarrow \neg \alpha$$

$$\text{divergent} \quad \text{if } T + \#_{\bar{n}} \text{ does not decide } \alpha$$

11.7. Lemma. *If T is a theory of infinite credibility extent then*

- (a) δ^T is an acceptable numbering ;
- (b) Δ^T is a Blum complexity measure for δ^T .

Proof. (a) Note that for each $n \in \omega$ the theory $T + \#_{\bar{n}}$ is consistent. By Corollary 2 in Smoryński [43] or by the Uniform Dual Semi-Representability Theorem of Smoryński [45] there exists a formula $\sigma(y,x)$ s.t. for all $i, m \in \omega$

$$T \vdash \sigma(\bar{i}, \bar{m})$$

$$\text{iff for some } n \in \omega \quad T + \#_{\bar{n}} \vdash \sigma(\bar{i}, \bar{m})$$

$$\text{iff for all } n \in \omega \quad T + \#_{\bar{n}} \vdash \sigma(\bar{i}, \bar{m})$$

$$\text{iff the } i\text{th } 0\text{-}1\text{-Turing machine halts in state } 0 \text{ on input } m$$

and

$$T \vdash \neg \sigma(\bar{i}, \bar{m})$$

$$\text{iff for some } n \in \omega \quad T + \#_{\bar{n}} \vdash \neg \sigma(\bar{i}, \bar{m})$$

$$\text{iff for all } n \in \omega \quad T + \#_{\bar{n}} \vdash \neg \sigma(\bar{i}, \bar{m})$$

$$\text{iff the } i\text{th } 0\text{-}1\text{-Turing machine halts in state } 1 \text{ on input } m$$

Let $\tau(i)$ be the formula

$$\forall x (\#_x \rightarrow \sigma(\bar{i}, x))$$

Then $\tau(i)$ is a δ^T -index for the function computed by the i th 0-1-Turing machine because

$$T \vdash \forall x \forall y (\#_x \wedge \#_y \rightarrow x = y)$$

and therefore

$$T \vdash \#_{\bar{n}} \rightarrow \forall x (\#_x \rightarrow \sigma(\bar{i}, x)) \leftrightarrow \sigma(\bar{i}, \bar{n})$$

(b) is obvious. ■

11.8. Proposition. *Let T be a Σ_1 -sound theory. There exists a locally positive diagonalizable algebra \mathfrak{D} with the strong disjunction property which is not isomorphic to a subalgebra of \mathfrak{D}_T .*

Proof. As usual we take \mathfrak{D} to be the quotient of the free diagonalizable algebra F modulo a suitable set \mathcal{D} of its elements. The generators of F are $\{q\} \cup \{p_i\}_{i \in \omega}$. We now describe \mathcal{D} .

Let X be a nonrecursive r.e. subset of ω .

First, we put in \mathcal{D} all the formulas

$$\#_n \rightarrow q \quad \text{for } n \in X$$

Second, formulas

$$\Box(\#_n \rightarrow q) \rightarrow \Box(\#_n \rightarrow p_i) \vee \Box(\#_n \rightarrow \neg p_i) \quad \text{for all } i, n \in \omega$$

Third, with the help of Lemma 11.7 we fix a sequence $\{g_i\}_{i \in \omega}$ of 0-1-valued partial recursive functions for which the conditions of Corollary 11.4 hold with φ replaced by δ^T and Φ replaced by Δ^T . Put in \mathcal{D} the formulas

$$\#_n \rightarrow p_i \quad \text{for all } i, n \in \omega \text{ s.t. } g_i(n) = 0$$

and

$$\#_n \rightarrow \neg p_i \quad \text{for all } i, n \in \omega \text{ s.t. } g_i(n) = 1$$


Let D_N be the conjunction of all formulas in \mathcal{D} that neither speak about $\#_m$ for $m > N$ nor about p_i with $i > N$.

Clearly $A(q, p_0, \dots) = \tau$ in F/\mathcal{D} if and only if for some $N \in \omega$ one has $D_N \vdash_L A$.

It can be shown that if $n \notin X$ then neither $\#_n \rightarrow p_i = \tau$ nor $\#_n \rightarrow \neg p_i = \tau$ for any $i \in \omega$.

Now we show that for each $N \in \omega$ the formula D_N is apt.

Consider the (q, p_0, \dots, p_N) -model \mathcal{H}_N (the forcing relation is not indicated in the picture):



We do not care what propositional letters are forced at the bottom node. Each of the rest nodes forces $\#_n$ for a unique $n \leq N$. Call this node a_n .

Let q be forced precisely at those nodes a_n for which $n \in X$.

For each $n \leq N$ s.t. $n \in X$ and each $i \in \omega$ just one of the formulas

$$\#_n \rightarrow p_i \quad \text{and} \quad \#_n \rightarrow \neg p_i$$

is a conjunct of D_N . In the first case let $a_n * p_i$ and in the second case let $a_n * \neg p_i$. The forcing of letters p at the nodes a_n with $n \notin X$ is quite irrelevant.

We leave it to the reader to graft a pair of models in which D_N holds above the root of \mathcal{K}_N and to check that D_N holds in the resulting model.

This having been done, one can apply Lemmas 5.14 and 5.16 to see that D_N is apt and that \mathbf{F}/\mathcal{D} enjoys the strong disjunction property. Moreover, \mathbf{F}/\mathcal{D} is clearly locally positive.

It remains to prove that \mathbf{F}/\mathcal{D} is not embeddable into \mathfrak{D}_T .

We need three total recursive functions n , mp and sd (necessitation, modus ponens and strong disjunction) with the following properties:

- n is such that if p is a proof in T of a sentence α and $p \leq x$ then there exists a proof q in T of $\Box\alpha$ satisfying $q \leq n(x)$. Clearly we can stipulate

that $n(x) \geq x$ for each x ;

- mp is such that if p_1 and p_2 are proofs in T of sentences α and $\alpha \rightarrow \beta$ and $p_1, p_2 \leq x$ then there is a proof q in T of β satisfying $q \leq mp(x)$;
- sd is such that if p is a proof in T of a sentence of the form $\Box\alpha \vee \Box\beta$ and $p \leq x$ then there exists a proof q in T either of α or of β satisfying $q \leq sd(x)$.

Note that by Parikh's Theorem cited in § 1 or, alternatively, by Proposition 1.1 the function sd can not be provably recursive whereas the functions n and mp can, under any reasonable gödel-numbering, be chosen primitive recursive.

Suppose $*$ is a hypothetical embedding of F/\mathcal{D} into \mathcal{D}_T . We want a contradiction.

Since p_i^* is a δ^T -index for g_i and because the functions g_i were chosen in accordance with Corollary 11.4 there is a number $i \in \omega$ s.t.

$$sd \circ mp \circ n \circ \Delta_{q^*}^T(n) < \Delta_{p_i^*}^T(n)$$

for all but finitely many $n \in X$ (we have that for each $n \in X$ both the l.h.s. and the r.h.s. converge).

Let D be a total recursive function s.t. for each $n \in \omega$ $D(n)$ is a proof in T of

$$\Box(\#_{\bar{n}} \rightarrow q^*) \rightarrow \Box(\#_{\bar{n}} \rightarrow p_i^*) \vee \Box(\#_{\bar{n}} \rightarrow \neg p_i^*)$$

By Lemma 11.5 we have

$$D(n) \leq \Delta_{q^*}^T(n) \leq n \circ \Delta_{q^*}^T(n)$$

for infinitely many $n \in X$. Therefore for infinitely many $n \in X$ we have that there are proofs $\leq n \circ \Delta_{q^*}^T(n)$ of both formulas

$$\Box(\#_{\bar{n}} \rightarrow q^*) \quad \text{and} \quad \Box(\#_{\bar{n}} \rightarrow q^*) \rightarrow \Box(\#_{\bar{n}} \rightarrow p_i^*) \vee \Box(\#_{\bar{n}} \rightarrow \neg p_i^*)$$

Therefore for infinitely many $n \in X$ there is a proof $\leq mp \circ n \circ \Delta_{q^*}^T(n)$ of the formula

$$\Box(\#_{\bar{n}} \rightarrow p_i^*) \vee \Box(\#_{\bar{n}} \rightarrow \neg p_i^*)$$

and hence for infinitely many $n \in X$ there is a proof $\leq \text{sd} \circ \text{mp} \circ n \circ \Delta_{q^*}^T(n)$ of one of the formulas

$$\#_{\bar{n}} \rightarrow p_i^* \quad \text{and} \quad \#_{\bar{n}} \rightarrow \neg p_i^*$$

that is,

$$\Delta_{p_i^*}^T(n) \leq \text{sd} \circ \text{mp} \circ n \circ \Delta_{q^*}^T(n)$$

for infinitely many $n \in X$. But this can not be the case by our assumption on p_i^* . Thus we have obtained the required contradiction and have therefore completed the proof. ■

Roughly speaking, the reason for Proposition 11.8's holding true is that the strong disjunction property in locally positive diagonalizable algebras can, in a sense, be even less effective than in the diagonalizable algebras of Σ_1 -sound theories. A reason for this reason lies in the fact that the conjunction of two apt formulas is not necessarily apt. If one, starting from Theorem 7.1 were to attempt disproving Proposition 11.8 following the lines that led us from Theorems 4.1 and 6.1 to Theorem 10.1 one would meet difficulties in arranging recursive sequences of formulas $\{D_i(m)\}_{m \in \omega}$ in such a way that for each $i, m \in \omega$

- (i) $D_i(m+1) \vdash_{\mathbf{L}} D_i(m)$;
- (ii) $\Box^+ D_{i+1}(m)$ is i -conservative over $\Box^+ D_i(m)$ and
- (iii) $D_i(m)$ is apt

This happens because we can not guarantee that $D_{i+1}(m) \wedge D_i(m+1)$ is apt from the condition that both $D_{i+1}(m) \wedge D_i(m)$ and $D_i(m+1)$ are. In other words, the introduction of a stronger apt formula $D_i(m+1)$ may create new disjunction problems for the sequence D_{i+1} and since in locally positive algebras D_i knows nothing about D_{i+1} this situation is generally unavoidable. (These sentiments also explain how the proof of Proposition 11.8 came to the author's mind.)

So the problem of characterizing subalgebras of diagonalizable algebras of Σ_1 -sound theories T remains open. It appears

that a locally positive algebra can be embedded in \mathfrak{D}_T if the generators of \mathfrak{D} abstain from trying to bury each other in increasingly complex disjunction problems. Thus, the free product of a countable number of positive diagonalizable algebras with the strong disjunction property is always embeddable in \mathfrak{D}_T .

A weak kind of consolation is presented by Corollary 11.10. Proposition 11.9 was kindly pointed out to me by Professor Franco Montagna.

11.9. Proposition. *Let T_1 and T_2 be Σ_1 -sound theories. Then \mathfrak{D}_{T_1} is embeddable in \mathfrak{D}_{T_2} .*

Proof. A straightforward consequence of Theorem 7.1. ■

11.10. Corollary. *Diagonalizable algebras of all Σ_1 -sound theories possess one and the same collection of subalgebras.* ■

References

- [1] Z.Adamowicz. *A recursion-theoretic characterization of instances of $\text{B}\Sigma_n$ provable in $\Pi_{n+1}(\text{N})$* . Fundamenta Mathematicae 129 (1988) 231-236.
- [2] С.Н.Артёмов. *Арифметически полные модальные теории*. В кн.: *Семиотика и информатика*, выпуск 14. ВИНТИ, Москва 1979/80, 115-133 (English translation: American Mathematical Society Translations, Series 2, 135 (1987) 39-54).
- [3] С.Н.Артёмов. *О модальных логиках, аксиоматизирующих доказуемость*. Известия Академии Наук СССР, Серия математическая 49 (1985) 1123-1154 (English translation: Mathematics of the USSR - Izvestiya 27 (1986) 401-429).
- [4] С.Н.Артёмов. *О локальной табличности пропозициональных логик доказуемости*. В кн.: *Логические методы построения эффективных алгоритмов*. Калининский государственный университет 1986, 9-13.
- [5] Л.Д.Беклемишев. *О классификации пропозициональных логик доказуемости*. Известия Академии Наук СССР, Серия математическая 53 (1989) 915-943.
- [6] L.D.Beklemishev. *For whom the doorbell rings or the provability logic with a constant for an infinitely credent Π_1 sentence over PA*, Manuscript, Москва 1990.
- [7] F.Bellissima. *On the modal logic corresponding to diagonalizable algebra theory*. Bollettino della Unione Matematica Italiana, Serie V, 15-B (1978) 915-930.
- [8] C.Bernardi. *On the equational class of diagonalizable algebras (the algebraization of the theories which express Theor; VI)*. Studia Logica 34 (1975) 321-331.
- [9] C.Bernardi & M.Mirolli. *Hyperdiagonalizable algebras*. Algebra Universalis 21 (1985) 89-102.
- [10] M.Blum. *A machine-independent theory of the complexity of recursive functions*. Journal of the Association for Computing Machinery 14 (1967) 322-336.
- [11] G.Boolos. *The Unprovability of Consistency. An essay in modal logic*. Cambridge University Press, Cambridge 1979.
- [12] G.Boolos. *Extremely undecidable sentences*. The Journal of

- Symbolic Logic 47 (1982) 191-196.
- [13] A.Carbone. *Provable fixed points in $\text{I}\Delta_0 + \Omega_1$* . ITLI Prepublication Series ML-89-09. Institute for Language Logic and Information, University of Amsterdam 1989.
- [14] A.Carbone & F.Montagna. *Much shorter proofs: a bimodal investigation*. Zeitschrift für mathematische Logik und Grundlagen der Mathematik 36 (1990) 47-66.
- [15] А. В. Чагров. *Неразрешимые свойства расширений логики доказуемости*, Рукопись, Тверь 1989. Будет опубликовано в Алгебре и Логике.
- [16] S.Feferman. *Arithmetization of metamathematics in a general setting*. Fundamenta Mathematicae 49 (1960) 35-92.
- [17] K.Fine. *Logics containing K4. Part I*. The Journal of Symbolic Logic 34 (1974) 31-42.
- [18] K.Fine. *Logics containing K4. Part II*. The Journal of Symbolic Logic 50 (1985) 619-651.
- [19] H.Friedman. *The disjunction property implies the numerical existence property*. Proceedings of the National Academy of Sciences of the United States of America 72 (1975) 2877-2878.
- [20] Z.Gleit & W.Goldfarb. *Characters and fixed points in provability logic*, Manuscript, Cambridge 1989. To appear in Notre Dame Journal of Formal Logic.
- [21] Р. Ш. Григолия. *Свободные алгебры Магари с конечным числом образующих*. В кн.: *Логико-методологические исследования*. Мецниереба, Тбилиси 1983, 135-149.
- [22] Р. Ш. Григолия. *Свободные алгебры неклассических логик*. Мецниереба, Тбилиси 1987.
- [23] D.Guaspari. *Partially conservative extensions of arithmetic*. Transactions of the American Mathematical Society 254 (1979) 47-68.
- [24] P.Hájek & A.Kucera. *On recursion theory in $\text{I}\Sigma_1$* . The Journal of Symbolic Logic 54 (1989) 576-589.
- [25] D.de Jongh & F.Montagna. *Provable fixed points*. Zeitschrift für mathematische Logik und Grundlagen der Mathematik 34 (1988) 229-250.
- [26] D.Jensen & A.Ehrenfeucht. *Some problem in elementary arith-*

- metics. *Fundamenta Mathematicae* 92 (1976) 223-245.
- [27] M.Jumelet. *On Solovay's completeness theorem*. ILLI Prepublication Series X-88-01. Institute for Language Logic and Information, University of Amsterdam 1988.
- [28] A.Macintyre & H.Simmons. *Gödel's diagonalization technique and related properties of theories*. *Colloquium Mathematicum* 28 (1973) 165-180.
- [29] R.Magari. *The diagonalizable algebras (the algebraization of the theories which express Theor.: II)*. *Bollettino della Unione Matematica Italiana, Serie IV, 12* (1975) suppl. fasc. 3, 117-125.
- [30] Г.Е.Минц. *Бескванторные и однокванторные системы*. В кн.: Ю.В.Матиясевич и А.О.Слисенко, ред. *Исследования по конструктивной математике и математической логике. IV*. Записки научных семинаров, том 20. Наука, Ленинград 1971, 115-133 (English translation: *Journal of Soviet Mathematics* 1 (1973) 211-226).
- [31] F.Montagna. *On the diagonalizable algebra of Peano arithmetic*. *Bollettino della Unione Matematica Italiana, Serie V, 16-B* (1979) 795-812.
- [32] F.Montagna. *Iterated extensional Rosser's fixed points and hyperhyperdiagonalizable algebras*. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik* 33 (1987) 293-303.
- [33] F.Montagna. *Shortening proofs in arithmetic by modal rules: polynomial shortenings and exponential shortenings*. *Rapporto Matematico N.129*. Dipartimento di Matematica, Università di Siena 1990.
- [34] F.Montagna & G.Sommaruga. *Rosser and Mostowski sentences*. *Archive for Mathematical Logic* 27 (1988) 115-133.
- [35] R.Parikh. *Existence and feasibility in arithmetic*. *The Journal of Symbolic Logic* 36 (1971) 494-508.
- [36] J.B.Paris & L.A.S.Kirby. Σ_n -collection schemas in arithmetic. In: A.Macintyre, L.Pacholski & J.Paris, eds. *Logic Colloquium '77*. North-Holland, Amsterdam 1978, 199-209.
- [37] M.B.Pour-El & S.Kripke. *Deduction-preserving "recursive isomorphisms" between theories*. *Fundamenta Mathematicae* 61

(1967) 141-163.

- [38] H.Rogers jr. *Theory of Recursive Functions and Effective Computability*. McGraw-Hill, New York 1967.
- [39] В.В.Рыбаков. *О допустимости правил вывода в модальной системе G*. В кн.: Ю.Л.Ершов, ред. *Математическая логика и алгоритмические проблемы*. Труды Института Математики, том 12. Наука, Новосибирск 1989, 120-138.
- [40] K.Segerberg. *An Essay in Classical Modal Logic*. Filosofiska Studier nr 13. Filosofiska Föreningen och Filosofiska Institutionen vid Uppsala Universitet, Uppsala 1971.
- [41] H.Simmons. *Large discrete parts of the E-tree*. *The Journal of Symbolic Logic* 53 (1988) 980-984.
- [42] T.Smiley. *The logical basis of ethics*. *Acta Philosophica Fennica* 16 (1963) 237-246.
- [43] C.Smoryński. *Avoiding self-referential statements*. *Proceedings of the American Mathematical Society* 70 (1978) 181-184.
- [44] C.Smoryński. *Beth's theorem and self-referential sentences*. In: A.Macintyre, L.Pacholski & J.Paris, eds. *Logic Colloquium '77*. North-Holland, Amsterdam 1978, 253-261.
- [45] C.Smoryński. *Calculating self-referential statements*. *Fundamenta Mathematicae* 109 (1980) 189-210.
- [46] C.Smoryński. *Self-Reference and Modal Logic*. Springer-Verlag, New York 1985.
- [47] R.M.Solovay. *Provability interpretations of modal logic*. *Israel Journal of Mathematics* 25 (1976) 287-304.
- [48] A.Visser. *Aspects of Diagonalization and Provability*, Proefschrift, Utrecht 1981.
- [49] A.Visser. *The provability logics of recursively enumerable theories extending Peano arithmetic at arbitrary theories extending Peano arithmetic*. *Journal of Philosophical Logic* 13 (1984) 97-113.

The ITLI Prepublication Series

1990

Logic, Semantics and Philosophy of Language

LP-90-01 Jaap van der Does
LP-90-02 Jeroen Groenendijk, Martin Stokhof
LP-90-03 Renate Bartsch
LP-90-04 Aarne Ranta
LP-90-05 Patrick Blackburn
LP-90-06 Gennaro Chierchia
LP-90-07 Gennaro Chierchia
LP-90-08 Herman Hendriks
LP-90-09 Paul Dekker

LP-90-10 Theo M.V. Janssen
LP-90-11 Johan van Benthem
LP-90-12 Serge Lapierre
LP-90-13 Zhisheng Huang
LP-90-14 Jeroen Groenendijk, Martin Stokhof
LP-90-15 Maarten de Rijke
LP-90-16 Zhisheng Huang, Karen Kwast
LP-90-17 Paul Dekker

Mathematical Logic and Foundations

ML-90-01 Harold Schellinx
ML-90-02 Jaap van Oosten
ML-90-03 Yde Venema
ML-90-04 Maarten de Rijke
ML-90-05 Domenico Zambella
ML-90-06 Jaap van Oosten

ML-90-07 Maarten de Rijke
ML-90-08 Harold Schellinx
ML-90-09 Dick de Jongh, Duccio Pianigiani
ML-90-10 Michiel van Lambalgen
ML-90-11 Paul C. Gilmore

Computation and Complexity Theory

CT-90-01 John Tromp, Peter van Emde Boas
CT-90-02 Sieger van Denneheuvel
Gerard R. Renardel de Lavalette
CT-90-03 Ricard Gavaldà, Leen Torenvliet
Osamu Watanabe, José L. Balcázar
CT-90-04 Harry Buhrman, Edith Spaan
Leen Torenvliet
CT-90-05 Sieger van Denneheuvel, Karen Kwast
CT-90-06 Michiel Smid, Peter van Emde Boas
CT-90-07 Kees Doets
CT-90-08 Fred de Geus, Ernest Rotterdam,
Sieger van Denneheuvel, Peter van Emde Boas

Other Prepublications

X-90-01 A.S. Troelstra
X-90-02 Maarten de Rijke
X-90-03 L.D. Beklemishev
X-90-04
X-90-05 Valentin Shehtman
X-90-06 Valentin Goranko, Solomon Passy
X-90-07 V.Yu. Shavrukov
X-90-08 L.D. Beklemishev
X-90-09 V.Yu. Shavrukov
X-90-10 Sieger van Denneheuvel
Peter van Emde Boas
X-90-11 Alessandra Carbone
X-90-12 Maarten de Rijke
X-90-13 K.N. Ignatiev

X-90-14 L.A. Chagrova
X-90-15 A.S. Troelstra

1991

Mathematical Logic and Foundations

ML-91-01 Yde Venema
ML-91-02 Alessandro Berarducci
Rineke Verbrugge

Other Prepublications

X-91-01 Alexander Chagrov
Michael Zakharyashev
X-91-02 Alexander Chagrov
Michael Zakharyashev
X-91-03 V. Yu. Shavrukov
X-91-04 K.N. Ignatiev

A Generalized Quantifier Logic for Naked Infinitives
Dynamic Montague Grammar
Concept Formation and Concept Composition
Intuitionistic Categorical Grammar
Nominal Tense Logic
The Variability of Impersonal Subjects
Anaphora and Dynamic Logic
Flexible Montague Grammar
The Scope of Negation in Discourse,
towards a flexible dynamic Montague grammar
Models for Discourse Markers
General Dynamics
A Functional Partial Semantics for Intensional Logic
Logics for Belief Dependence
Two Theories of Dynamic Semantics
The Modal Logic of Inequality
Awareness, Negation and Logical Omniscience
Existential Disclosure, Implicit Arguments in Dynamic Semantics

Isomorphisms and Non-Isomorphisms of Graph Models
A Semantical Proof of De Jongh's Theorem
Relational Games
Unary Interpretability Logic
Sequences with Simple Initial Segments
Extension of Lifschitz' Realizability to Higher Order Arithmetic,
and a Solution to a Problem of F. Richman
A Note on the Interpretability Logic of Finitely Axiomatized Theories
Some Syntactical Observations on Linear Logic
Solution of a Problem of David Guaspari
Randomness in Set Theory
The Consistency of an Extended NaDSet

Associative Storage Modification Machines
A Normal Form for PCSJ Expressions

Generalized Kolmogorov Complexity
in Relativized Separations
Bounded Reductions

Efficient Normalization of Database and Constraint Expressions
Dynamic Data Structures on Multiple Storage Media, a Tutorial
Greatest Fixed Points of Logic Programs
Physiological Modelling using RL

Remarks on Intuitionism and the Philosophy of Mathematics,
Revised Version
Some Chapters on Interpretability Logic
On the Complexity of Arithmetical Interpretations of Modal Formulae
Annual Report 1989
Derived Sets in Euclidean Spaces and Modal Logic
Using the Universal Modality: Gains and Questions
The Lindenbaum Fixed Point Algebra is Undecidable
Provability Logics for Natural Turing Progressions of Arithmetical
Theories
On Rosser's Provability Predicate
An Overview of the Rule Language RL/1

Provable Fixed points in $\text{ID}_0 + \Omega_1$, revised version
Bi-Unary Interpretability Logic
Dzhaparidze's Polymodal Logic: Arithmetical Completeness,
Fixed Point Property, Craig's Property
Undecidable Problems in Correspondence Theory
Lectures on Linear Logic

Cylindric Modal Logic
On the Metamathematics of Weak Theories

The Disjunction Property of Intermediate Propositional Logics

On the Undecidability of the Disjunction Property of Intermediate
Propositional Logics
Subalgebras of Diagonizable Algebras of Theories containing Arithmetic
Partial Conservativity and Modal Logics