

ADMISSIBLE BASES VIA STABLE CANONICAL RULES

NICK BEZHANISHVILI, DAVID GABELAIA, SILVIO GHILARDI, MAMUKA JIBLADZE

ABSTRACT. We establish the dichotomy property of [17] for stable canonical multi-conclusion rules of [1] for **IPC**, **K4**, and **S4**. This yields an alternative proof of existence of explicit bases of admissible rules for these logics.

1. INTRODUCTION

An inference rule is *admissible* in a given logical system L if no new theorems are derived by adding this rule to the rules of inference of L . Friedman [9] raised the question whether admissibility of rules in the intuitionistic propositional calculus (**IPC**) is decidable. A solution to this problem for **IPC**, as well as for well-known systems of modal logic such as **K4** and **S4**, was first given by Rybakov (see the comprehensive book [23] and the references therein). An alternative solution via projectivity and unification was supplied in [10, 11]. Explicit bases for admissible rules were built in [22, 24, 20, 14, 16]. We refer to Goudsmit [13] for a modern historic account of the admissibility problem.

Recently Jeřábek [17] developed a new technique for building bases for admissible rules by generalizing Zakharyashev's canonical formulas [26] to multi-conclusion canonical rules, and by developing the *dichotomy property* for canonical rules. This property states that a canonical multi-conclusion rule is either admissible or equivalent to an assumption-free rule. Our goal is to establish the same property for *stable* multi-conclusion canonical rules for **IPC**, **K4**, and **S4**. These rules were recently introduced in [1], where it was shown that each normal modal multi-conclusion consequence relation is axiomatizable by stable multi-conclusion canonical rules. The same result for intuitionistic multi-conclusion consequence relations was established in [2].

The proof methodology we follow is similar to [17] and goes through a semantic characterization of non-admissible stable canonical rules in terms of the finite domains they are built from. In spite of the similarities, the semantic characterization we obtain is different than the one given in [17]. As a simple corollary of our main theorem, similarly to [17], we obtain decidability of the admissibility problem for **IPC**, **K4** and **S4**. Finally, we note that admissibility for the basic modal logic **K** is a long standing open problem. While the proofs of this paper do not directly apply to **K**, we observe that the method of stable canonical rules, unlike that of canonical rules of [17], is not limited to the transitive case. Therefore, our method is potentially applicable to non-transitive logics such as **K**.

The paper is organised as follows: In Section 2 we recall Esakia duality for Heyting algebras, multi-conclusion consequence relations and stable canonical rules for **IPC**. In Section 3 we obtain an explicit basis of admissible rules for **IPC** via stable canonical rules and prove that the latter have the dichotomy property. In Section 4 we recall duality for modal algebras, modal multi-conclusion consequence relations and stable canonical rules for modal logic. Finally, in Section 5 we obtain explicit bases of admissible rules for **K4** and **S4** via stable canonical rules and prove their dichotomy property.

2. PRELIMINARIES ON HEYTING ALGEBRAS AND IPC

2.1. Esakia duality for Heyting algebras. We recall that a *Heyting algebra* is a bounded distributive lattice with an additional binary operation \rightarrow that is residual of \wedge . For Heyting algebras

A and B , a *Heyting homomorphism* is a bounded lattice homomorphism $h : A \rightarrow B$ such that $h(a \rightarrow b) = h(a) \rightarrow h(b)$ for each $a, b \in A$. Let **Heyt** be the category of Heyting algebras and Heyting homomorphisms. It is well known (see, e.g., [19, Ch. IX] or [6, Ch. 7]) that Heyting algebras provide an adequate algebraic semantics for superintuitionistic logics. In fact, there is a dual isomorphism between the (complete) lattice of superintuitionistic logics and the (complete) lattice of varieties of Heyting algebras.

In order to introduce topological duality for **Heyt**, we need to fix some notation for posets. If X is a poset (partially ordered set), we denote the partial order on X by \leq . For $Y \subseteq X$, we recall that the *down-set* of Y is the set $\downarrow Y = \{x \in X : \exists y \in Y \text{ with } x \leq y\}$. The *up-set* of Y is defined dually and is denoted by $\uparrow Y$. If Y is a singleton set $\{y\}$, then we use $\downarrow y$ and $\uparrow y$ instead of $\downarrow\{y\}$ and $\uparrow\{y\}$, respectively. We call $U \subseteq X$ an *up-set* if $x \in U$ and $x \leq y$ imply $y \in U$. A *down-set* of X is defined dually. For $Y \subseteq X$ we denote by $\max Y$, resp $\min Y$ the set of its *maximal*, resp. *minimal* points. That is, $\max Y = \{y \in Y \mid Y \cap \uparrow y = \{y\}\}$ and $\min Y = \{y \in Y \mid Y \cap \downarrow y = \{y\}\}$.

An *Esakia space* is a Priestley space X such that $\downarrow U$ is clopen for each clopen U of X ; recall that a poset X is a *Priestley space* if X is a compact space and for each $x, y \in X$, from $x \not\leq y$ it follows that there is a clopen (closed and open) up-set U of X such that $x \in U$ and $y \notin U$. It follows easily from e. g. [7, 11.15(i)] that for any Priestley space (X, \leq) , any closed subset $Y \subseteq X$ and any $y \in Y$ there are $y_1 \in \min Y$, $y_2 \in \max Y$ with $y_1 \leq y \leq y_2$.

For posets X and Y , a map $f : X \rightarrow Y$ is *order-preserving* if $x \leq y$ implies $f(x) \leq f(y)$ for all $x, y \in X$; an order-preserving f is said to be a *bounded morphism* (or *p-morphism*) iff for each $x \in X$ and $y \in Y$, from $f(x) \leq y$ it follows that there exists $z \in X$ such that $x \leq z$ and $f(z) = y$.

For Esakia spaces X and Y , a map f is an *Esakia morphism* if it is a bounded morphism which is also continuous. Let **Esa** be the category of Esakia spaces and Esakia morphisms.

By Esakia duality [8], **Heyt** is dually equivalent to **Esa** (the dual of a Heyting algebra A is indicated with A_*). The functors $(-)_* : \mathbf{Heyt} \rightarrow \mathbf{Esa}$ and $(-)^* : \mathbf{Esa} \rightarrow \mathbf{Heyt}$ that establish this dual equivalence are constructed as follows. For a Heyting algebra A , let $A_* = (X, \leq)$, where X is the space of all prime filters of A (topologized by the subbasis $\{\alpha(a), X \setminus \alpha(a) : a \in A\}$, where $\alpha(a) = \{x \in X : a \in x\}$) and $x \leq y$ iff $x \subseteq y$. For a Heyting algebra homomorphism h , let $h_* = h^{-1}$. For an Esakia space (X, \leq) , let $(X, \leq)^* = A$, where A is the Heyting algebra of clopen up-sets of X , with meet and join given by intersection and union respectively and with implication given by $U \rightarrow V = X \setminus \downarrow(U \setminus V)$. For an Esakia morphism f , let $f^* = f^{-1}$.

It follows from Esakia duality that onto Heyting homomorphisms dually correspond to 1-1 Esakia morphisms, and 1-1 Heyting homomorphisms to onto Esakia morphisms. In particular, homomorphic images of $A \in \mathbf{Heyt}$ correspond to closed up-sets of the Esakia dual of A .

2.2. Intuitionistic multi-conclusion consequence relations. We use greek letters γ, δ, \dots , φ, ψ, \dots to denote formulas built up from propositional variables using the connectives $\neg, \wedge, \vee, \rightarrow, \perp, \top$. A valuation on a Heyting algebra A is a map associating an element of A with every propositional variable. It is then extended to all formulas in a standard way. An *intuitionistic Kripke model* is a triple (X, \leq, V) where (X, \leq) is a poset and V is a valuation on the Heyting algebra of its up-sets. We use letters $\mathfrak{M}, \mathfrak{N}, \dots$ for Kripke models and the notation $\mathfrak{M}, x \models \varphi$ to mean that x belongs to $V(\varphi)$, where V is the valuation on the Kripke model \mathfrak{M} . The notation $\mathfrak{M} \models \varphi$ means that $\mathfrak{M}, x \models \varphi$ holds for all x from the underlying poset of \mathfrak{M} .

A *multi-conclusion rule* is an expression Γ/Δ , where Γ, Δ are finite sets of formulas. If $\Delta = \{\varphi\}$, then Γ/Δ is called a *single-conclusion rule* and is written Γ/φ . If $\Gamma = \emptyset$, then Γ/Δ is called an *assumption-free rule* and is written Δ . Assumption-free single-conclusion rules Δ/φ can be identified with formulas φ .

Definition 2.1. *An intuitionistic multi-conclusion consequence relation is a set \mathcal{S} of multiple conclusion rules such that*

- (1) $\varphi/\varphi \in \mathcal{S}$.

- (2) $\varphi, \varphi \rightarrow \psi / \psi \in \mathcal{S}$.
- (3) $\varphi \in \mathcal{S}$ for each theorem φ of **IPC** (i.e. of intuitionistic propositional calculus).
- (4) If $\Gamma/\Delta \in \mathcal{S}$, then $\Gamma, \Gamma'/\Delta, \Delta' \in \mathcal{S}$.
- (5) If $\Gamma/\Delta, \varphi \in \mathcal{S}$ and $\Gamma, \varphi/\Delta \in \mathcal{S}$, then $\Gamma/\Delta \in \mathcal{S}$.
- (6) If $\Gamma/\Delta \in \mathcal{S}$ and σ is a substitution, then $\sigma(\Gamma)/\sigma(\Delta) \in \mathcal{S}$.

We denote the smallest intuitionistic multi-conclusion consequence relation by $\mathcal{S}_{\mathbf{IPC}}$. For a set \mathcal{R} of multi-conclusion rules, let $\mathcal{S}_{\mathbf{IPC}} + \mathcal{R}$ be the smallest intuitionistic multi-conclusion consequence relation containing \mathcal{R} . If $\mathcal{S} = \mathcal{S}_{\mathbf{IPC}} + \mathcal{R}$, then we say that \mathcal{S} is *axiomatized by \mathcal{R}* or that \mathcal{R} is a *basis for \mathcal{S}* . Whenever Γ/Δ belongs to $\mathcal{S}_{\mathbf{IPC}} + \mathcal{R}$ we say that Γ/Δ is *derivable from \mathcal{R}* .

A Heyting algebra A *validates* a multi-conclusion rule Γ/Δ provided for every valuation v on A , if $v(\gamma) = 1$ for all $\gamma \in \Gamma$, then $v(\delta) = 1$ for some $\delta \in \Delta$. If A validates Γ/Δ , we write $A \models \Gamma/\Delta$. The following result is proved in [17, 4]:

Theorem 2.2. Γ/Δ is derivable from \mathcal{R} iff every Heyting algebra validating all rules in \mathcal{R} validates also Γ/Δ .

We will say that rules ρ_1 and ρ_2 are *equivalent* if ρ_1 is derivable from $\{\rho_2\}$ and ρ_2 is derivable from $\{\rho_1\}$. By Theorem 2.2 this means that a Heyting algebra validates ρ_1 if and only if it validates ρ_2 .

Derivability should be contrasted with admissibility; we will call a rule Γ/Δ *admissible* in **IPC** (or *admissible tout court*) iff it is valid in the free Heyting algebra with countably many generators. It is known (see e. g. [21, 15]) that this is equivalent to either one of the following conditions: (1) every substitution making all members of Γ a theorem in **IPC** makes also some member of Δ a theorem of **IPC**, and (2) adding Γ/Δ to **IPC** does not lead to the derivability of new theorems.

A set of rules \mathcal{R} is said to form an *admissible basis* for a logic L if every rule admissible in L is derivable from \mathcal{R} .

2.3. Closed domain condition and stable canonical rules for Heyting algebras. We recall some definitions and results from [1].

Definition 2.3. Let $\mathfrak{X} = (X, \leq)$ and $\mathfrak{Y} = (Y, \leq)$ be Esakia spaces and let $f : X \rightarrow Y$ be a map. We call f *stable* if it is continuous and order-preserving.

It can be shown that Definition 2.3 can be dualized in the following way. Let A and B be Heyting algebras; then $h : A \rightarrow B$ is a bounded lattice morphism iff the dual Esakia morphism $h_* : B_* \rightarrow A_*$ is stable.

Definition 2.4. Let $\mathfrak{X} = (X, \leq)$ and $\mathfrak{Y} = (Y, \leq)$ be Esakia spaces, $f : X \rightarrow Y$ be a map, and U be a clopen subset of Y . We say that f satisfies the *closed domain condition (CDC)* for U if

$$U \cap \uparrow f(x) \neq \emptyset \Rightarrow U \cap f(\uparrow x) \neq \emptyset$$

holds for all $x \in X$. Let \mathfrak{D} be a collection of clopen subsets of Y . We say that $f : X \rightarrow Y$ satisfies the *closed domain condition (CDC)* for \mathfrak{D} if f satisfies CDC for each $U \in \mathfrak{D}$.

Stable canonical rules are introduced in the following definition:

Definition 2.5. Let A be a finite Heyting algebra and let $D \subseteq A^2$. For every $a \in A$ let p_a be a propositional letter, and define the *stable canonical rule* $\rho(A, D)$ associated with A and D as $\gamma(A, D) = \Gamma/\Delta$, where

$$\begin{aligned} \Gamma = & \{p_0 \leftrightarrow 0\} \cup \{p_1 \leftrightarrow 1\} \cup \\ & \{p_{a \vee b} \leftrightarrow (p_a \vee p_b) \mid a, b \in A\} \cup \\ & \{p_{a \wedge b} \leftrightarrow (p_a \wedge p_b) \mid a, b \in A\} \cup \\ & \{p_{a \rightarrow b} \leftrightarrow (p_a \rightarrow p_b) \mid (a, b) \in D\} \end{aligned}$$

and

$$\Delta = \{p_a \leftrightarrow p_b \mid a, b \in A \text{ with } a \neq b\}.$$

Sometimes, if F is the dual space of A , we might write $\gamma(F, D)$ instead of $\gamma(A, D)$.

Theorem 2.6. *Let A be a finite Heyting algebra, $D \subseteq A^2$, and B be an arbitrary Heyting algebra. Then the following are equivalent:*

- (i) $B \not\models \gamma(A, D)$;
- (ii) *there is a bounded lattice embedding $h : A \rightarrow B$ such that $h(a \rightarrow b) = h(a) \rightarrow h(b)$ for each $(a, b) \in D$;*
- (iii) *there is a stable onto map $f : B_* \rightarrow A_*$ satisfying CDC for $\mathfrak{D} := \{\alpha(a) \setminus \alpha(b) : (a, b) \in D\}$.*

The interesting point about stable rules is the following completeness theorem:

Theorem 2.7. [2, Prop. 3.4] *Any intuitionistic multi-conclusion consequence relation can be axiomatized by stable canonical rules.*

3. DICHOTOMY PROPERTY AND ADMISSIBLE BASIS FOR **IPC**

Let V_n be the rule:

$$(V_n) \quad \frac{((\bigvee_{i=1}^n p_i) \rightarrow q) \rightarrow \bigvee_{i=1}^n p_i}{q \rightarrow p_1 \mid \dots \mid q \rightarrow p_n}$$

(denoted by V'_n in [16]).

Theorem 3.1. *The rule V_n is admissible for each $n \in \omega$.*

Proof. We have to show that if σ is a substitution such that none of $\sigma q \rightarrow \sigma p_1, \dots, \sigma q \rightarrow \sigma p_n$ is a theorem of **IPC**, then $(\bigvee_{i=1}^n \sigma p_i \rightarrow \sigma q) \rightarrow \bigvee_{i=1}^n \sigma p_i$ is not a theorem either. By the finite model property of **IPC** there are finite rooted Kripke models $\mathfrak{M}_1, \dots, \mathfrak{M}_n$ such that $\mathfrak{M}_1 \models \sigma q, \dots, \mathfrak{M}_n \models \sigma q$ and $\mathfrak{M}_1 \not\models \sigma p_1, \dots, \mathfrak{M}_n \not\models \sigma p_n$ (a Kripke model is said to be rooted iff its underlying poset has a root, i. e. a smallest element). Consider the disjoint union of $\mathfrak{M}_1, \dots, \mathfrak{M}_n$ and add a new root r to it. Extend the valuation to the resulting frame by making each variable false at r . Denote the new model by \mathfrak{M} . Then $\mathfrak{M}, r \not\models \sigma p_1, \dots, \sigma p_n$. So, $\mathfrak{M}, r \not\models \bigvee_{i=1}^n \sigma p_i$ and $\mathfrak{M}, r \models \bigvee_{i=1}^n \sigma p_i \rightarrow \sigma q$. Thus $\mathfrak{M}, r \not\models (\bigvee_{i=1}^n \sigma p_i \rightarrow \sigma q) \rightarrow \bigvee_{i=1}^n \sigma p_i$. Hence $(\bigvee_{i=1}^n \sigma p_i \rightarrow \sigma q) \rightarrow \bigvee_{i=1}^n \sigma p_i$ is not a theorem of **IPC**. \square

Lemma 3.2. *Suppose that a stable canonical rule $\gamma(F, D)$ has the following property. Given an Esakia space W and a clopen up-set $Y \subseteq W$ and a stable surjective map $f : Y \rightarrow F$ satisfying CDC for D , there is stable surjective $\bar{f} : W \rightarrow F$ with $f \subseteq \bar{f}$ satisfying CDC for D . Then $\gamma(F, D)$ is equivalent to an assumption-free rule.*

Proof. Let $\gamma(A, D)$ be the rule

$$(1) \quad \frac{\varphi}{\varphi_1 \mid \dots \mid \varphi_n}$$

We will show that under the assumption of the lemma this rule is equivalent to

$$(2) \quad \frac{}{\varphi \rightarrow \varphi_1 \mid \dots \mid \varphi \rightarrow \varphi_n}.$$

(2) \Rightarrow (1) is clear. Now assume that (the Heyting algebra dual to the Esakia space) W does not validate (2). We show that then it does not validate (1). Let V be a valuation on W such that $V(\varphi) \not\subseteq V(\varphi_1), \dots, V(\varphi) \not\subseteq V(\varphi_n)$. We set $Y = V(\varphi) \subseteq W$. Then $Y \not\models (1)$. This means that there is a stable surjective $f : Y \rightarrow F$. By the condition of the lemma f is extended to stable surjective $\bar{f} : W \rightarrow F$, implying $W \not\models (1)$. \square

The following definition will be our main ingredient for a semantic characterization of admissibility of a stable canonical rule:¹

Definition 3.3. *A stable canonical rule $\gamma(A, D)$ is called trivial if for all $S \subseteq A_*$ there is $x \in A_*$ such that*

- (1) $S \subseteq \uparrow x$
- (2) For all $d \in D$ if $d \cap \uparrow x \neq \emptyset$, then $d \cap (\{x\} \cup \uparrow S) \neq \emptyset$.

We will see below that the triviality condition plays the same role for stable canonical rules as the existence of tight predecessors in the context of [17, Theorem 4.9 (iv)].

Theorem 3.4. *The following are equivalent:*

- (1) $\gamma(A, D)$ is admissible.
- (2) $\gamma(A, D)$ is derivable from $\{V_n : n \in \omega\}$.
- (3) $\gamma(A, D)$ is not trivial.
- (4) $\gamma(A, D)$ is not equivalent to an assumption-free rule.

Proof. (2) \Rightarrow (1). We know that all V_n are admissible, i. e. valid in the free Heyting algebra on infinitely many generators. Since moreover $\gamma(A, D)$ is derivable from $\{V_n : n \in \omega\}$, we conclude that $\gamma(A, D)$ is also valid on this algebra, i. e. is admissible.

(3) \Rightarrow (2). Let $F = A_*$ and suppose $\gamma(F, D)$ is not derivable from $\{V_n : n \in \omega\}$. Then, by Theorem 2.2, there is an Esakia space W validating all V_n 's and refuting $\gamma(F, D)$. The latter means that there is a stable surjective $f : W \rightarrow F$ satisfying CDC for D . We will now show that $\gamma(F, D)$ is trivial. In what follows, we will employ the Heyting algebra W^* ; in particular, implication will be understood in the sense of this algebra. Fix $S \subseteq F$. For $s \in S$ let $p_s = f^{-1}(F \searrow \downarrow s) \subseteq W$, $q = f^{-1}(\uparrow S)$. Since f is stable, p_s and q are up-sets. For all $s \in S$ we have $q \not\subseteq p_s$. Indeed, if $x_s \in f^{-1}(s)$, then we have that $x_s \in q$ but $x_s \notin p_s$.

Since W validates the rules V_n for each $n \in \omega$, and none of $q \rightarrow p_s$ are the whole of W , it follows that neither $((\bigcup_{s \in S} p_s) \rightarrow q) \rightarrow \bigcup_{s \in S} p_s$ is the whole of W ; in particular, $((\bigcup_{s \in S} p_s) \rightarrow q) \setminus (\bigcup_{s \in S} p_s)$ is not empty. As the topology on F is discrete, p_s and q are clopen sets. Thus both $(\bigcup_{s \in S} p_s) \rightarrow q$ and $\bigcup_{s \in S} p_s$ are clopen too, and we may actually pick a maximal element y of $((\bigcup_{s \in S} p_s) \rightarrow q) \setminus \bigcup_{s \in S} p_s$.

We claim that then for each $y' > y$ we have $y' \in q$. Indeed since $(\bigcup_{s \in S} p_s) \rightarrow q$ is an upset and y belongs to it, also y' will belong to it. But then $y' \notin \bigcup_{s \in S} p_s$ is impossible by maximality of y , so $y' \in \bigcup_{s \in S} p_s$, hence $y' \in q$.

Let us now check that $f(y)$ fulfils the triviality conditions for S . For the first condition just note that $y \notin \bigcup_{s \in S} p_s$ iff for all $s \in S$ we have $y \notin f^{-1}(F \searrow \downarrow s)$, i. e. $y \in \bigcap_{s \in S} f^{-1}(\downarrow s)$, which is equivalent to $\uparrow f(y) \supseteq S$. For the second condition, suppose $d \cap \uparrow f(y) \neq \emptyset$ for $d \in D$, then by the CDC of f we have that there is $y' \geq y$ such that $f(y') \in d$. Thus, either $y' = y$ and then $f(y') = f(y) \in d \cap \{f(y)\}$ or $y' > y$ and then, as we have seen, $y' \in q = f^{-1}(\uparrow S)$, so $f(y') \in d \cap \uparrow S$. Thus $\gamma(A, D)$ is trivial.

(4) \Rightarrow (3) Suppose $\gamma(A, D)$ is trivial. We show that then it is equivalent to an assumption-free rule. We use Lemma 3.2. Let W be an Esakia space, $Y \subseteq W$ a clopen up-set and $f : Y \rightarrow F$ a stable surjective map satisfying CDC for D . We extend f to some $f^l : W \rightarrow F$ with the same properties. For $w \in W$ let $f_w = f(Y \cap \uparrow w)$. If $S \subseteq F$ is of the kind f_w , let $Y_S \subseteq W$ be $Y_S = \{w \in W \setminus Y : f_w = S\}$. We take a minimal $S \subseteq F$ such that $Y_S \neq \emptyset$ and extend f to $Y \cup Y_S$.

Claim 3.5. *$Y \cup Y_S$ is a clopen up-set.*

Proof. It follows from the minimality of S that $Y \cup Y_S$ is an up-set. Indeed, if $x \in Y \cup Y_S$ and $x \leq y$, then either $y \in Y$ and then we are done, or, provided $y \notin Y$, in view of minimality of S , $f_y = S$.

¹The triviality notion below was independently introduced by J. Goudsmit in his thesis [13, Def. 4.76] under the name of 'adequate extendibility'. The author uses this notion when revisiting Rybakov results on admissible rules via universal (finite variable) models.

Indeed since Y is an up-set, $y \notin Y$ implies $x \notin Y$, so $x \in Y_S$, i. e. $f_x = S$. Moreover $\uparrow y \subseteq \uparrow x$, hence

$$f_y = f(Y \cap \uparrow y) \subseteq f(Y \cap \uparrow x) = f_x = S,$$

thus, as S is minimal, $f_y = S$, i. e. $y \in Y_S$.

To show that $Y \cup Y_S$ is clopen it suffices to show that Y_S is clopen. Now for any $w \in W$ we have that $w \in (W \setminus \downarrow f^{-1}(F \setminus S)) \setminus Y$ if and only if $f(Y \cap \uparrow w) \subseteq S$ and $w \notin Y$, which by minimality of S is equivalent to $f(Y \cap \uparrow w) = S$ and $w \notin Y$. Thus $Y_S = (W \setminus \downarrow f^{-1}(F \setminus S)) \setminus Y$ is clopen. This finishes the proof of the claim. \square

We now extend f to \bar{f} with $\text{dom}(\bar{f}) = Y \cup Y_S$. We put

$$\bar{f}(w) = \begin{cases} f(w), & \text{if } w \in Y, \\ s, & \text{if } w \notin Y, \end{cases}$$

where s is such that $S \subseteq \uparrow s$ and for all $d \in D$, $d \cap \uparrow s \neq \emptyset \Rightarrow d \cap (\{s\} \cup \uparrow S) \neq \emptyset$.

It is easy to see that \bar{f} is order-preserving. Now we also show that \bar{f} is continuous. Indeed, for every $x \in F$ we have $\bar{f}^{-1}(x) = f^{-1}(x)$ or $\bar{f}^{-1}(x) = f^{-1}(x) \cup Y_S$. Since Y_S is a clopen set the continuity follows.

Finally, we show that \bar{f} satisfies CDC. The relevant case is when $d \cap \uparrow \bar{f}(w) \neq \emptyset$ for $d \in D$, $w \in Y_S$. Now $\uparrow \bar{f}(w) = \uparrow s$. Thus, we have $d \cap (\{s\} \cup \uparrow S) \neq \emptyset$ by the choice of s . Hence either $\bar{f}(w) \in d$ or there is $s' \in S$ with $d \cap \uparrow s' \neq \emptyset$. But $f_w = S$ (because $w \in Y_S$), hence $f(Y \cap \uparrow w) = S$. So there is $w' \geq w$ such that $w' \in Y$ and $f(w') = s'$. We can then apply CDC for f to get $w'' \geq w'$ with $f(w'') \in d$. Thus, $w'' \geq w$ and $\bar{f}(w'') = f(w'') \in d$.

So we extended f to \bar{f} on $Y \cup Y_S$. We need to show that by repeating this procedure we cover the whole of W . This holds since the following is true: if some $S \subseteq F$ has been used for further extension of the map according to the above procedure, then this same S can never occur again during any subsequent extensions.

Indeed let f^k , resp. f^n be any further extensions of f to Y^k , resp. Y^n , $k < n < \omega$. Suppose we have used some S for f^k ; then it cannot happen that S can be also used for f^n .

Suppose, to the contrary, that S occurs as one of the candidates to build f^n . Then in particular $S = f_w^{n-1}$ for some $w \in Y^n \setminus Y^{n-1}$. Then also $w \notin Y^k$, so $S \neq f_w^{k-1}$ (since $Y^k = Y^{k-1} \cup Y_S^{k-1}$ and Y_S^{k-1} consists precisely of those v for which $f_v^{k-1} = S$). In fact by the minimality of S , f_w^{k-1} cannot be included in S , so $f_w^{k-1} \setminus S$ is nonempty. Now note that since f^{n-1} is an extension of f^{k-1} , one has $f_w^{n-1} \supseteq f_w^{k-1}$, hence also $f_w^{n-1} \setminus S$ is nonempty, which contradicts the equality $S = f_w^{n-1}$ above.

It thus follows that after each next extension at least one subset of F is excluded from all subsequent extension steps. Thus after some step n there will be no $w \notin Y^n$ and no S left with the property $f_w^n = S$. Which just means that there is no w outside Y^n , i. e. $Y^n = W$.

(1) \Rightarrow (4) Suppose $\gamma(A, D)$ is admissible and equivalent to an assumption-free rule $\diagup \Delta$. Then by the definition of admissibility any substitution makes one of the formulas in Δ a theorem of **IPC**. Hence $\diagup \Delta$ is valid on any Heyting algebra. However, $A \not\# \gamma(A, D)$, which is a contradiction. \square

Corollary 3.6. *A stable canonical rule $\gamma(A, D)$ has the following dichotomy property: it is either admissible or equivalent to an assumption-free rule.*

Corollary 3.7. *Admissibility in **IPC** is decidable.*

Proof. Given a rule ρ , we effectively compute the stable canonical rules $\gamma(A_1, D_1), \dots, \gamma(A_n, D_n)$ which are equivalent to ρ over **IPC** [1, 2]. Hence ρ is admissible if and only if each of the $\gamma(A_1, D_1), \dots, \gamma(A_n, D_n)$ is admissible. By Theorem 3.4, each $\gamma(A_i, D_i)$ is admissible iff it is

not trivial. Obviously, triviality of a rule $\gamma(A_i, D_i)$ can be checked in finite time. The result follows.² \square

Corollary 3.8. *The rules $\{V_n : n \in \omega\}$ form an admissible basis for IPC.*

Proof. By Theorem 2.7 and the above. \square

4. PRELIMINARIES ON MODAL ALGEBRAS AND MODAL LOGICS

4.1. Duality for modal algebras. We use [6, 18, 5, 25] as our main references for the basic theory of normal modal logics, including their algebraic and relational semantics, and the dual equivalence between modal algebras and modal spaces (descriptive Kripke frames).

A *modal algebra* is a pair $\mathfrak{A} = (A, \diamond)$, where A is a Boolean algebra and \diamond is a unary operator on A that commutes with finite joins. As usual, the dual operator \square is defined as $\neg \diamond \neg$. A *modal homomorphism* between two modal algebras is a Boolean homomorphism h satisfying $h(\diamond a) = \diamond h(a)$. Let \mathbf{MA} be the category of modal algebras and modal homomorphisms.

A *modal space* (or *descriptive Kripke frame*) is a pair $\mathfrak{X} = (X, R)$, where X is a Stone space (zero-dimensional compact Hausdorff space) and R is a binary relation on X satisfying the conditions:

$$R[x] := \{y \in X : xRy\}$$

is closed for each $x \in X$ and

$$R^{-1}[U] := \{x \in X : \exists y \in U \text{ with } xRy\}$$

is clopen (closed and open) for each clopen U of X . A *bounded morphism* (or *p-morphism*) $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ between two modal spaces is a continuous map $f : X \rightarrow Y$ such that $f(R[x]) = R[f(x)]$ for all $x \in X$. Let \mathbf{MS} be the category of modal spaces and bounded morphisms.

It is a well-known theorem in modal logic that \mathbf{MA} is dually equivalent to \mathbf{MS} . The functors $(-)_* : \mathbf{MA} \rightarrow \mathbf{MS}$ and $(-)^* : \mathbf{MS} \rightarrow \mathbf{MA}$ that establish this dual equivalence are constructed as follows. For a modal algebra $\mathfrak{A} = (A, \diamond)$, let $\mathfrak{A}_* = (A_*, R)$, where A_* is the Stone space of A (that is, the set of ultrafilters of A topologized by the basis $\{\beta(a) : a \in A\}$, where $\beta(a) = \{x \in A_* : a \in x\}$) and xRy iff $(\forall a \in A)(a \in y \Rightarrow \diamond a \in x)$. We call R the *dual* of \diamond . For a modal homomorphism h , let $h_* = h^{-1}$. For a modal space $\mathfrak{X} = (X, R)$, let $\mathfrak{X}^* = (A, \diamond)$, where A is the Boolean algebra of clopens of X and $\diamond(U) = R^{-1}[U]$. For a bounded morphism f , let $f^* = f^{-1}$.

Let $\mathfrak{A} = (A, \diamond)$ be a modal algebra and let $\mathfrak{X} = (X, R)$ be its dual space. Then it is well known that R is reflexive iff $a \leq \diamond a$ for all $a \in A$, and R is transitive iff $\diamond \diamond a \leq \diamond a$ for all $a \in A$. A modal algebra \mathfrak{A} is a **K4**-algebra if $\diamond \diamond a \leq \diamond a$ holds in \mathfrak{A} , and it is an **S4**-algebra if in addition $a \leq \diamond a$ holds in \mathfrak{A} . **S4**-algebras are also known as closure algebras, interior algebras, or topological Boolean algebras. Let $\mathbf{K4}$ be the full subcategory of \mathbf{MA} consisting of **K4**-algebras, and let $\mathbf{S4}$ be the full subcategory of $\mathbf{K4}$ consisting of **S4**-algebras. A modal space $\mathfrak{X} = (X, R)$ is a *transitive space* if R is transitive, and it is a *quasi-ordered space* if R is reflexive and transitive.

For a clopen subset $Y \subseteq X$ of a transitive space (X, R) , a point $y \in Y$ is called *quasi-maximal* if for any $x \in Y$ with yRx we have xRy . It is known that any point of any clopen subset sees a quasi-maximal point of this subset (see e. g. [6, Theorem 10.36]).

Let \mathbf{TS} be the full subcategory of \mathbf{MS} consisting of transitive spaces, and let \mathbf{QS} be the full subcategory of \mathbf{TS} consisting of quasi-ordered spaces. Then the dual equivalence of \mathbf{MA} and \mathbf{MS} restricts to the dual equivalence of $\mathbf{K4}$ and \mathbf{TS} , which restricts further to the dual equivalence of $\mathbf{S4}$ and \mathbf{QS} .

²An alternative proof can be given as follows: rule admissibility is Π_1^0 and derivability from a recursive set of rules $\{V_n : n \in \omega\}$ is Σ_1^0 . Thus, admissibility is decidable.

4.2. Multi-conclusion modal rules. We use greek letters $\gamma, \delta, \dots, \varphi, \psi, \dots$ to denote formulas built up from propositional variables using the connectives $\neg, \wedge, \vee, \rightarrow, \perp, \top, \diamond$. A valuation on a modal algebra $\mathfrak{A} = (A, \diamond)$ is a map associating an element of A with every propositional variable. It is then extended to all modal formulas in a standard way. A *Kripke frame* is a pair (X, R) where X is a set and R is a binary relation on X . A *Kripke model* is a triple (X, R, V) , where (X, R) is a Kripke frame and V is a valuation on the powerset Boolean algebra of X with $\diamond := R^{-1}$. We use letters $\mathfrak{M}, \mathfrak{N}, \dots$ for Kripke models and the notation $\mathfrak{M}, x \models \varphi$ to mean that x belongs to $V(\varphi)$, where V is the valuation of the Kripke model \mathfrak{M} . The notation $\mathfrak{M} \models \varphi$ (' φ is valid in \mathfrak{M} ') means that $\mathfrak{M}, x \models \varphi$ holds for all x from the underlying frame of \mathfrak{M} . We let $\mathbf{K}, \mathbf{K4}, \mathbf{S4}$ stand for the set of formulas which are valid in all modal algebras, $\mathbf{K4}$ -modal algebras, $\mathbf{S4}$ -modal algebras, respectively (as it is well-known, we can equivalently use validity in the corresponding classes of Kripke models).

A *transitive normal modal multi-conclusion consequence relation* is a set \mathcal{S} of modal rules such that

- (1) $\varphi/\varphi \in \mathcal{S}$.
- (2) $\varphi, \varphi \rightarrow \psi/\psi \in \mathcal{S}$.
- (3) $\varphi/\Box\varphi \in \mathcal{S}$.
- (4) $\varphi \in \mathcal{S}$ for each φ in $\mathbf{K4}$.
- (5) If $\Gamma/\Delta \in \mathcal{S}$, then $\Gamma, \Gamma'/\Delta, \Delta' \in \mathcal{S}$.
- (6) If $\Gamma/\Delta, \varphi \in \mathcal{S}$ and $\Gamma, \varphi/\Delta \in \mathcal{S}$, then $\Gamma/\Delta \in \mathcal{S}$.
- (7) If $\Gamma/\Delta \in \mathcal{S}$ and σ is a substitution, then $\sigma(\Gamma)/\sigma(\Delta) \in \mathcal{S}$.

We denote the least transitive normal modal multi-conclusion consequence relation by $\mathbf{S}_{\mathbf{K4}}$. For a set \mathcal{R} of multi-conclusion modal rules, let $\mathbf{S}_{\mathbf{K4}} + \mathcal{R}$ be the least transitive normal modal multi-conclusion consequence relation containing \mathcal{R} . If $\mathcal{S} = \mathbf{S}_{\mathbf{K4}} + \mathcal{R}$, then we say that \mathcal{S} is *axiomatized* by \mathcal{R} or that \mathcal{R} is a *basis* for \mathcal{S} . Whenever Γ/Δ belongs to $\mathbf{S}_{\mathbf{K4}} + \mathcal{R}$ we say that Γ/Δ is *derivable* from \mathcal{R} .

A $\mathbf{K4}$ algebra A *validates* a multi-conclusion rule Γ/Δ provided for every valuation v on A , if $v(\gamma) = 1$ for all $\gamma \in \Gamma$, then $v(\delta) = 1$ for some $\delta \in \Delta$. If A validates Γ/Δ , we write $A \models \Gamma/\Delta$. The following result is proved in [17, 4]:

Theorem 4.1. Γ/Δ is derivable from \mathcal{R} iff every $\mathbf{K4}$ -algebra validating all rules in \mathcal{R} also validates Γ/Δ .

Admissibility of rules in modal calculi is defined similarly to the intuitionistic case (described in 2.2) and has similar properties.

4.3. Closed domain conditions and stable canonical rules for modal algebras. We now introduce the key concepts of stable homomorphisms and the closed domain condition, and show how the two relate to each other. For the proofs of the results stated in this subsection, the reader is referred to [1].

Definition 4.2. Let $\mathfrak{A} = (A, \diamond)$ and $\mathfrak{B} = (B, \diamond)$ be $\mathbf{K4}$ -algebras and let $h : A \rightarrow B$ be a Boolean homomorphism. We call h a *stable homomorphism* provided $\diamond h(a) \leq h(\diamond a)$ for each $a \in A$.

It is easy to see that $h : A \rightarrow B$ is stable iff $h(\Box a) \leq \Box h(a)$ for each $a \in A$. Stable homomorphisms were considered in [3] under the name of semi-homomorphisms and in [12] under the name of continuous morphisms.

Definition 4.3. Let $\mathfrak{X} = (X, R)$ and $\mathfrak{Y} = (Y, R)$ be transitive modal spaces and let $f : X \rightarrow Y$ be a map. We call f *stable* if it is continuous and xRy implies $f(x)Rf(y)$.

Lemma 4.4. Let $\mathfrak{A} = (A, \diamond)$ and $\mathfrak{B} = (B, \diamond)$ be $\mathbf{K4}$ -algebras, $\mathfrak{X} = (X, R)$ be the dual of \mathfrak{A} , $\mathfrak{Y} = (Y, R)$ be the dual of \mathfrak{B} , and $h : A \rightarrow B$ be a Boolean homomorphism. Then $h : A \rightarrow B$ is stable iff $h_* : Y \rightarrow X$ is stable.

Definition 4.5. Let $\mathfrak{X} = (X, R)$ and $\mathfrak{Y} = (Y, R)$ be transitive modal spaces, $f : X \rightarrow Y$ be a map, and U be a clopen subset of Y . We say that f satisfies the closed domain condition (CDC) for U if

$$R[f(x)] \cap U \neq \emptyset \Rightarrow f(R[x]) \cap U \neq \emptyset.$$

Let \mathfrak{D} be a collection of clopen subsets of Y . We say that $f : X \rightarrow Y$ satisfies the closed domain condition (CDC) for \mathfrak{D} if f satisfies CDC for each $U \in \mathfrak{D}$.

Theorem 4.6. Let $\mathfrak{A} = (A, \diamond)$ and $\mathfrak{B} = (B, \diamond)$ be **K4**-algebras, $h : A \rightarrow B$ be a stable homomorphism, and $a \in A$. The following two conditions are equivalent:

- (1) $h(\diamond a) = \diamond h(a)$.
- (2) $h_* : B_* \rightarrow A_*$ satisfies CDC for $\beta(a)$.

Theorem 4.6 motivates the following definition.

Definition 4.7. Let $\mathfrak{A} = (A, \diamond)$ and $\mathfrak{B} = (B, \diamond)$ be **K4**-algebras and let $h : A \rightarrow B$ be a stable homomorphism.

- (1) We say that h satisfies the closed domain condition (CDC) for $a \in A$ if $h(\diamond a) = \diamond h(a)$.
- (2) We say that h satisfies the closed domain condition (CDC) for $D \subseteq A$ if h satisfies CDC for each $a \in D$.

We now come to stable canonical rules:

Definition 4.8. Let $\mathfrak{A} = (A, \diamond)$ be a finite **K4**-algebra and let D be a subset of A . For each $a \in A$ we introduce a new propositional letter p_a and define the stable canonical rule $\rho(\mathfrak{A}, D)$ associated with \mathfrak{A} and D as Γ/Δ , where:

$$\begin{aligned} \Gamma &= \{p_{a \vee b} \leftrightarrow p_a \vee p_b : a, b \in A\} \cup \\ &\quad \{p_{\neg a} \leftrightarrow \neg p_a : a \in A\} \cup \\ &\quad \{\diamond p_a \rightarrow p_{\diamond a} : a \in A\} \cup \\ &\quad \{p_{\diamond a} \rightarrow \diamond p_a : a \in D\}, \end{aligned}$$

and

$$\Delta = \{p_a : a \in A, a \neq 1\}.$$

Stable canonical rules are characterized in terms of refutations as follows:

Theorem 4.9. Let $\mathfrak{A} = (A, \diamond)$ be a finite **K4**-algebra, $D \subseteq A$, and $\mathfrak{B} = (B, \diamond)$ be a **K4**-algebra. Then $\mathfrak{B} \neq \rho(\mathfrak{A}, D)$ iff there is a stable embedding $h : A \rightarrow B$ satisfying CDC for D .

It was proved in [1] that every multi-conclusion consequence relation above **K** is axiomatizable by stable canonical rules (relative to arbitrary finite modal algebras - not only to those validating **K4**-axiom). The same proof can easily be extended to our multi-conclusion consequence relations above **K4**. Thus, we have the following theorem.

Theorem 4.10. Any transitive normal modal multi-conclusion consequence relation can be axiomatized by canonical rules $\rho(\mathfrak{A}, D)$ (where $\mathfrak{A} = (A, \diamond)$ is a finite **K4**-algebra and $D \subseteq A$).

5. DICHOTOMY PROPERTY AND ADMISSIBLE BASIS FOR **K4**

From now on, all Kripke frames and modal spaces are assumed to be transitive. Below $\Box^+\varphi$ abbreviates $\varphi \wedge \Box\varphi$; in a transitive Kripke frame/modal space (X, R) , R^+ abbreviates $R \cup id$ and $\uparrow S$ stands for $\{w \in X \mid \exists s \in S \ sR^+w\}$. We may also use the notation $\uparrow S$ for $\{w \in X \mid \exists s \in S \ sRw\}$. When we say that S is an up-set we mean $S = \uparrow S$. If S is a singleton set $\{y\}$, then we use $\uparrow y$ and $\uparrow y$ instead of $\uparrow\{y\}$ and $\uparrow\{y\}$, respectively. Notations $\downarrow S, \downarrow S, \downarrow\{y\}$ and $\downarrow\{y\}$ are defined dually (notice that $R^{-1}(S)$ is the same as $\downarrow S$).

Let $F = (W, R)$ be a frame dual to a finite **K4**-algebra $\mathfrak{A} = (A, \diamond)$. We denote the set $\{\beta(a) : a \in D\}$ by D . We will also denote (abusing notation) the stable canonical rule $\rho(\mathfrak{A}, D)$ by $\rho(F, D)$.

Let $S_n^{\ell, m}$ be the rule:

$$(S_n^{\ell, m}) \quad \frac{(\bigwedge_{l=1}^{\ell} (\Box v_l \rightarrow v_l) \wedge \bigwedge_{k=1}^m \Box(r_k \rightarrow \Box(r_k \vee \Box^+ q))) \rightarrow \bigvee_{i=1}^n \Box p_i}{\Box^+ q \rightarrow p_1 \mid \cdots \mid \Box^+ q \rightarrow p_n}$$

and T_n^m be the rule:

$$(T_n^m) \quad \frac{\bigwedge_{k=1}^m (\diamond r_k \rightarrow \diamond(r_k \wedge \Box^+ q)) \rightarrow \bigvee_{i=1}^n \Box p_i}{\Box^+ q \rightarrow p_1 \mid \cdots \mid \Box^+ q \rightarrow p_n}$$

Theorem 5.1.

- (1) The rule $(S_n^{\ell, m})$ is admissible in **K4** for each $n, m, \ell \in \omega$.
- (2) The rule (T_n^m) is admissible in **K4** for all $m \in \omega$.

Proof. The foregoing proof is essentially an adjustment of the proof of Theorem 3.1.

(1) We have to show that if σ is a substitution such that none of $\Box^+ \sigma q \rightarrow \sigma p_1, \dots, \Box^+ \sigma q \rightarrow \sigma p_n$ is a theorem of **K4**, then $(\bigwedge_{l=1}^{\ell} (\Box \sigma v_l \rightarrow \sigma v_l) \wedge \bigwedge_{k=1}^m \Box(\sigma r_k \rightarrow \Box(\sigma r_k \vee \Box^+ \sigma q))) \rightarrow \bigvee_{i=1}^n \Box \sigma p_i$ is not a theorem either.

By the finite model property of **K4** there are finite rooted Kripke models $\mathfrak{M}_1, \dots, \mathfrak{M}_n$ such that $\mathfrak{M}_i \models \Box^+ \sigma q$ and $\mathfrak{M}_i \not\models \sigma p_i$ for all $i = 1, \dots, n$. Consider the disjoint union of $\mathfrak{M}_1, \dots, \mathfrak{M}_n$ and add a new reflexive root ϱ to it. Extend the valuation to the resulting frame by making each variable false at ϱ . Denote the new model by \mathfrak{M} . Then by reflexivity of ϱ we will have $\mathfrak{M}, \varrho \models \bigwedge_{l=1}^{\ell} (\Box \sigma v_l \rightarrow \sigma v_l)$. Moreover $\mathfrak{M}, \varrho \not\models \bigvee_{i=1}^n \Box \sigma p_i$, because for every i , ϱ sees a point in \mathfrak{M}_i where σp_i is not true. Thus we will be done if $\mathfrak{M}, \varrho \models \bigwedge_{k=1}^m \Box(\sigma r_k \rightarrow \Box(\sigma r_k \vee \Box^+ \sigma q))$. This means that for any k and any w in \mathfrak{M} with $\mathfrak{M}, w \models \sigma r_k$ one has $\mathfrak{M}, w' \models \sigma r_k \vee \Box^+ \sigma q$ for all w' with wRw' . But any such w' is either in some \mathfrak{M}_i and then $\mathfrak{M}, w' \models \Box^+ \sigma q$, or $w' = \varrho$, and then because of wRw' also $w = \varrho$, so $\mathfrak{M}, w' \models \sigma r_k$. In both cases $\mathfrak{M}, w' \models \sigma r_k \vee \Box^+ \sigma q$.

(2) The rule (T_n^m) is proved to be admissible in a similar way (this time, an irreflexive extra root is needed). \square

Lemma 5.2. *Suppose that a stable canonical rule $\rho(F, D)$ has the following property. Given a transitive modal space (W, R) and a clopen up-set $Y \subseteq W$ and a stable surjective map $f : Y \rightarrow F$ satisfying CDC for D , there is stable surjective $\bar{f} : W \rightarrow F$ with $f \subseteq \bar{f}$ satisfying CDC for D . Then $\rho(F, D)$ is equivalent to an assumption-free rule.*

Proof. Let $\rho(A, D)$ be the rule

$$(1) \quad \frac{\varphi}{\varphi_1 \mid \cdots \mid \varphi_n}$$

We will show that under the assumption of the lemma this rule is equivalent to

$$(2) \quad \frac{}{\Box^+ \varphi \rightarrow \varphi_1 \mid \cdots \mid \Box^+ \varphi \rightarrow \varphi_n}$$

(2) \Rightarrow (1) is clear. Now assume that a transitive modal space (W, R) does not validate (2). We show that then it does not validate (1). Let V be a valuation on W such that $V(\Box^+ \varphi) \not\subseteq V(\varphi_1), \dots, V(\Box^+ \varphi) \not\subseteq V(\varphi_n)$. We set $Y = V(\Box^+ \varphi) \subseteq W$. Then $Y \not\models (1)$. This means that there is a stable surjective $f : Y \rightarrow F$ satisfying CDC for D . By the condition of the lemma f can be extended to a stable surjective map $\bar{f} : W \rightarrow F$ satisfying CDC for D , implying $W \not\models (1)$. \square

The following is a modal analogue of Definition 3.3.

Definition 5.3. *A stable canonical rule $\rho(A, D)$ is called trivial^o if for all $S \subseteq A_*$ there is a reflexive $x^\circ \in A_*$ such that*

$$(1): S \subseteq \uparrow x^\circ$$

(2): For all $d \in D$, if $d \cap \uparrow x^\circ \neq \emptyset$ then $d \cap (\{x^\circ\} \cup \uparrow S) \neq \emptyset$.

A stable canonical rule $\rho(A, D)$ is called *trivial[•]* if for all $S \subseteq A_*$ there is $x^\bullet \in A_*$ such that

(3): $S \subseteq \uparrow x^\bullet$

(4): For all $d \in D$, if $d \cap \uparrow x^\bullet \neq \emptyset$ then $d \cap \uparrow S \neq \emptyset$.

A stable canonical rule $\rho(A, D)$ is called *trivial* iff it is both *trivial^P* and *trivial[•]*.³

Theorem 5.4. *The following are equivalent:*

(1) $\rho(A, D)$ is admissible.

(2) $\rho(A, D)$ is derivable from $\{S_n^{\ell, m} : m, n, \ell \in \omega\} \cup \{T_n^m : m, n \in \omega\}$.

(3) $\rho(A, D)$ is not trivial.

(4) $\rho(A, D)$ is not equivalent to an assumption-free rule.

Proof. (2) \Rightarrow (1): We know that all $S_n^{\ell, m}$ and T_n^m are admissible, i. e. valid in the free **K4**-algebra on infinitely many generators. Since moreover $\rho(A, D)$ is derivable from $\{S_n^{\ell, m}, T_n^m : \ell, m, n \in \omega\}$, we conclude that $\rho(A, D)$ is also valid on this algebra, i. e. is admissible.

(3) \Rightarrow (2):

Suppose $\rho(F, D)$ is not derivable from $\{S_n^{\ell, m} : m, n, \ell \in \omega\} \cup \{T_n^m : m, n \in \omega\}$ with $F = A_*$. Then there is a transitive modal space (W, R) validating all $S_n^{\ell, m}$'s and all T_n^m 's and refuting $\rho(F, D)$. The latter means that there is a stable surjective $f : W \rightarrow F$ satisfying *CDC* for D . Fix $S \subseteq F$.

We will first show that there exist x° and x^\bullet satisfying the conditions of Definition 5.3(1)-(2). In what follows we are working in the modal algebra $(W, R)^*$; all connectives and modal operators are taken in this algebra. For $s \in S$ let $p_s = W \setminus f^{-1}(s) \subseteq W$, let $q = f^{-1}(\uparrow S)$ and let $r_k = f^{-1}(k)$ for $k \in F$. Let $C = \{v_1, \dots, v_\ell\}$ be a finite set of clopens of W . Since f is continuous and F is discrete, p_s and r_k are clopens, while q is a clopen up-set in W since f is also stable. In particular, q and $\Box^+ q$ have the same underlying set. Moreover, for all $s \in S$ we have $\Box^+ q \not\subseteq p_s$. Indeed, for any $w_s \in f^{-1}(s)$ we have that $w_s \in q$ but $w_s \notin p_s = W \setminus f^{-1}(s)$. This means that the conclusion of the rule $S_n^{\ell, m}$ is falsified on W . It follows that W falsifies the premise of that rule as well. Hence there exists $w_C \in W$ such that $w_C \in \bigcap_{l=1}^\ell (\Box v_l \rightarrow v_l)$, $w_C \in \bigcap_{k \in F} \Box(r_k \rightarrow \Box(r_k \cup \Box^+ q))$ and $w_C \notin \bigcup_{s \in S} \Box p_s$. The latter can be equivalently written as $w_C \in \bigcap_{s \in S} \Diamond f^{-1}(s)$. We thus obtain that the set $\{\Box v \rightarrow v \mid v \in W^*\} \cup \{\bigcap_{k \in F} \Box(r_k \rightarrow \Box(r_k \cup \Box^+ q)) \cap \bigcap_{s \in S} \Diamond f^{-1}(s)\}$ of clopens of W has finite intersection property. Since W is compact, the intersection of all these clopens is nonempty, i. e. there is $w \in W$ that belongs to all of these clopens. That is, w belongs to all clopens of the form $\Box v \rightarrow v$ (which means that w is reflexive), and also $w \in \bigcap_{k \in F} \Box(r_k \rightarrow \Box(r_k \cup \Box^+ q))$ and $w \in \bigcap_{s \in S} \Diamond f^{-1}(s)$. By the latter, we have that for every $s \in S$ there is a w' such that wRw' and $f(w') = s$. In other words, f being stable, $\uparrow f(w) \supseteq S$. Let $x := f(w)$. Then condition (1) of Definition 5.3 is met (notice that x is reflexive because w is reflexive and f is stable). We now show that condition (2) is met as well.

Since $w \in \bigcap_{k \in F} \Box(r_k \rightarrow \Box(r_k \cup \Box^+ q))$, in particular we have $w \in \Box(r_x \rightarrow \Box(r_x \cup \Box^+ q))$. Since $w \in r_x$, we obtain that $w \in \Box(r_x \cup \Box^+ q) = \Box(\neg r_x \rightarrow \Box^+ q)$. This means that any w' such that wRw' and $f(w') \neq x$ will be necessarily in $\Box^+ q$.

Now if $d \cap \uparrow x \neq \emptyset$ for some $d \in D$, then as $x = f(w)$, by the CDC of f there is w' such that wRw' and $f(w') \in d$. Then, either $f(w') = x$ and then $f(w') \in d \cap \{x\}$, or $f(w') \neq x$ and then as we have seen $f(w') \in \uparrow S$. Thus $f(w') \in d \cap (\{x\} \cup \uparrow S)$, so that $d \cap (\{x\} \cup \uparrow S) \neq \emptyset$. This implies that $\rho(F, D)$ is *trivial^o* (putting $x^\circ = x$).

Next we show that there exists an x^\bullet satisfying the conditions of Definition 5.3(3)-(4). As above, for $s \in S$ let $p_s = W \setminus f^{-1}(s) \subseteq W$, let $q = f^{-1}(\uparrow S)$ and let $r_k = f^{-1}(k)$ for $k \in F$. Again, the conclusion of the rule T_n^m is falsified on W and consequently W falsifies the premise of that rule as

³Notice that the points x° and x^\bullet may coincide.

well. Thus there is $w \in \bigcap_{k \in F} (\diamond r_k \rightarrow \diamond(r_k \wedge \square^+ q))$ and $w \notin \bigcup_{s \in S} \square p_s$. By the latter, we have that for every $s \in S$ there is a w' such that wRw' and $f(w') = s$. In other words, f being stable, $\uparrow f(w) \supseteq S$. Let $x := f(w)$. Then condition (3) of Definition 5.3 is met. For condition (4), consider $d \in D$ such that $d \cap \uparrow f(w) \neq \emptyset$. Then, since f satisfies CDC for D , there is an $u \in W$ with wRu and $f(u) \in d$. Thus $w \in \diamond r_k$ for $k = f(u)$, as $r_k = f^{-1}(k)$; since $w \in \diamond r_k \rightarrow \diamond(r_k \wedge \square^+ q)$, there is w' such that wRw' and $w' \in r_k \cap \square^+ q$, which means in particular that $f(w') = k \in d \cap \uparrow S$, as wanted. Putting $x^\bullet = x$ we deduce that $\rho(F, D)$ is trivial[•] and hence, trivial.

(4) \Rightarrow (3):

Suppose $\rho(A, D)$ is trivial. We show that then it is equivalent to an assumption-free rule. Using Lemma 5.2, it suffices to extend any stable surjective map $f : Y \rightarrow F$ from a clopen up-set $Y \subseteq W$ of a transitive modal space (W, R) to F satisfying CDC for D to an $\bar{f} : W \rightarrow F$ with the same properties.

For $w \in W$ let $f_w = f(Y \cap \uparrow w)$. If $S \subseteq F$ is of the kind f_w , let $Y_S \subseteq W$ be $Y_S = \{w \in W \setminus Y : f_w = S\}$. We take a minimal $S \subseteq F$ such that $Y_S \neq \emptyset$ (i. e. that $S = f_w$ for some $w \notin Y$) and extend f to $Y \cup Y_S$.

Claim 5.5. $Y \cup Y_S$ is a clopen up-set.

Proof. That $Y \cup Y_S$ is an up-set follows from minimality of S . Indeed, if $x \in Y \cup Y_S$ and xRy , then either $y \in Y$ and then we are done, or, provided $y \notin Y$, then, since Y is an up-set, also $x \notin Y$, so $x \in Y_S$, i. e. $f_x = S$. Moreover $\uparrow y \subseteq \uparrow x$, hence

$$f_y = f(Y \cap \uparrow y) \subseteq f(Y \cap \uparrow x) = f_x = S,$$

so by minimality of S necessarily $f_y = S$. The latter means $y \in Y_S$, so $y \in Y \cup Y_S$. Thus $Y \cup Y_S$ is an up-set.

To show that $Y \cup Y_S$ is clopen it suffices to show that Y_S is clopen. Indeed, for any $w \in W$ we have that $w \in (\square^+(Y \rightarrow f^{-1}S)) \setminus Y$ if and only if $f(Y \cap \uparrow w) \subseteq S$ and $w \notin Y$, which by minimality of S implies that actually $f(Y \cap \uparrow w) = S$. Thus $Y_S = (\square^+(Y \rightarrow f^{-1}S)) \setminus Y$ is clopen. This finishes the proof of the claim. \square

We now extend f to \bar{f} with $Y \subsetneq \text{dom } \bar{f} = Y \cup Y_S$. Recall that, by the triviality of (F, D) , there exist two (not necessarily distinct) points s^\bullet, s° such that (i) $S \subseteq \uparrow s^\bullet$ and $d \cap \uparrow s^\bullet \neq \emptyset \Rightarrow d \cap \uparrow S \neq \emptyset$ for all $d \in D$; (ii) s° is reflexive, $S \subseteq \uparrow s^\circ$ and $d \cap \uparrow s^\circ \neq \emptyset \Rightarrow d \cap (\{s^\circ\} \cup \uparrow S) \neq \emptyset$ for all $d \in D$. We distinguish two cases, depending whether S has a reflexive root or not.

Case (I): S has a reflexive root $s \in S$. We put:

$$\bar{f}(w) = \begin{cases} f(w), & \text{if } w \in Y, \\ s, & \text{if } w \in Y_S \setminus Y. \end{cases}$$

It is easy to see that \bar{f} is stable (s is reflexive). Now we also show that \bar{f} is continuous. Indeed, for every $x \in F$ we have $\bar{f}^{-1}(x) = f^{-1}(x)$ (if $x \neq s$) or $\bar{f}^{-1}(x) = f^{-1}(x) \cup Y_S$ (if $x = s$). Since the latter is a clopen set, continuity follows. Also, \bar{f} satisfies CDC: the relevant case is when $d \cap \uparrow \bar{f}(w) \neq \emptyset$ for $d \in D$, $w \in Y_S$. We have $\bar{f}(w) = s$. But $f_w = S$ (because $w \in Y_S$), i. e. $f(Y \cap \uparrow w) = S$. Thus, there is $w' \in Y$ with wR^+w' and $f(w') = s$. Since $w \notin Y$ and $w' \in Y$, we have wRw' . We can use the fact that f satisfies the CDC: since $w' \in Y = \text{dom}(f)$ and $\uparrow \bar{f}(w) = \uparrow s = \uparrow f(w')$, we get $\uparrow f(w') \cap d \neq \emptyset$ and also $f(\uparrow w') \cap d \neq \emptyset$; as a consequence $\bar{f}(\uparrow w) \cap d$ is also not empty.

Case (II): S does not have a reflexive root. We further distinguish two sub-cases, depending whether there are irreflexive R^+ -quasi-maximal points in Y_S or not. Notice that such points form the clopen antichain $Y_S^\bullet = Y_S \setminus \downarrow Y_S$.

Subcase (II.1): suppose $Y_S^\bullet = \emptyset$, i. e. there are no irreflexive quasi-maximal points in Y_S . Then, as noted above, every point in Y_S can see a quasi-maximal reflexive point in it. We put:

$$\bar{f}(w) = \begin{cases} f(w), & \text{if } w \in Y, \\ s^\circ, & \text{if } w \in Y_S. \end{cases}$$

It is easy to see that \bar{f} is stable (s° is reflexive). Now we also show that \bar{f} is continuous. Indeed, for every $x \in F$ we have $\bar{f}^{-1}(x) = f^{-1}(x)$ (if $x \neq s^\circ$) or $\bar{f}^{-1}(x) = f^{-1}(x) \cup Y_S$ (if $x = s^\circ$). Since the latter is a clopen set, continuity follows. Also, \bar{f} satisfies CDC: the relevant case is when $d \cap \uparrow \bar{f}(w) \neq \emptyset$ for $d \in D$, $w \in Y_S$. We have $\bar{f}(w) = s^\circ$ by construction and $d \cap (\{s^\circ\} \cup \uparrow S) \neq \emptyset$ by the choice of s° . That is, either (i) $s^\circ \in d$ or (ii) there is an $s' \in S$ such that $d \cap \uparrow s' \neq \emptyset$. In case (i), we pick a quasi-maximal reflexive w' in Y_S such that wR^+w' : since $\bar{f}(w') = s^\circ$, we have that $\bar{f}(\uparrow w) \cap d$ contains s° and is not empty. In case (ii), recall that $f_w = S$ (because $w \in Y_S$), i. e. $f(Y \cap \uparrow w) = S$. So there is w' such that wR^+w' , $w' \in Y$ and $f(w') = s'$. Since $w \notin Y$, we must have wRw' . We can then apply CDC for f to get w'' such that $w'Rw''$ with $f(w'') \in d$. Thus, wRw'' by transitivity and $\bar{f}(w'') = f(w'') \in d$: again, $\bar{f}(\uparrow w) \cap d$ is not empty.

Subcase (II.2): Y_S^\bullet is not empty, i. e. Y_S has irreflexive quasi-maximal points. Here we first make a preliminary extension f_0 of f to $Y \cup Y_S^\bullet$ in order to include such points into the domain of the map. We then extend the new f_0 to \bar{f} on $Y_S \setminus \downarrow Y_S^\bullet$.

Notice that $s^\bullet \notin S$ because $S \subseteq \uparrow s^\bullet$ and S does not have a reflexive root. We put:

$$f_0(w) = \begin{cases} f(w), & \text{if } w \in Y, \\ s^\bullet, & \text{if } w \in Y_S^\bullet. \end{cases}$$

It is easy to see that f_0 is stable (points in Y_S^\bullet are irreflexive). Now we also show that f_0 is continuous. Indeed, for every $x \in F$ we have $(f_0)^{-1}(x) = f^{-1}(x)$ (if $x \neq s^\bullet$) or $(f_0)^{-1}(x) = f^{-1}(x) \cup Y_S^\bullet$ (if $x = s^\bullet$). Since the latter is a clopen set, continuity follows. Also, f_0 satisfies CDC: the relevant case is when $d \cap \uparrow f_0(w) \neq \emptyset$ for $d \in D$, $w \in Y_S^\bullet$. We have $f_0(w) = s^\bullet$ by construction. From $d \cap \uparrow s^\bullet \neq \emptyset$ we get that there is an $s' \in d \cap \uparrow S$, i. e. there is $s'' \in S$ such that $s''Rs' \in d$. Then, since $S = f_w = f(Y \cap \uparrow w)$, there is w'' such that $w'' \in Y$, wR^+w'' and $f(w'') = s''$. Since $w \notin Y$, in fact we have wRw'' and by the CDC for f ($w'' \in \text{dom}(f) = Y$), since $f(w'') = s''Rs' \in d$, we get $f(\uparrow w'') \cap d \neq \emptyset$. Thus $\uparrow f_0(w) \cap d \supseteq \uparrow f_0(w'') \cap d = \uparrow f(w'') \cap d \neq \emptyset$.

If we compute Y_S with f_0 instead of f , we now get $Y_S' = Y_S \setminus \downarrow Y_S^\bullet$ instead of Y_S : since $s^\bullet \notin S$, for any w we will have $f_0((Y \cup Y_S^\bullet) \cap \uparrow w) = S$ if and only if $f(Y \cap \uparrow w) = S$ and $\uparrow w \cap Y_S^\bullet = \emptyset$, i. e. $w \in Y_S \setminus \downarrow Y_S^\bullet$. It follows that quasi-maximal points in Y_S' are all reflexive. We then can continue as in Subcase (II.1) above and get an extension \bar{f} .

So we extended f to \bar{f} . We need to show that by repeating this procedure we cover the whole of W . This holds since the following is true: if some $S \subseteq F$ has been used for further extension of the map according to the above procedure, then this same S can never occur again during any subsequent extensions.

Indeed let f^k , resp. f^n be any further extensions of f to Y^k , resp. Y^n , $k < n$. Suppose we have used some S for f^k ; then it cannot happen that S can be also used for f^n .

Suppose, to the contrary, that S occurs as one of the candidates to build f^n . Then in particular $S = f_w^{n-1}$ for some $w \in Y^n \setminus Y^{n-1}$. Then also $w \notin Y^k$, so $S \neq f_w^{k-1}$ (since $Y^k = Y^{k-1} \cup Y_S^{k-1}$ and Y_S^{k-1} consists precisely of those $v \notin Y^{k-1}$ for which $f_v^{k-1} = S$). In fact by minimality of S , f_w^{k-1} cannot be included in S , so $f_w^{k-1} \setminus S$ is nonempty. Now note that since f^{n-1} is an extension of f^{k-1} , one has $f_w^{n-1} \supseteq f_w^{k-1}$, hence also $f_w^{n-1} \setminus S$ is nonempty, which contradicts the equality $S = f_w^{n-1}$ above.

It thus follows that after each next extension at least one subset of F is excluded from all subsequent extension steps. Thus after some step n there will be no $w \notin Y^n$ and no S left with the property $f_w^n = S$. Which just means that there is no w outside Y^n , i. e. $Y^n = W$.

(1) \Rightarrow (4): The proof is exactly the same as in Theorem 3.4.

□

Corollary 5.6. *A canonical rule $\rho(F, D)$ has the following dichotomy property: it is either admissible or equivalent to an assumption-free rule.*

Corollary 5.7. *Admissibility is decidable for **K4**.*

Proof. The proof is similar to the proof of Corollary 3.7. □

Corollary 5.8. *The rules $\{S_n^{\ell, m} : m, n \in \omega\} \cup \{T_n^m : m \in \omega\}$ form an admissible basis for **K4**.*

Proof. The proof is similar to the proof of Corollary 3.8. □

To conclude, we mention that the above results also hold for **S4**, with the following modifications: (i) rules (T_n^m) should be removed from the admissible basis; (ii) rules $(S_n^{\ell, m})$ are kept, but can be simplified (we do not need the parameter ℓ either, because the conjuncts $\bigwedge_{l=1}^{\ell} (\Box v_l \rightarrow v_l)$ are now valid formulas); (iii) in Definition 5.3, conditions (3)-(4) are removed (thus a stable canonical rule is trivial in the new **S4** sense iff it was just trivial^o in the old sense).

Remark 5.9. Recall that an algebra P in a variety \mathcal{V} is called *projective* if for any surjective homomorphism $p : A \rightarrow B$ of \mathcal{V} -algebras and any homomorphism $b : P \rightarrow B$ in the diagram

$$\begin{array}{ccc} & & A \\ & \nearrow a & \downarrow p \\ P & \xrightarrow{b} & B \end{array}$$

there exists a lift, i. e. a homomorphism $a : P \rightarrow A$ with $b = pa$. It is well known that free algebras are projective, that a retract of a projective algebra is projective, and that an algebra is projective if and only if it is a retract of a free algebra.

For modal and Heyting algebras we can generalise the notion of projectivity to D -projectivity. We will discuss only the modal **K4**-case here. Let (P, D) be a pair where P is a **K4**-algebra and $D \subseteq P$. For brevity, let us call a map $h : P \rightarrow A$ a *D -morphism* if h is a stable homomorphism satisfying CDC for D . We will denote D -morphisms by $h : P \rightsquigarrow A$.

For a subset $D \subseteq P$ of a **K4**-algebra P we will call the algebra P *D -projective* if any diagram

$$\begin{array}{ccc} & & A \\ & \nearrow a & \downarrow p \\ P & \rightsquigarrow b & B \end{array}$$

of **K4**-algebras has a D -lift, that is, for any surjective modal homomorphism p and any D -morphism b there is a D -morphism a with $pa = b$. It can be shown that P is D -projective if and only if it is a D -retract of a free **K4**-algebra. The latter means that there exists a modal homomorphism $p : F \rightarrow P$ from a free **K4**-algebra to P and a D -morphism $f : P \rightsquigarrow F$ with $pf = id_P$.

Then our main theorem 5.4 is nothing but a characterisation of finite D -projective **K4**-algebras. Namely it follows from the main theorem that for a finite **K4**-algebra P and $D \subseteq P$, TFAE: (1) P is D -projective, (2) $\rho(P, D)$ is not admissible, (3) The dual of P satisfies the triviality conditions of Definition 5.3. Thus, in terms of D -projectivity we have the following dichotomy property: for any finite **K4**-algebra P and any subset $D \subseteq P$, the stable canonical rule $\rho(P, D)$ is not admissible if and only if P is D -projective.

Remark 5.10. Admissibility and unification over the basic (non-transitive) modal logic **K** are long-standing open problems. Although the proofs of this paper do not apply to **K** directly, we note that unlike the canonical rules of [17], stable canonical rules axiomatize consequence relations over

K. It remains open whether stable canonical rules could be applicable in analysing admissibility for non-transitive logics. In particular, whether they could be used in obtaining some dichotomy property for **K**.

REFERENCES

- [1] G. Bezhanishvili, N. Bezhanishvili, and R. Iemhoff. Stable canonical rules. *J. Symbolic Logic*. To appear.
- [2] G. Bezhanishvili, N. Bezhanishvili, and J. Ilin. Cofinal stable logics. Submitted, 2015.
- [3] G. Bezhanishvili, R. Mines, and P. J. Morandi. Topo-canonical completions of closure algebras and Heyting algebras. *Algebra Universalis*, 58(1):1–34, 2008.
- [4] N. Bezhanishvili and S. Ghilardi. Multiple-conclusion rules, hypersequents syntax and step frames. In *Advances in Modal Logic (AiML 2014)*, 2014. To appear. An extended version available as ILLC Prepublication Series Report PP-2014-05.
- [5] P. Blackburn, M. de Rijke, and Y. Venema. *Modal Logic*. Cambridge University Press, 2001.
- [6] A. Chagrov and M. Zakharyashev. *Modal Logic*. The Clarendon Press, 1997.
- [7] B. A. Davey and H. A. Priestley. *Introduction to lattices and order*. Cambridge University Press, New York, second edition, 2002.
- [8] L. Esakia. Topological Kripke models. *Soviet Math. Dokl.*, 15:147–151, 1974.
- [9] H. Friedman. One hundred and two problems in mathematical logic. *J. Symbolic Logic*, 40:113–129, 1975.
- [10] S. Ghilardi. Unification in intuitionistic logic. *J. Symbolic Logic*, 64(2):859–880, 1999.
- [11] S. Ghilardi. Best solving modal equations. *Ann. Pure Appl. Logic*, 102(3):183–198, 2000.
- [12] S. Ghilardi. Continuity, freeness, and filtrations. *Journal of Applied Non-Classical Logics*, 20(3):193–217, 2010.
- [13] J. Goudsmit. *Intuitionistic Rules, Admissible Rules of Intermediate Logics*. PhD thesis, University of Utrecht, 2015.
- [14] R. Iemhoff. On the admissible rules of intuitionistic propositional logic. *J. Symbolic Logic*, 66(1):281–294, 2001.
- [15] R. Iemhoff. A note on consequence. Submitted, 2015.
- [16] E. Jeřábek. Independent bases of admissible rules. *Log. J. IGPL*, 16(3):249–267, 2008.
- [17] E. Jeřábek. Canonical rules. *J. Symbolic Logic*, 74(4):1171–1205, 2009.
- [18] M. Kracht. *Tools and techniques in modal logic*, volume 142 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., 1999.
- [19] H. Rasiowa and R. Sikorski. *The mathematics of metamathematics*. Monografie Matematyczne, Tom 41. Państwowe Wydawnictwo Naukowe, Warsaw, 1963.
- [20] P. Rozière. *Regles admissibles en calcul propositionnel intuitionniste*. PhD thesis, Université Paris VII, 1992.
- [21] V. Rybakov. *Admissibility of logical inference rules*, volume 136 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 1997.
- [22] V. V. Rybakov. Bases of admissible rules of the logics S4 and Int. *Algebra i Logika*, 24(1):87–107, 123, 1985.
- [23] V. V. Rybakov. *Admissibility of logical inference rules*, volume 136 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, 1997.
- [24] V. V. Rybakov. Construction of an explicit basis for rules admissible in modal system S4. *MLQ Math. Log. Q.*, 47(4):441–446, 2001.
- [25] Y. Venema. Algebras and coalgebras. In P. Blackburn, J. van Benthem, and F. Wolter, editors, *Handbook of modal logic*, volume 3 of *Studies in Logic and Practical Reasoning*, pages 331–426. Elsevier, 2007.
- [26] M. Zakharyashev. Canonical formulas for K4. I. Basic results. *J. Symbolic Logic*, 57(4):1377–1402, 1992.