


Institute for Logic, Language and Computation

HOW TO BROADEN YOUR HORIZON

Harold Schellinx

ILLC Prepublication Series
for Mathematical Logic and Foundations ML-92-07



University of Amsterdam

The ILLC Prepublication Series

1990

Logic, Semantics and Philosophy of Language

- LP-90-01 Jaap van der Does A Generalized Quantifier Logic for Naked Infinitives
 LP-90-02 Jeroen Groenendijk, Martin Stokhof Dynamic Montague Grammar
 LP-90-03 Renate Bartsch Concept Formation and Concept Composition
 LP-90-04 Aarne Ranta Intuitionistic Categorical Grammar
 LP-90-05 Patrick Blackburn Nominal Tense Logic
 LP-90-06 Gennaro Chierchia The Variability of Impersonal Subjects
 LP-90-07 Gennaro Chierchia Anaphora and Dynamic Logic
 LP-90-08 Herman Hendriks Flexible Montague Grammar
 LP-90-09 Paul Dekker The Scope of Negation in Discourse, towards a Flexible Dynamic Montague grammar
 LP-90-10 Theo M.V. Janssen Models for Discourse Markers
 LP-90-11 Johan van Benthem General Dynamics
 LP-90-12 Serge Lapierre A Functional Partial Semantics for Intensional Logic
 LP-90-13 Zhisheng Huang Logics for Belief Dependence
 LP-90-14 Jeroen Groenendijk, Martin Stokhof Two Theories of Dynamic Semantics
 LP-90-15 Maarten de Rijke The Modal Logic of Inequality
 LP-90-16 Zhisheng Huang, Karen Kwast Awareness, Negation and Logical Omniscience
 LP-90-17 Paul Dekker Existential Disclosure, Implicit Arguments in Dynamic Semantics

Mathematical Logic and Foundations

- ML-90-01 Harold Schellinx Isomorphisms and Non-Isomorphisms of Graph Models
 ML-90-02 Jaap van Oosten A Semantical Proof of De Jongh's Theorem
 ML-90-03 Yde Venema Relational Games
 ML-90-04 Maarten de Rijke Unary Interpretability Logic
 ML-90-05 Domenico Zambella Sequences with Simple Initial Segments
 ML-90-06 Jaap van Oosten Extension of Lifschitz' Realizability to Higher Order Arithmetic, and a Solution to a Problem of F. Richman
 ML-90-07 Maarten de Rijke A Note on the Interpretability Logic of Finitely Axiomatized Theories
 ML-90-08 Harold Schellinx Some Syntactical Observations on Linear Logic
 ML-90-09 Dick de Jongh, Duccio Pianigiani Solution of a Problem of David Guaspari
 ML-90-10 Michiel van Lambalgen Randomness in Set Theory
 ML-90-11 Paul C. Gilmore The Consistency of an Extended NaDSet

Computation and Complexity Theory

- CT-90-01 John Tromp, Peter van Emde Boas Associative Storage Modification Machines
 CT-90-02 Sieger van Denneheuvel, Gerard R. Renardel de Lavalette A Normal Form for PCSJ Expressions
 CT-90-03 Ricard Gavaldà, Leen Torenvliet, Osamu Watanabe, José L. Balcázar Generalized Kolmogorov Complexity in Relativized Separations
 CT-90-04 Harry Buhman, Edith Spaan, Leen Torenvliet Bounded Reductions
 CT-90-05 Sieger van Denneheuvel, Karen Kwast Efficient Normalization of Database and Constraint Expressions
 CT-90-06 Michiel Smid, Peter van Emde Boas Dynamic Data Structures on Multiple Storage Media, a Tutorial
 CT-90-07 Kees Doets Greatest Fixed Points of Logic Programs
 CT-90-08 Fred de Geus, Ernest Rotterdam, Sieger van Denneheuvel, Peter van Emde Boas Physiological Modelling using RL
 CT-90-09 Roel de Vrijer Unique Normal Forms for Combinatory Logic with Parallel Conditional, a case study in conditional rewriting

Other Prepublications

- X-90-01 A.S. Troelstra Remarks on Intuitionism and the Philosophy of Mathematics, Revised Version
 X-90-02 Maarten de Rijke Some Chapters on Interpretability Logic
 X-90-03 L.D. Beklemishev On the Complexity of Arithmetical Interpretations of Modal Formulae
 X-90-04 Annual Report 1989
 X-90-05 Valentin Shehtman Derived Sets in Euclidean Spaces and Modal Logic
 X-90-06 Valentin Goranko, Solomon Passy Using the Universal Modality: Gains and Questions
 X-90-07 V.Yu. Shavrukov The Lindenbaum Fixed Point Algebra is Undecidable
 X-90-08 L.D. Beklemishev Provability Logics for Natural Turing Progressions of Arithmetical Theories
 X-90-09 V.Yu. Shavrukov On Rosser's Provability Predicate
 X-90-10 Sieger van Denneheuvel, Peter van Emde Boas An Overview of the Rule Language RL/1
 X-90-11 Alessandra Carbone Provable Fixed points in $\Lambda_0 + \Omega_1$, revised version
 X-90-12 Maarten de Rijke Bi-Unary Interpretability Logic
 X-90-13 K.N. Ignatiev Dzhaparidze's Polymodal Logic: Arithmetical Completeness, Fixed Point Property, Craig's Property
 X-90-14 L.A. Chagrova Undecidable Problems in Correspondence Theory
 X-90-15 A.S. Troelstra Lectures on Linear Logic


1991

Logic, Semantics and Philosophy of Language

- LP-91-01 Wiebe van der Hoek, Maarten de Rijke Generalized Quantifiers and Modal Logic
 LP-91-02 Frank Veltman Defaults in Update Semantics
 LP-91-03 Willem Groeneveld Dynamic Semantics and Circular Propositions
 LP-91-04 Makoto Kanazawa The Lambek Calculus enriched with Additional Connectives
 LP-91-05 Zhisheng Huang, Peter van Emde Boas The Schoenmakers Paradox: Its Solution in a Belief Dependence Framework
 LP-91-06 Zhisheng Huang, Peter van Emde Boas Belief Dependence, Revision and Persistence
 LP-91-07 Henk Verkuyl, Jaap van der Does The Semantics of Plural Noun Phrases
 LP-91-08 Víctor Sánchez Valencia Categorical Grammar and Natural Reasoning
 LP-91-09 Arthur Nieuwendijk Semantics and Comparative Logic
 LP-91-10 Johan van Benthem Logic and the Flow of Information

Mathematical Logic and Foundations

- ML-91-01 Yde Venema Cylindric Modal Logic
 ML-91-02 Alessandro Berarducci, Rineke Verbrugge On the Metamathematics of Weak Theories
 ML-91-03 Domenico Zambella On the Proofs of Arithmetical Completeness for Interpretability Logic
 ML-91-04 Raymond Hoofman, Harold Schellinx Collapsing Graph Models by Preorders
 ML-91-05 A.S. Troelstra History of Constructivism in the Twentieth Century
 ML-91-06 Inge Bethke Finite Type Structures within Combinatory Algebras
 ML-91-07 Yde Venema Modal Derivation Rules
 ML-91-08 Inge Bethke Going Stable in Graph Models
 ML-91-09 V.Yu. Shavrukov A Note on the Diagonalizable Algebras of PA and ZF



Institute for Logic, Language and Computation

Plantage Muidergracht 24

1018TV Amsterdam

Telephone 020-525.6051, Fax: 020-525.5101

HOW TO BROADEN YOUR HORIZON

Harold Schellinx

Department of Mathematics and Computer Science
University of Amsterdam

How to broaden your horizon*

by

Harold Schellinx

Department of Mathematics and Computer Science

University of Amsterdam

[harold@fwi.uva.nl]

Abstract

We generalize the proofs in Meyer and Ono(1992) of the finite model property for **BCK** and **BCIW** to an infinite collection of extensions of **BCI** with a ‘knotted’ rule of contraction or expansion. As a corollary we get that **BCI** has the finite model property (with respect to the class of models under consideration) only if it is equal to an intersection of these extensions.

1 Introduction

Linear implicational logic or **BCI** is the fragment of intuitionistic implicational logic obtained by deleting the structural rules of weakening and contraction from the formulation of intuitionistic implicational logic as a sequent-calculus. So we have the identity axiom:

$$[\text{ax}] \quad A \Rightarrow A,$$

the logical rules:

$$[\supset R] \quad \frac{\Gamma, A \Rightarrow B}{\Gamma \Rightarrow AB} \quad [\supset L] \quad \frac{\Gamma_1 \Rightarrow A \quad \Gamma_2, B \Rightarrow C}{\Gamma_1, \Gamma_2, AB \Rightarrow C}$$

and the cut-rule:

$$[\text{cut}] \quad \frac{\Gamma_1 \Rightarrow A \quad \Gamma_2, A \Rightarrow B}{\Gamma_1, \Gamma_2 \Rightarrow B}.$$

(Here Γ, Γ_i denote *multisets* (so *exchange* is implicit), while AB abbreviates $A \supset B$.)

*EXERCISE: Explain the title.

An equivalent Hilbert-type formulation is given by the axiom-schemes

$$\begin{array}{ll} \mathbf{[B]} & (\mathbf{BC})((\mathbf{AB})(\mathbf{AC})) \\ \mathbf{[C]} & (\mathbf{A}(\mathbf{BC}))(\mathbf{B}(\mathbf{AC})) \\ \mathbf{[I]} & \mathbf{AA} \end{array}$$

and *modus ponens* as a single rule of inference.

If we extend **BCI** with the structural rule of *weakening*

$$[\text{weak}] \quad \frac{\Gamma \Rightarrow B}{\Gamma, A \Rightarrow B}$$

(for the Hilbert-system with the axiom-scheme **[K]** : $A(\mathbf{BA})$) we obtain the implicational fragment of *affine* logic, a.k.a.¹ **BCK**.

If we extend **BCI** with the structural rule of *contraction*

$$[\text{cont}] \quad \frac{\Gamma, A, A \Rightarrow B}{\Gamma, A \Rightarrow B}$$

(for the Hilbert-system with the axiom-scheme **[W]** : $(\mathbf{A}(\mathbf{AB}))(\mathbf{AB})$) we get the implicational fragment of the *relevant* logic \mathcal{R} , a.k.a. **BCIW**.

Let us write A^n to denote n copies of some formula² A . We then can define for any pair of natural numbers (n, k) a structural rule as follows:

$$[n \rightsquigarrow k] \quad \frac{\Gamma, A^n \Rightarrow B}{\Gamma, A^k \Rightarrow B}$$

We will denote the logical system thus obtained by \mathbf{BCI}_k^n . Note that \mathbf{BCI}_k^n can equivalently be obtained by extending the Hilbert-system for **BCI** with an axiom-scheme $[n \rightsquigarrow k]$ defined as

$$\underbrace{(X \supset (X \supset (\dots (X \supset A) \dots)))}_n \supset \underbrace{(X \supset (X \supset \dots (X \supset A) \dots))}_k,$$

(which for $n = 0$ should be read as $A \supset (X \supset (X \supset \dots (X \supset A) \dots))$, for $k = 0$ as $(X \supset (X \supset \dots (X \supset A) \dots)) \supset A$.)

Clearly \mathbf{BCI}_k^k is just **BCI**, while \mathbf{BCI}_0^n for $n \neq 0$ is weird, to say the least: in \mathbf{BCI}_0^1 any formula ϕ is derivable; also, for all $n > 0$ the instance $(p^n \supset p) \supset p$ of the axiom-scheme $[n \rightsquigarrow 0]$ is not valid classically. On the other hand, for all pairs (n, k) such that $k = 0 \rightarrow n = 0$ what we have is an *extension* of **BCI** that is a fragment of intuitionistic implicational logic. In fact, for $n < k$ each logic

¹I.e. *also known as*

²By 'formula' we will of course always mean an *implicational* formula.

\mathbf{BCI}_k^n is a fragment of \mathbf{BCI}_1^0 (which is just \mathbf{BCK}), while for $n > k > 0$ clearly \mathbf{BCI}_k^n is a fragment of \mathbf{BCI}_1^2 (which is just \mathbf{BCIW}).

In Meyer and Ono(1992) it is shown that \mathbf{BCK} and \mathbf{BCIW} have the *finite model property* with respect to a certain class of ordered monoids. The purpose of this note is to show that Meyer and Ono's proof of the finite model property for \mathbf{BCK} uniformly generalizes to a proof for \mathbf{BCI}_k^n , for all $n < k$, and that their proof of the finite model property for \mathbf{BCIW} uniformly generalizes to a proof for \mathbf{BCI}_k^n , for all $n > k$. I.e. we will show that for a suitable class of structures, if a formula ϕ is valid on all its *finite* members, then it is derivable in \mathbf{BCI}_k^n . Crucial in this generalization is the construction of *finite* so-called \mathbf{BCI}_k^n -monoids in section 3.

All these logics in fact are *proper* extensions of \mathbf{BCI} (it is easy to show that $\mathbf{BCI} \vdash \Rightarrow [n \rightsquigarrow k]$ iff $n = k$) and it seems not too farfetched to conjecture that \mathbf{BCI} is their intersection. In the final section we observe that it is an easy corollary of our proof of the finite model property for all the extensions that in fact this is equivalent to the statement that \mathbf{BCI} *itself* has the finite model property (with respect to this class of models).

2 \mathbf{BCI}_k^n : structures, validity and completeness

We begin by introducing the notion of \mathbf{BCI}_k^n -monoid, being an obvious generalization of the \mathbf{BCI} -, \mathbf{BCK} -, and \mathbf{BCIW} -structures of Meyer and Ono(1992).

2.1. DEFINITION. A \mathbf{BCI} -structure is a quadruple $(\mathbf{M}, \cdot, 1, \leq)$, with \mathbf{M} a set and $1 \in \mathbf{M}$, such that \cdot is a binary operation on \mathbf{M} that is monotonous with respect to the binary relation \leq on \mathbf{M} (i.e. for all $x, y, z \in \mathbf{M}$ we have that $x \leq y$ implies $x \cdot z \leq y \cdot z$) and $(\mathbf{M}, \cdot, 1)$ a commutative monoid with unity 1.

A \mathbf{BCI}_k^n -monoid is a \mathbf{BCI} -structure of which we moreover demand that, for all $x \in \mathbf{M}$, $x^n \leq x^k$ (taking x^0 to denote the unity 1). \square

Note that we do not ask anything special of the relation \leq . However, we observe the following.

2.2. LEMMA. A finite \mathbf{BCI} -structure (\mathcal{X}, \leq) is both a \mathbf{BCI}_k^n -monoid and a \mathbf{BCI}_n^k -monoid for some $(n \neq k)$ if and only if \leq is reflexive.

PROOF: If \mathcal{X} is finite then there are $n \neq k$ such that $x^n = x^k$, for all x . By reflexivity then both $x^n \leq x^k$ and $x^k \leq x^n$, for all x . For the converse, observe that all \mathbf{BCI}_k^n -monoids are reflexive. \square

We define *validity* on \mathbf{BCI}_k^n -monoids just as validity on \mathbf{BCI} -structures: a

valuation \models on a \mathbf{BCI}_k^n -monoid is a relation between elements of \mathbf{M} and propositional variables satisfying the monotonicity condition

$$[\text{mon}] \quad x \models p \quad \text{and} \quad x \leq y \quad \text{implies} \quad y \models p.$$

Each valuation has a canonical extension to a relation between elements of \mathbf{M} and implicational formulas by $x \models AB$ iff $\forall y. y \models A \Rightarrow xy \models B$.

A formula ϕ is said to be *valid* on a given \mathbf{BCI}_k^n -monoid \mathcal{X} (written as $\mathcal{X} \models \phi$) if $1 \models \phi$ for all valuations \models .

With respect to validity on \mathbf{BCI} -structures we observe:

2.3. LEMMA. *A formula ϕ is valid on all (finite) \mathbf{BCI} -structures iff it is valid on all (finite) reflexive (even: all discrete) \mathbf{BCI} -structures.*

PROOF: Suppose $(\mathcal{X}, \leq) \not\models \phi$ for some valuation \models . Define \models^* on $(\mathcal{X}, =)$ by $x \models^* p$ iff $x \models p$, for propositional variables p . Then obviously $x \models^* \psi$ iff $x \models \psi$ for any formula ψ . Therefore $1 \not\models^* \phi$, so ϕ is not valid on a reflexive (more so, a discrete) \mathbf{BCI} -structure. \square

2.4. REMARK. It seems that in general the condition $x^n \leq x^k$ does *not* ensure validity of $[n \rightsquigarrow k]$, so in fact we can not guarantee *soundness* of *all* logics \mathbf{BCI}_k^n with respect to validity in \mathbf{BCI}_k^n -monoids. The reader might want to verify however that soundness *does* hold in case $k = 1$.

2.5. DEFINITION. Let Γ, Δ be multisets of formulas. We say that $\Gamma \stackrel{n}{|}_k \Delta$ iff $\Delta \Rightarrow x$ is obtainable from $\Gamma \Rightarrow x$ by means of some (maybe no) applications of rule $[n \rightsquigarrow k]$ (x being an arbitrary formula). \square

So $\stackrel{n}{|}_k$ is a partial order on the collection of multisets of formulas. Note that for $k = n$ we have $\Gamma \stackrel{n}{|}_k \Delta$ iff $\Gamma = \Delta$. Also obviously $\Gamma \stackrel{n}{|}_k \Delta$ iff $\Delta \stackrel{k}{|}_n \Gamma$. So Γ and Δ are $\stackrel{n}{|}_k$ -comparable if and only if they are $\stackrel{k}{|}_n$ -comparable.

2.6. PROPOSITION. *\mathbf{BCI}_k^n -logic is complete with respect to the class of all \mathbf{BCI}_k^n -monoids, for any n, k .*

PROOF: (This is standard, and as in Meyer and Ono(1992). We give an outline of the proof, for we will need some of the notions and details later on.)

Suppose $\mathbf{BCI}_k^n \not\vdash A$. Let $S = \{S_1, \dots, S_m\}$ be the set of all subformulas of A . Let Σ denote the collection of all finite multisets with elements in S .

(Note that each element can be unambiguously represented by a vector $\langle a_1, \dots, a_m \rangle$ with entries in the set \mathbb{N} of natural numbers, and a_i denoting the number of occurrences of S_i . Clearly, $(\Sigma, \cup, \emptyset, \stackrel{n}{|}_k)$ is a \mathbf{BCI}_k^n -monoid, with as isomorphic representation the set of all m -dimensional vectors with entries in

\mathbb{N} , pointwise addition as operation, $\mathbf{0} := \langle 0, \dots, 0 \rangle$ as unity and the obvious interpretation of the relation \models_k^n .)

Then, for propositional variables p appearing in A , define

$$\Gamma \models p \quad \text{iff} \quad \mathbf{BCI}_k^n \vdash \Gamma \Rightarrow p.$$

One then shows by induction on the complexity of $S_i \in S$, that for all subformulas S_i of A we have that

$$\Gamma \models S_i \quad \text{iff} \quad \mathbf{BCI}_k^n \vdash \Gamma \Rightarrow S_i.$$

(If $\Gamma \models B \supset C$, note that by induction hypothesis $\{B\} \models B$, so $\Gamma, B \models C$, a.w.a.d.³ by induction hypothesis. Conversely, if $\mathbf{BCI}_k^n \vdash \Gamma \Rightarrow B \supset C$ and $\Delta \models B$, then $\mathbf{BCI}_k^n \vdash \Delta \Rightarrow B$ by inductive hypothesis. So $\mathbf{BCI}_k^n \vdash \Gamma, \Delta \Rightarrow C$ by an application of cut, and $\Gamma, \Delta \models C$ by inductive hypothesis.)

We conclude that $\emptyset \not\models A$. \square

3 Finite \mathbf{BCI}_k^n -monoids

We are going to construct finite \mathbf{BCI}_k^n -monoids on initial segments of the natural numbers, for any $n \neq k$.

Let $\text{mod}(a, b)$ denote the remainder of a on division by b . For any pair of natural numbers $R \geq 0$ and $r \geq 1$ define an operation $[R, \cdot]_r$ on natural numbers as follows:

$$[R, a]_r = \begin{cases} a - R & \text{if } a < R \\ \text{mod}(a - R, r) & \text{otherwise.} \end{cases}$$

It is easy to see that then $R + [R, a]_r$ maps the natural numbers onto the initial segment $\{0, 1, \dots, R + r - 1\}$; moreover, it is the identity on this segment, as for $a \leq R + r - 1$ we have that $[R, a]_r = a - R$.

We define on $\{0, 1, \dots, R + r - 1\}$ an operation \oplus_r by:

$$a \oplus_r b = R + [R, a + b]_r.$$

3.1. PROPOSITION. $(\{0, 1, \dots, R + r - 1\}, \oplus_r, 0)$ is a commutative monoid with unity 0.

PROOF: Commutativity is clear because of commutativity of $+$; neutrality of 0 follows from the remarks above. So it remains to show associativity of \oplus_r .

³I.e. and we are done

By definition $(a \oplus_r b) \oplus_r c = R + [R, (a \oplus_r b) + c]_r$. First suppose that $a + b + c < R$. Then obviously $a + b < R$, so $(a \oplus_r b) \oplus_r c = R + [R, a + b + c]_r$. Otherwise $a + b + c \geq R$. If $a + b < R$ we again have $(a \oplus_r b) \oplus_r c = R + [R, a + b + c]_r$. So let us suppose that $a + b \geq R$. Then

$$\begin{aligned} (a \oplus_r b) \oplus_r c &= R + [R, R + [R, a + b]_r + c]_r \\ &= R + \text{mod}(\text{mod}(a + b - R, r) + c, r) \\ &= R + \text{mod}(a + b + c - R, r) \\ &= R + [R, a + b + c]_r \end{aligned}$$

So $(a \oplus_r b) \oplus_r c = R + [R, a + b + c]_r$ in all cases. Similarly one shows that $a \oplus_r b \oplus_r c = R + [R, a + b + c]_r$, a.w.a.d. \square

The following two lemma's give some properties of the operation \oplus_r .

3.2. LEMMA. *Let $a, b, h \in \{0, 1, \dots, R + r - 1\}$. Then (i) $(a + b) - (a \oplus_r b) \equiv 0 \pmod r$; (ii) if $h \leq (a \oplus_r b)$ and $(a + b) - h \equiv 0 \pmod r$, also $(a \oplus_r b) - h \equiv 0 \pmod r$.*

PROOF: For (i), either $a \oplus_r b = a + b$, or $a \oplus_r b = R + \text{mod}(a + b - R, r)$. For (ii), just note that $(a \oplus_r b) - h = (a + b) - h - ((a + b) - (a \oplus_r b))$ and use (i). \square

Let $\oplus_r^n a$ stand for $\underbrace{a \oplus_r a \oplus_r \dots \oplus_r a}_n$.

3.3. LEMMA. *For all $x, a \in \{0, 1, \dots, R + r - 1\}$ we have $(\oplus_r^n a) \oplus_r x \geq x$; moreover $((\oplus_r^n a) \oplus_r x) - x \equiv 0 \pmod r$.*

PROOF: By the above $\oplus_r^n a$ equals either na or $R + \text{mod}(na - R, r)$ which in turn equals $R + \text{mod}(-R, r)$. In the first case we obtain $x \oplus_r na$, which either equals $x + na \geq x$, or $R + \text{mod}(x + na - R, r) = R + \text{mod}(x - R, r)$.

In the second case we obtain

$$\begin{aligned} x \oplus_r (R + \text{mod}(-R, r)) &= R + \text{mod}(x + R + \text{mod}(-R, r) - R, r) \\ &= R + \text{mod}(x + \text{mod}(-R, r), r) \\ &= R + \text{mod}(x - R, r). \end{aligned}$$

It therefore suffices to show that for any $x \in \{0, 1, \dots, R + r - 1\}$ we have that $x \leq R + \text{mod}(x - R, r)$. If $x < R$ this is obvious. Otherwise $R \leq x \leq R + r - 1$ and $R + \text{mod}(x - R, r) = R + x - R = x$.

The second claim follows by noting that $\text{mod}(x - R, r) = r - \text{mod}(R - x, r)$, so $R + \text{mod}(x - R, r) - x = (R - x) - \text{mod}(R - x, r) + r \equiv 0 \pmod r$. \square

We will write \mathcal{M} for $(\{0, 1, \dots, R + r - 1\}, \oplus_r, 0)$, suppose $r = |n - k|$ for some $n \neq k$ and consider the elements of \mathcal{M} as representing the number of occurrences of some formula ϕ . Let Γ, Δ range over multisets having ϕ as their sole element (when non-empty). Γ, Δ then are uniquely represented by their cardinality and we have a relation $|_k^n$ between natural numbers by defining $x |_k^n y$ iff Γ, Δ have x, y elements and $\Gamma |_k^n \Delta$ in the sense of definition 2.5.

So the following hold:

- if $k > n$ then $x |_k^n y$ iff $(x = y \text{ or } n \leq x \leq y \text{ and } y - x \equiv 0 \pmod{k - n})$;
- if $k < n$ then $x |_k^n y$ iff $y |_n^k x$ iff $(x = y \text{ or } k \leq y \leq x \text{ and } x - y \equiv 0 \pmod{n - k})$.

One easily verifies that for any $n \neq k$ any subset of \mathbb{N} without $|_k^n$ -comparable elements is *finite*. Moreover, for $k < n$, there obviously are no infinite ascending chains, while for $n < k$ there are no infinite descending chains. (I.e. for $n < k$ the relation $|_k^n$ is a *well-quasi-ordering* on the set of natural numbers.)

3.4. PROPOSITION. *If $\min(n, k) \leq R$, then $(\mathcal{M}, |_k^n)$ is a finite \mathbf{BCI}_k^n -monoid.*

PROOF: We show monotonicity of $|_k^n$ with respect to \oplus_r . Suppose $x |_k^n y$.

If $k > n$ and $x = y$, then $x \oplus_r z = y \oplus_r z$. So let us assume that $n \leq x \leq y$ and $(y - x) \equiv 0 \pmod{r}$. Then $y = x + ar$ for some natural number a . Now

$$x \oplus_r z = R + [R, x + z]_r \quad \text{and}$$

$$y \oplus_r z = R + [R, y + z]_r = R + [R, x + ar + z]_r.$$

If $y + z < R$ then $x \oplus_r z = x + z$, $y \oplus_r z = y + z$ a.w.a.d., as obviously $x + z |_k^n y + z$. Otherwise $y \oplus_r z = R + \text{mod}(x + ar + z - R, r) = R + \text{mod}(x + z - R, r)$. If $x + z \geq R$, then in fact $y \oplus_r z = x \oplus_r z$ a.w.a.d. Otherwise $x \oplus_r z = x + z$. Then $n \leq x \oplus_r z < y \oplus_r z$ and moreover $(y \oplus_r z) - (x \oplus_r z) = R + \text{mod}(x + z - R, r) - (x + z) = R - (x + z) - \text{mod}(R - (x + z), r) + r \equiv 0 \pmod{r}$.

So $x \oplus_r z |_k^n y \oplus_r z$ in all cases, q.e.d.

If $k < n$ we reason by duality, as in that case

$$x |_k^n y \Leftrightarrow y |_n^k x \Rightarrow y \oplus_r z |_n^k x \oplus_r z \Leftrightarrow x \oplus_r z |_k^n y \oplus_r z.$$

To complete the proof we need $(\oplus_r^n x) |_k^n (\oplus_r^k x)$ for all $x \in \mathcal{M}$. Let us assume that $k > n$. Then $(\oplus_r^k x) = (\oplus_r^n x) \oplus_r (\oplus_r^r x)$. Put $y := (\oplus_r^n x)$. We will show that $y |_k^n (\oplus_r^r x) \oplus_r y$. If $x = 0$ we have equality. Otherwise $y = nx \geq n$ or $y = R + \text{mod}(nx - R, r) \geq R \geq n$ by assumption. S.w.a.d.⁴ by lemma 3.3.

For $n > k$ we use once more duality. (Here we need the assumption that $R \geq k$). \square

⁴I.e. *So we are done*

3.5. REMARK. Though the condition “ $\min(n, k) \leq R$ ” is sufficient, it is *not* necessary. E.g. for $k > n$ we may have $n > R$, while nevertheless $(\mathcal{M}, |\frac{n}{k}|)$ is a finite \mathbf{BCI}_k^n -monoid. This can be the case if $\text{mod}(nx - R, r) > 0$, for all $nx \geq R$.

4 The finite model property

After having shown how one may obtain finite \mathbf{BCI}_k^n -monoids, we can continue exploiting Meyer and Ono(1992), in order to generalize the proofs there given of the finite model property for \mathbf{BCK} (which for us is nothing but \mathbf{BCI}_1^0) and \mathbf{BCIW} (which is \mathbf{BCI}_1^2) to a proof of the finite model property for \mathbf{BCI}_k^n for all $n < k$, respectively to a proof of the finite model property for \mathbf{BCI}_k^n for all $n > k > 0$.

As a starter, we will generalize their notion of ϕ -critical formula.

4.1. DEFINITION. Let ϕ be a formula, Γ a multiset of formulas.

- if $n < k$ we say that Γ is (ϕ, n, k) -critical iff
 - (i) $\mathbf{BCI}_k^n \vdash \Gamma \Rightarrow \phi$;
 - (ii) If $\Delta |\frac{n}{k}| \Gamma$ and $\Delta \neq \Gamma$, then $\mathbf{BCI}_k^n \not\vdash \Delta \Rightarrow \phi$.
- if $k < n$ we say that Γ is (ϕ, k, n) -critical iff
 - (i) $\mathbf{BCI}_k^n \not\vdash \Gamma \Rightarrow \phi$;
 - (ii) If $\Gamma |\frac{n}{k}| \Delta$ and $\Delta \neq \Gamma$, then $\mathbf{BCI}_k^n \vdash \Delta \Rightarrow \phi$. □

4.2. PROPOSITION. Let Σ be the collection of all finite multisets with elements from some finite set of formulas S . Then the set

$$\mathcal{R}(\phi) := \{\Gamma \in \Sigma \mid \Gamma \text{ is } (\phi, n, k)\text{-critical}\}$$

is finite, for any $n \neq k$.

PROOF: If Γ, Δ are distinct elements in $\mathcal{R}(\phi)$, then they are neither $|\frac{n}{k}|$ - nor $|\frac{k}{n}|$ -comparable. The claim then follows from the fact that for $n \neq k$ either the relation $|\frac{n}{k}|$ or the relation $|\frac{k}{n}|$ is a well-quasi-ordering, as it is equivalent to the pointwise lifting of the ‘singleton-order’ to $|S|$ -tuples. See e.g. Rosenstein(1982). □

We can now more or less copy the finite model property proofs in Meyer and Ono(1992), with just some additional arguments using properties of \oplus_r . To keep this note self-contained, we will nevertheless work through the details.

Suppose $\mathbf{BCI}_k^n \not\Rightarrow A$. Let S, Σ be as in the proof of 2.6. Then $\bigcup_{\phi \in S} \mathcal{R}(\phi)$ is *finite*, by proposition 4.2.

First we assume that $n < k$.

Let $R := \max\{n, \max\{a_i \mid \langle a_1, \dots, a_m \rangle \in \bigcup \mathcal{R}(\phi)\}\}$, where $\langle a_1, \dots, a_m \rangle$ is the vector representing the multiset $\{S_1^{a_1}, \dots, S_m^{a_m}\}$. Let $r = k - n$, and take the set \mathcal{V} of m -dimensional vectors with entries $\leq R + r - 1$. Define an operation \oplus_r on \mathcal{V} by $\langle a_1, \dots, a_m \rangle \oplus_r \langle b_1, \dots, b_m \rangle = \langle a_1 \oplus_r b_1, \dots, a_m \oplus_r b_m \rangle$, and a relation $|_k^n$ by $\langle a_1, \dots, a_m \rangle |_k^n \langle b_1, \dots, b_m \rangle$ iff $\forall i. a_i |_k^n b_i$.

(In what follows we will write \mathbf{a} for $\langle a_1, \dots, a_m \rangle$, etc.)

By the results of the previous section it is clear that $(\mathcal{V}, \oplus_r, \mathbf{0}, |_k^n)$ is a finite \mathbf{BCI}_k^n -monoid. Take the refuting valuation \models used in the proof of 2.6. Define a valuation \models^* on $(\mathcal{V}, |_k^n)$ by $\mathbf{a} \models^* p$ iff $\mathbf{a} \models p$, for any variable p in A . (Observe that \models^* satisfies [mon], because \models satisfies it.)

Then we apply induction on the complexity of $S_i \in S$ in order to show that for *all* subformulas S_i of A we have that $\mathbf{a} \models^* S_i$ iff $\mathbf{a} \models S_i$.

Let $S_i = B \supset C$. Suppose $\mathbf{a} \models B \supset C$ and $\mathbf{b} \models^* B$. Then by inductive hypothesis $\mathbf{b} \models B$, and therefore $\mathbf{a} + \mathbf{b} \models C$. Let \mathbf{c} be (C, n, k) -critical and such that $\mathbf{c} |_k^n \mathbf{a} + \mathbf{b}$. (Such \mathbf{c} has to exist as there are but finitely many vectors \mathbf{x} with the property that $\mathbf{x} |_k^n \mathbf{a} + \mathbf{b}$.)

CLAIM: $c_i |_k^n (a_i \oplus_r b_i)$. Indeed, as $c_i |_k^n a_i + b_i$, $c_i < n$ would imply that $c_i = a_i + b_i = a_i \oplus_r b_i$; otherwise, by criticality, $n \leq c_i \leq R$; if $a_i + b_i \leq R + r - 1$, then $a_i \oplus_r b_i = a_i + b_i$; otherwise $c_i \leq a_i \oplus_r b_i$ (because $R + r - 1 \geq a_i \oplus_r b_i \geq R$). The second part of lemma 3.2 then gives $(a_i \oplus_r b_i) - c_i \equiv 0 \pmod r$, a.w.a.d.

As $\mathbf{c} \models C$, by inductive hypothesis $\mathbf{c} \models^* C$, and thus $\mathbf{a} \oplus_r \mathbf{b} \models^* C$ by the above claim and [mon]. We thus showed that $\mathbf{a} \models^* B \supset C$.

Conversely, let $\mathbf{a} \models^* B \supset C$ and $\mathbf{b} \models B$. Take a (B, n, k) -critical \mathbf{c} such that $\mathbf{c} |_k^n \mathbf{b}$.

As $\forall i. c_i \leq R + r - 1$ we have $\mathbf{c} \models^* B$, so $\mathbf{a} \oplus_r \mathbf{c} \models^* C$. The inductive hypothesis then tells us that $\mathbf{a} \oplus_r \mathbf{c} \models C$.

But $a_i \oplus_r c_i |_k^n a_i + b_i$: for $c_i |_k^n b_i$ implies $a_i \oplus_r c_i |_k^n a_i \oplus_r b_i$ (by monotonicity of $|_k^n$ with respect to \oplus_r); also $a_i \oplus_r b_i |_k^n a_i + b_i$, and the result follows by transitivity of $|_k^n$. Therefore $\mathbf{a} + \mathbf{b} \models C$ by [mon], and we showed $\mathbf{a} \models B \supset C$.

But then $(\mathcal{V}, |_k^n)$ (for $n < k$) is a refuting *finite* structure for A .

We proceed to the case that $n > k$.

We now put $R := \max\{k, \max\{a_i \mid \langle a_1, \dots, a_m \rangle \in \bigcup \mathcal{R}(\phi)\}\}$ and similar to the previous case we take the finite \mathbf{BCI}_k^n -monoid $(\mathcal{V}, \oplus_r, \mathbf{0}, |_k^n)$, with $r = n - k$ and the refuting valuation \models used in the proof of 2.6. Again we define a valuation \models^* on $(\mathcal{V}, |_k^n)$ by $\mathbf{a} \models^* p$ iff $\mathbf{a} \models p$, for any variable p in A .

By induction on the complexity of $S_i \in S$ we show that $\mathbf{a} \models^* S_i$ iff $\mathbf{a} \models S_i$ for *all* subformulas S_i of A .

Let $S_i = B \supset C$, $\mathbf{a} \models B \supset C$ and $\mathbf{b} \models^* B$. By inductive hypothesis $\mathbf{b} \models B$, so $\mathbf{a} + \mathbf{b} \models C$. As $\mathbf{a} \oplus_r \mathbf{b} \mid_n^k \mathbf{a} + \mathbf{b}$, we have $\mathbf{a} + \mathbf{b} \mid_k^n \mathbf{a} \oplus_r \mathbf{b}$, and therefore ([mon]) $\mathbf{a} \oplus_r \mathbf{b} \models C$. By inductive hypothesis $\mathbf{a} \oplus_r \mathbf{b} \models^* C$, so $\mathbf{a} \models^* B \supset C$.

Conversely, suppose $\mathbf{a} \not\models B \supset C$. This means we have $\mathbf{b} \in \mathcal{V}$ such that $\mathbf{b} \models B$ and $\mathbf{a} + \mathbf{b} \not\models C$. There is a (C, k, n) -critical \mathbf{c} such that $\mathbf{a} + \mathbf{b} \mid_k^n \mathbf{c}$, i.e. $\mathbf{c} \mid_n^k \mathbf{a} + \mathbf{b}$. As before we conclude $\mathbf{c} \mid_k^n \mathbf{a} \oplus_r \mathbf{b}$, so $\mathbf{a} \oplus_r \mathbf{b} \mid_k^n \mathbf{c}$, and $\mathbf{a} \oplus_r \mathbf{b} \not\models C$. Therefore $\mathbf{a} \oplus_r \mathbf{b} \not\models^* C$ by inductive hypothesis.

Now say $\mathbf{b} = \langle b_1, \dots, b_m \rangle$. We define $\mathbf{b}' = \langle b'_1, \dots, b'_m \rangle$, where $b'_i = R + [R, b_i]_r$. Then $\mathbf{b}' \in \mathcal{V}$ and $\mathbf{b} \mid_k^n \mathbf{b}'$ (for if $b_i \leq R + r - 1$, then $b'_i = b_i$; otherwise $b'_i = R + \text{mod}(b_i - R, r)$, so $k \leq b'_i < b_i$ and $b_i - b'_i = b_i - R - \text{mod}(b_i - R, r) \equiv 0 \pmod{r}$). Moreover

$$\begin{aligned} \mathbf{a} \oplus_r \mathbf{b}' &= R + [R, a_i + b'_i]_r \\ &= R + [R, a_i + R + [R, b_i]_r]_r \\ &= R + [R, a_i + b_i]_r \\ &= \mathbf{a} \oplus_r \mathbf{b}. \end{aligned}$$

So $\mathbf{b}' \models B$ and by inductive hypothesis $\mathbf{b}' \models^* B$. However $\mathbf{a} \oplus_r \mathbf{b}' \not\models^* C$. So $\mathbf{a} \not\models^* B \supset C$.

Thus we found also for $k < n$ that (\mathcal{V}, \mid_k^n) is a refuting *finite* structure for A and we have shown:

4.3. THEOREM. (Finite model property for \mathbf{BCI}_k^n) \mathbf{BCI}_k^n is complete with respect to the collection of all finite \mathbf{BCI}_k^n -monoids, for all $n \neq k$. \square

4.4. PROPOSITION. \mathbf{BCI}_k^n is complete with respect to the collection of all finite \mathbf{BCI} -structures, i.e. if ϕ is valid on all finite \mathbf{BCI} -structures then it is derivable in \mathbf{BCI}_k^n , for any $n \neq k$.

PROOF: If ϕ is valid on all finite reflexive \mathbf{BCI} -structures, it is valid on all finite \mathbf{BCI}_k^n -monoids. \square

4.5. COROLLARY. \mathbf{BCI} has the finite model property only if it is the same as $\bigcap_{\mathcal{J}} \mathbf{BCI}_k^n$, for some $\mathcal{J} \subseteq \{(n, k) \mid n \neq k\}$. \square

4.6. REMARK. The argument given follows a quite general pattern, that might be instructive to sketch.

Let \mathcal{L} be some logic, and suppose we have a collection of logics $L_i \supset \mathcal{L}$. Let moreover classes of structures $\mathcal{M}(\mathcal{L})$ for \mathcal{L} and $\mathcal{M}(L_i)$ for L_i be given, such that $\mathcal{M}(L_i) \subset \mathcal{M}(\mathcal{L})$ and each *finite* \mathcal{L} -structure is an L_i -structure for some i . Then one easily shows:

1. \mathcal{L} has fmp⁵ $\Rightarrow \bigcap_i L_i = \mathcal{L}$;

2. Each L_i has fmp and $\bigcap_i L_i = \mathcal{L} \Rightarrow \mathcal{L}$ has fmp,

provided (for 1) that L_i is *sound*, (for 2) that L_i is *complete* with respect to the given collection of L_i -structures.

Acknowledgement

I would like to thank prof. Anne Troelstra and Andreja Prijatelj, whose work on logics with ‘knotted’ structural rules inspired this note.

References

MEYER, R. K. AND ONO, H. (1992). The finite model property for BCK and BCIW. Manuscript.

ROSENSTEIN, J. G. (1982). *Linear orderings*. Academic Press.

⁵I.e. *finite model property*

The ILLC Prepublication Series

- ML-91-10 Maarten de Rijke, Yde Venema Sahlqvist's Theorem for Boolean Algebras with Operators
 ML-91-11 Rineke Verbrugge Feasible Interpretability
 ML-91-12 Johan van Benthem Modal Frame Classes, revisited
- Computation and Complexity Theory*
 CT-91-01 Ming Li, Paul M.B. Vitányi Kolmogorov Complexity Arguments in Combinatorics
 CT-91-02 Ming Li, John Tromp, Paul M.B. Vitányi How to Share Concurrent Wait-Free Variables
 CT-91-03 Ming Li, Paul M.B. Vitányi Average Case Complexity under the Universal Distribution Equals Worst Case Complexity
- CT-91-04 Sieger van Denneheuvel, Karen Kwast Weak Equivalence
 CT-91-05 Sieger van Denneheuvel, Karen Kwast Weak Equivalence for Constraint Sets
 CT-91-06 Edith Spaan Census Techniques on Relativized Space Classes
 CT-91-07 Karen L. Kwast The Incomplete Database
 CT-91-08 Kees Doets Levationis Laus
 CT-91-09 Ming Li, Paul M.B. Vitányi Combinatorial Properties of Finite Sequences with high Kolmogorov Complexity
- CT-91-10 John Tromp, Paul Vitányi A Randomized Algorithm for Two-Process Wait-Free Test-and-Set
 CT-91-11 Lane A. Hemachandra, Edith Spaan Quasi-Injective Reductions
 CT-91-12 Krzysztof R. Apt, Dino Pedreschi Reasoning about Termination of Prolog Programs
- Computational Linguistics*
 CL-91-01 J.C. Scholtes Kohonen Feature Maps in Natural Language Processing
 CL-91-02 J.C. Scholtes Neural Nets and their Relevance for Information Retrieval
 CL-91-03 Hub Prüst, Remko Scha, Martin van den Berg A Formal Discourse Grammar tackling Verb Phrase Anaphora
- Other Prepublications*
 X-91-01 Alexander Chagrov, Michael Zakharyashev The Disjunction Property of Intermediate Propositional Logics
 X-91-02 Alexander Chagrov, Michael Zakharyashev On the Undecidability of the Disjunction Property of Intermediate Propositional Logics
 X-91-03 V. Yu. Shavrukov Subalgebras of Diagonalizable Algebras of Theories containing Arithmetic
 X-91-04 K.N. Ignatiev Partial Conservativity and Modal Logics
 X-91-05 Johan van Benthem Temporal Logic
 X-91-06 Annual Report 1990
 X-91-07 A.S. Troelstra Lectures on Linear Logic, Errata and Supplement
 X-91-08 Giorgie Dzhaparidze Logic of Tolerance
 X-91-09 L.D. Beklemishev On Bimodal Provability Logics for Π_1 -axiomatized Extensions of Arithmetical Theories
 X-91-10 Michiel van Lambalgen Independence, Randomness and the Axiom of Choice
 X-91-11 Michael Zakharyashev Canonical Formulas for K4. Part I: Basic Results
 X-91-12 Herman Hendriks Flexibele Categoriale Syntaxis en Semantiek: de proefschriften van Frans Zwarts en Michael Moortgat
 X-91-13 Max I. Kanovich The Multiplicative Fragment of Linear Logic is NP-Complete
 X-91-14 Max I. Kanovich The Horn Fragment of Linear Logic is NP-Complete
 X-91-15 V. Yu. Shavrukov Subalgebras of Diagonalizable Algebras of Theories containing Arithmetic, revised version
 X-91-16 V.G. Kanovei Undecidable Hypotheses in Edward Nelson's Internal Set Theory
 X-91-17 Michiel van Lambalgen Independence, Randomness and the Axiom of Choice, Revised Version
 X-91-18 Giovanna Cepparello New Semantics for Predicate Modal Logic: an Analysis from a standard point of view
 X-91-19 Papers presented at the Provability Interpretability Arithmetic Conference, 24-31 Aug. 1991, Dept. of Phil., Utrecht University
 Annual Report 1991
- 1992**
Logic, Semantics and Philosophy of Language
 LP-92-01 Víctor Sánchez Valencia Lambek Grammar: an Information-based Categorical Grammar
 LP-92-02 Patrick Blackburn Modal Logic and Attribute Value Structures
 LP-92-03 Szabolcs Mikulás The Completeness of the Lambek Calculus with respect to Relational Semantics
 LP-92-04 Paul Dekker An Update Semantics for Dynamic Predicate Logic
 LP-92-05 David I. Beaver The Kinematics of Presupposition
 LP-92-06 Patrick Blackburn, Edith Spaan A Modal Perspective on the Computational Complexity of Attribute Value Grammar
 LP-92-07 Jeroen Groenendijk, Martin Stokhof A Note on Interrogatives and Adverbs of Quantification
 LP-92-08 Maarten de Rijke A System of Dynamic Modal Logic
 LP-92-09 Johan van Benthem Quantifiers in the world of Types
- Mathematical Logic and Foundations*
 ML-92-01 A.S. Troelstra Comparing the theory of Representations and Constructive Mathematics
 ML-92-02 Dmitrij P. Skvortsov, Valentin B. Shehtman Maximal Kripke-type Semantics for Modal and Superintuitionistic Predicate Logics
 ML-92-03 Zoran Marković On the Structure of Kripke Models of Heyting Arithmetic
 ML-92-04 Dimiter Vakarelov A Modal Theory of Arrows, Arrow Logics I
 ML-92-05 Domenico Zambella Shavrukov's Theorem on the Subalgebras of Diagonalizable Algebras for Theories containing $IA_0 + EXP$
 ML-92-06 D.M. Gabbay, Valentin B. Shehtman Undecidability of Modal and Intermediate First-Order Logics with Two Individual Variables
 ML-92-07 Harold Schellinx How to Broaden your Horizon
- Computation and Complexity Theory*
 CT-92-01 Erik de Haas, Peter van Emde Boas Object Oriented Application Flow Graphs and their Semantics
 CT-92-02 Karen L. Kwast, Sieger van Denneheuvel Weak Equivalence: Theory and Applications
- Other prepublications*
 X-92-01 Heinrich Wansing The Logic of Information Structures
 X-92-02 Konstantin N. Ignatiev The Closed Fragment of Dzhaparidze's Polymodal Logic and the Logic of Σ_1 -conservativity
 X-92-03 Willem Groeneveld Dynamic Semantics and Circular Propositions, revised version