

# Cardinal spaces and topological representations of bimodal logics

Benedikt Löwe

*Institute for Logic, Language and Computation, Universiteit van Amsterdam,  
Plantage Muidergracht 24, 1018 TV Amsterdam, The Netherlands,  
bloewe@science.uva.nl*

Darko Sarenac

*Department of Philosophy, Stanford University, Stanford, CA 94305-2155, United  
States of America, sarenac@stanford.edu*

---

## Abstract

We look at bimodal logics interpreted by cartesian products of topological spaces and discuss the validity of certain bimodal formulae in products of so-called cardinal spaces. This solves an open problem of van Benthem *et al.*

*Keywords.* Bimodal logic, topological interpretations of modal logic, ordinals, commutativity, Church-Rosser

*2000 AMS Mathematics Subject Classification.* **54B99 03B45** 54A99 03E75.

---

## Introduction

Topological interpretations of modal logics have been introduced by McKinsey and Tarski [2] long before the advent of Kripke semantics. The authors of [1] have introduced an interpretation of bimodal logics on cartesian products of topological spaces: you have a modal language with two modalities,  $\Box_1$  and  $\Box_2$ , and interpret them as interior operators on horizontal and vertical sections of the cartesian product of two topological spaces. It is clear that both the  $\Box_1$  fragment and the  $\Box_2$  fragment satisfy the axioms of **S4**.

In [1, § 2], there is a list of results on validity of mixed formulas, in particular the mixed formulas

- $\Diamond_1 \Diamond_2 p \rightarrow \Diamond_2 \Diamond_1 p$  (**left commutativity**,  $\text{com}_{\leftarrow}$ ),

- $\diamond_2 \diamond_1 p \rightarrow \diamond_1 \diamond_2 p$  (**right commutativity**,  $\text{com}_{\rightarrow}$ ), and
- $\diamond_1 \square_2 p \rightarrow \square_2 \diamond_1 p$  (**Church-Rosser**,  $\text{chr}$ ).

**Theorem 1 (van Benthem, Bezhanishvili, ten Cate, Sarenac)** *For first-countable spaces  $\mathbf{X}$  and  $\mathbf{Y}$ , the following equivalences hold:*

- $\mathbf{X}$  is Alexandroff  $\iff \mathbf{X}, \mathbf{Y} \models \text{com}_{\leftarrow}$ ,
- $\mathbf{Y}$  is Alexandroff  $\iff \mathbf{X}, \mathbf{Y} \models \text{com}_{\rightarrow}$ , and
- at least one of  $\mathbf{X}$  and  $\mathbf{Y}$  is Alexandroff  $\iff \mathbf{X}, \mathbf{Y} \models \text{chr}$ .

Moreover, in each of the equivalences, the forward direction (“ $\implies$ ”) holds for all topological spaces.<sup>1</sup>

**Question 2 (van Benthem, Bezhanishvili, ten Cate, Sarenac)** *Do the backwards directions (“ $\impliedby$ ”) of the equivalences in Theorem 1 hold for arbitrary topological spaces  $\mathbf{X}$  and  $\mathbf{Y}$ ?*

In this note, we answer Question 2 negatively.

## Definitions

Let  $\mathbf{X} = \langle X, \tau_X \rangle$  and  $\mathbf{Y} = \langle Y, \tau_Y \rangle$  be topological spaces. If  $A \subseteq X \times Y$ ,  $x^* \in X$  and  $y^* \in Y$ , we can look at **vertical sections**  $A^{x^*} := \{y \in Y; \langle x^*, y \rangle \in A\}$  and **horizontal sections**  $A_{y^*} := \{x \in X; \langle x, y^* \rangle \in A\}$ . Vertical and horizontal sections are subsets of  $Y$  and  $X$ , respectively, and hence we can look at their closures and interiors in the spaces  $\mathbf{Y}$  and  $\mathbf{X}$ . We define the **horizontal (vertical) closure (interior)** of  $A$  as follows:<sup>2</sup>

$$\begin{aligned} \langle x, y \rangle \in \text{hcl}(A) &: \iff x \in \text{cl}_{\tau_X}(A_y), \\ \langle x, y \rangle \in \text{hint}(A) &: \iff x \in \text{int}_{\tau_X}(A_y), \\ \langle x, y \rangle \in \text{vcl}(A) &: \iff y \in \text{cl}_{\tau_Y}(A^x), \\ \langle x, y \rangle \in \text{vint}(A) &: \iff y \in \text{int}_{\tau_Y}(A^x). \end{aligned}$$

Now look at the modal language with two modalities  $\square_1$  and  $\square_2$ . The cartesian product interpretation of  $\square_1$  and  $\square_2$  is given by the following recursion<sup>3</sup>: suppose we have already defined the meaning of  $\mathbf{X}, \mathbf{Y}, x, y \models \varphi$  for all  $x \in X$

<sup>1</sup> Corollary 6.11, Proposition 6.14, Proposition 6.1, and Proposition 6.9 of [1].

<sup>2</sup> Note that these operations are closure and interior in the topology  $\text{discrete}_X \otimes \tau_Y$  and  $\tau_X \otimes \text{discrete}_Y$ , respectively.

<sup>3</sup> For details, see [1].

and  $y \in Y$ , then we let

$$\mathbf{X}, \mathbf{Y}, x, y \models \Box_1 \varphi \iff \langle x, y \rangle \in \text{hint}(\{\langle v, w \rangle; \mathbf{X}, \mathbf{Y}, v, w \models \varphi\}), \text{ and}$$

$$\mathbf{X}, \mathbf{Y}, x, y \models \Box_2 \varphi \iff \langle x, y \rangle \in \text{vint}(\{\langle v, w \rangle; \mathbf{X}, \mathbf{Y}, v, w \models \varphi\}).$$

The derived modalities  $\Diamond_1 = \neg\Box_1\neg$  and  $\Diamond_2 = \neg\Box_2\neg$  then correspond to horizontal and vertical closure. As usual, we write

$$\mathbf{X}, \mathbf{Y} \models \varphi$$

if the formula  $\varphi$  holds at all points. Consequently, for topological spaces  $\mathbf{X}$  and  $\mathbf{Y}$ , the mentioned bimodal formulae transform into the following topological statements

$$\begin{aligned} \text{com}_{\rightarrow} &\rightsquigarrow \forall A \subseteq X \times Y (\text{hcl}(\text{vcl}(A)) \subseteq \text{vcl}(\text{hcl}(A))), \\ \text{com}_{\leftarrow} &\rightsquigarrow \forall A \subseteq X \times Y (\text{vcl}(\text{hcl}(A)) \subseteq \text{hcl}(\text{vcl}(A))), \text{ and} \\ \text{chr} &\rightsquigarrow \forall A \subseteq X \times Y (\text{hcl}(\text{vint}(A)) \subseteq \text{vint}(\text{hcl}(A))), \end{aligned}$$

respectively. Note that chr is symmetric:

$$\begin{aligned} \mathbf{X}, \mathbf{Y} \models \text{chr} &\iff \mathbf{X}, \mathbf{Y} \models \Diamond_1 \Box_2 p \rightarrow \Box_2 \Diamond_1 p \\ &\iff \mathbf{Y}, \mathbf{X} \models \Diamond_2 \Box_1 p \rightarrow \Box_1 \Diamond_2 p \\ &\iff \mathbf{Y}, \mathbf{X} \models \neg \Box_2 \Diamond_1 \neg p \rightarrow \neg \Diamond_1 \Box_2 \neg p \\ &\iff \mathbf{Y}, \mathbf{X} \models \Diamond_1 \Box_2 \neg p \rightarrow \Box_2 \Diamond_1 \neg p \\ &\iff \mathbf{Y}, \mathbf{X} \models \text{chr}. \end{aligned}$$

We call a topological space **Alexandroff** if arbitrary intersections of open sets are open. Examples are the discrete or the indiscrete topologies.

As mentioned in the introduction, we are looking for non-Alexandroff spaces that validate the bimodal formulae  $\text{com}_{\rightarrow}$ ,  $\text{com}_{\leftarrow}$ , and/or chr.

For this, we define the **cardinal space** of cardinality  $\kappa$ , denoted by  $\mathbf{card}_{\kappa}$  as follows: The underlying set of the space is  $\kappa \cup \{\infty\}$  where  $\infty \notin \kappa$ . The open neighbourhood base for each  $\alpha \in \kappa$  is  $\{\{\alpha\}\}$ , the open neighbourhood base for  $\infty$  is

$$\{\{\xi; \alpha < \xi < \kappa\} \cup \{\infty\}; \alpha \in \kappa\}.$$

The topology of  $\mathbf{card}_{\kappa}$  is the discrete topology on  $\kappa$  and a point at infinity that is infinitely far away (can be reached only by sequences cofinal in  $\kappa$ ). An alternative way of viewing these spaces is as the ordinal topology on the ordinal  $\kappa + 1$  with all limit points below  $\kappa$  removed.

Note that for infinite cardinals  $\kappa$ , the cardinal space  $\mathbf{card}_{\kappa}$  is not Alexandroff, and for uncountable cardinals  $\kappa$ , it is not first-countable.

## Bimodal formulae in products of cardinal spaces

If  $\mu \leq \nu$  are ordinals, and  $\gamma \in \nu$ , we can form the Cantor Normal Form of  $\gamma$  to the base  $\mu$ :

$$\gamma = \mu^{\alpha_n} \cdot \gamma_n + \mu^{\alpha_{n-1}} \cdot \gamma_{n-1} + \dots + \mu^{\alpha_1} \cdot \gamma_1 + \gamma_0.$$

We write  $S_\mu(\gamma) := \gamma_0$  and call it the **scalar term** of  $\gamma$  to the base  $\mu$ .

**Lemma 3** *If  $\mu \leq \nu$  are cardinals,  $\gamma < \mu$  and  $\beta < \nu$ , then there is some  $\beta < \eta < \nu$  such that  $S_\mu(\eta) \geq \gamma$ .*

**Proof.** Let  $\xi := S_\mu(\beta)$ . If  $\xi \geq \gamma$ , then  $\eta := \beta + 1$  does the job. Otherwise, there is a unique  $0 < \sigma < \mu$  such that  $\gamma = \xi + \sigma$ . Let  $\eta := \beta + \sigma$ . q.e.d.

**Theorem 4** *Let  $\kappa$  and  $\lambda$  be cardinals. Then the following are equivalent:*

- (i)  $\mathbf{card}_\kappa, \mathbf{card}_\lambda \models \diamond_1 \diamond_2 p \rightarrow \diamond_2 \diamond_1 p$ , and
- (ii)  $\lambda < \text{cf } \kappa$ .

**Proof.** “(i) $\Rightarrow$ (ii)”. Suppose  $\text{cf } \kappa \leq \lambda$ . We’ll construct a subset of  $\mathbf{card}_\kappa \times \mathbf{card}_\lambda$  that constitutes a counterexample to  $\text{com}_\rightarrow$ . Let  $A = \{\alpha_\gamma; \gamma < \text{cf } \kappa\} \subseteq \kappa$  be an increasing enumeration of a cofinal subset of  $\kappa$ . We define a subset of  $\kappa \times \lambda$  as follows:

$$\langle \alpha_\gamma, \beta \rangle \in X : \iff \gamma \leq S_{\text{cf } \kappa}(\beta).$$

Note that  $S_{\text{cf } \kappa}(\beta) < \text{cf } \kappa$ , so if you fix an element  $\beta \in \lambda$  and look at the horizontal section  $X_\beta = \{\alpha; \langle \alpha, \beta \rangle \in X\}$ , then each of these sets has cardinality less than  $\text{cf } \kappa$ . In particular, none of these can be cofinal in  $\kappa$  ( $\star$ ).

Moreover, if you fix  $\alpha_\gamma \in A$  and look at the vertical section

$$X^{\alpha_\gamma} = \{\beta; \langle \alpha_\gamma, \beta \rangle \in X\},$$

then this set is cofinal in  $\lambda$  ( $\star\star$ ) by the following argument: Take an arbitrary  $\beta < \lambda$ . By Lemma 3 applied to  $\text{cf } \kappa \leq \lambda$ , we find  $\beta < \eta < \lambda$  such that  $S_{\text{cf } \kappa}(\eta) \geq \gamma$ . But that means that  $\langle \alpha_\gamma, \eta \rangle \in X$ , so  $\beta < \eta \in X^{\alpha_\gamma}$ .

By ( $\star$ ), the horizontal closure of  $X$  is  $X$  itself: none of the elements of the form  $\langle \infty, \beta \rangle$  are reached by horizontal sections of  $X$ . By ( $\star\star$ ), the vertical closure of  $X$  is  $X \cup A \times \{\infty\}$ . Of course, since  $A$  is cofinal in  $\kappa$ , the horizontal closure of  $A \times \{\infty\}$  includes the point  $\langle \infty, \infty \rangle$ .

But then

$$\begin{aligned} \text{vcl}(\text{hcl}(X)) &= X \cup A \times \{\infty\}, \text{ yet} \\ \text{hcl}(\text{vcl}(X)) &= X \cup A \times \{\infty\} \cup \{\langle \infty, \infty \rangle\}. \end{aligned}$$

But this means that

$$\mathbf{card}_\kappa, \mathbf{card}_\lambda \not\models \diamond_1 \diamond_2 p \rightarrow \diamond_2 \diamond_1 p.$$

“(ii) $\Rightarrow$ (i)”. Assume that  $\lambda < \text{cf } \kappa$ . We have to show that  $\mathbf{card}_\kappa, \mathbf{card}_\lambda \models \diamond_1 \diamond_2 p \rightarrow \diamond_2 \diamond_1 p$ , so we have to show for every subset  $X$  of the product that  $\text{hcl}(\text{vcl}(X)) \subseteq \text{vcl}(\text{hcl}(X))$ . Note that the only point for which the order of horizontal and vertical closures matters is the point  $\langle \infty, \infty \rangle$ , so the we are done if we can show that

$$\langle \infty, \infty \rangle \in \text{hcl}(\text{vcl}(X)) \text{ implies } \langle \infty, \infty \rangle \in \text{vcl}(\text{hcl}(X)).$$

Without loss of generality,  $X \subseteq \kappa \times \lambda$ .

If  $\langle \infty, \infty \rangle \in \text{hcl}(\text{vcl}(X))$ , there is a cofinal set  $C \subseteq \kappa$  of cardinality  $\text{cf } \kappa$  such that for all  $\gamma \in C$ , we have

$$\langle \gamma, \infty \rangle \in \text{vcl}(X).$$

This in turn means that for each such  $\gamma$ , the vertical section  $X^\gamma = \{\beta; \langle \gamma, \beta \rangle \in X\}$  must be cofinal in  $\lambda$ . In other words, if you fix  $\eta \in \lambda$ , then

$$X_{>\eta}^* := \{\langle \alpha, \beta \rangle \in X; \beta > \eta\} \cap (\kappa \times C)$$

must have cardinality at least  $\text{cf } \kappa$ .

For each  $\beta \in \lambda$ , let

$$P_\beta := \{\langle \alpha, \beta \rangle \in X; \alpha \in C\}.$$

The family  $\{P_\beta; \eta < \beta < \lambda\}$  is a partition of  $X_{>\eta}^*$  into at most  $\lambda$  many pieces. Consequently, by the pigeon hole principle, there must be a  $\beta^* > \eta$  such that  $P_{\beta^*}$  has  $\text{cf } \kappa$  many elements. But since  $P_{\beta^*} \subseteq X_{\beta^*}$  and  $C$  was cofinal in  $\kappa$ , this means that  $\langle \infty, \beta^* \rangle \in \text{hcl}(X)$ .

Since  $\eta$  was arbitrary, we just showed that the set of such  $\beta^*$  is cofinal in  $\lambda$ , and thus  $\langle \infty, \infty \rangle \in \text{vcl}(\text{hcl}(X))$ . This was the claim. q.e.d.

**Corollary 5** For  $\aleph_0 \leq \lambda < \text{cf } \kappa$ ,  $\mathbf{card}_\kappa$  and  $\mathbf{card}_\lambda$  are non-Alexandroff spaces such that  $\text{com}_{\leftarrow}$  holds in  $\mathbf{card}_\kappa \times \mathbf{card}_\lambda$ . In particular, this is true in  $\mathbf{card}_{\aleph_1} \times \mathbf{card}_{\aleph_0}$ . Also,  $\text{com}_{\rightarrow}$  holds in  $\mathbf{card}_{\aleph_0} \times \mathbf{card}_{\aleph_1}$ .

**Theorem 6** Let  $\kappa$  and  $\lambda$  be cardinals. Then the following are equivalent:

- (i)  $\mathbf{card}_\kappa, \mathbf{card}_\lambda \models \diamond_1 \square_2 p \rightarrow \square_2 \diamond_1 p$ , and
- (ii)  $\text{cf } \lambda \neq \text{cf } \kappa$ .

**Proof.** “(i) $\Rightarrow$ (ii)”. Suppose that  $\vartheta := \text{cf } \kappa = \text{cf } \lambda$ . Let  $A = \{\alpha_\gamma; \gamma < \vartheta\} \subseteq \kappa$  and  $B = \{\beta_\gamma; \gamma < \vartheta\} \subseteq \lambda$  be increasing enumerations of cofinal subsets. Define

$$X := \{\langle \alpha_\gamma, \beta \rangle; \beta \geq \beta_\gamma, \gamma < \vartheta\} \cup \{\langle \alpha_\gamma, \infty \rangle; \gamma < \vartheta\}.$$

Then for each  $\gamma < \vartheta$ ,  $\{\infty\} \cup \{\beta; \beta_\gamma \leq \beta < \lambda\} \subseteq X^{\alpha_\gamma}$  which is an open neighbourhood of  $\infty$  in  $\mathbf{card}_\lambda$ . Consequently,  $\langle \alpha_\gamma, \infty \rangle \in \text{vint}(X)$ . Since  $A$  was cofinal in  $\kappa$ , this means that  $\langle \infty, \infty \rangle \in \text{hcl}(\text{vint}(X))$ .

Yet, for each  $\beta \in \lambda$  there is an upper bound for  $X_\beta$ : if  $\beta_\gamma \leq \beta < \beta_{\gamma+1}$ , then

$$X_\beta \subseteq \{\alpha \in \kappa; 0 \leq \alpha < \alpha_{\gamma+1}\}.$$

That means that  $\text{hcl}(X)$  doesn't contain any element of the form  $\langle \infty, \beta \rangle$ , and so  $\langle \infty, \infty \rangle \notin \text{vint}(\text{hcl}(X))$ .

“(ii) $\Rightarrow$ (i)”. The symmetry of  $\text{chr}$  makes sure that we only have to check the case  $\text{cf } \kappa < \text{cf } \lambda$ .

To start, let us notice that for subsets  $X$  of  $\mathbf{card}_\kappa \times \mathbf{card}_\lambda$ , we always have that

$$\text{hcl}(\text{vint}(X)) \setminus \{\langle \infty, \infty \rangle\} \subseteq \text{vint}(\text{hcl}(X)),$$

since the elements of  $\kappa \times \lambda$  are not affected by any of the interior and closure operations. Thus, we only have to show

$$\langle \infty, \infty \rangle \in \text{hcl}(\text{vint}(X)) \text{ implies } \langle \infty, \infty \rangle \in \text{vint}(\text{hcl}(X)).$$

Fix  $X$  such that  $\langle \infty, \infty \rangle \in \text{hcl}(\text{vint}(X))$ . This means that there is some cofinal set  $A = \{\alpha_\gamma; \gamma < \text{cf } \kappa\} \subseteq \kappa$  such that  $A \times \{\infty\} \subseteq \text{vint}(X)$ , so for each  $\gamma$ , there is some  $\beta_\gamma < \lambda$  such that

$$\{\infty\} \cup \{\beta; \beta_\gamma \leq \beta < \lambda\} \subseteq X^{\alpha_\gamma}.$$

The set  $\{\beta_\gamma; \gamma < \text{cf } \kappa\}$  has cardinality  $\text{cf } \kappa < \text{cf } \lambda$ , so  $\beta^* := \sup\{\beta_\gamma; \gamma < \text{cf } \kappa\} < \lambda$ . But then for every  $\beta > \beta^*$ , we have that  $A \subseteq X_\beta$ , and so  $\langle \infty, \beta \rangle \in \text{hcl}(X)$ . This means that  $\{\infty\} \cup \{\beta; \beta^* < \beta\}$  is an  $\mathbf{card}_\lambda$ -open neighbourhood contained in  $(\text{hcl}(X))^\infty$ , so  $\langle \infty, \infty \rangle \in \text{vint}(\text{hcl}(X))$ . q.e.d.

**Corollary 7** For  $\aleph_0 \leq \text{cf } \kappa < \text{cf } \lambda$ ,  $\mathbf{card}_\kappa$  and  $\mathbf{card}_\lambda$  are non-Alexandroff spaces such that  $\text{chr}$  holds in  $\mathbf{card}_\kappa \times \mathbf{card}_\lambda$ . In particular, this is true in  $\mathbf{card}_{\aleph_0} \times \mathbf{card}_{\aleph_1}$ .

Corollaries 5 and 7 together answer Question 2 negatively:

$$\mathbf{card}_{\aleph_0}, \mathbf{card}_{\aleph_1} \models \text{chr} \ \& \ \text{com}_\rightarrow \ \& \ \neg \text{com}_\leftarrow, \text{ and}$$

$$\mathbf{card}_{\aleph_1}, \mathbf{card}_{\aleph_0} \models \text{chr} \ \& \ \text{com}_\leftarrow \ \& \ \neg \text{com}_\rightarrow.$$

## References

- [1] Johan **van Benthem**, Guram **Bezhanishvili**, Balder **ten Cate**, Darko **Sarenac**, Modal Logics for Products of Topologies, *in preparation*
- [2] John Charles Chenoweth **McKinsey**, Alfred **Tarski**, The algebra of topology, **Annals of Mathematics** 45 (1944), p. 141–191