


Institute for Logic, Language and Computation

**ALGORITHMIC PROBLEMS CONCERNING FIRST-ORDER
DEFINABILITY OF MODAL FORMULAS
ON THE CLASS OF ALL FINITE FRAMES**

A.V. Chagrov, L.A. Chagrova

ILLC Prepublication Series
for Mathematical Logic and Foundations ML-93-07



University of Amsterdam

The ILLC Prepublication Series

1990

Logic, Semantics and Philosophy of Language

- LP-90-01 Jaap van der Does A Generalized Quantifier Logic for Naked Infinitives
 LP-90-02 Jeroen Groenendijk, Martin Stokhof Dynamic Montague Grammar
 LP-90-03 Renate Bartsch Concept Formation and Concept Composition
 LP-90-04 Aarne Ranta Intuitionistic Categorical Grammar
 LP-90-05 Patrick Blackburn Nominal Tense Logic
 LP-90-06 Gennaro Chierchia The Variability of Impersonal Subjects
 LP-90-07 Gennaro Chierchia Anaphora and Dynamic Logic
 LP-90-08 Herman Hendriks Flexible Montague Grammar
 LP-90-09 Paul Dekker The Scope of Negation in Discourse, towards a Flexible Dynamic Montague grammar
 LP-90-10 Theo M.V. Janssen Models for Discourse Markers
 LP-90-11 Johan van Benthem General Dynamics
 LP-90-12 Serge Lapierre A Functional Partial Semantics for Intensional Logic
 LP-90-13 Zhisheng Huang Logics for Belief Dependence
 LP-90-14 Jeroen Groenendijk, Martin Stokhof Two Theories of Dynamic Semantics
 LP-90-15 Maarten de Rijke The Modal Logic of Inequality
 LP-90-16 Zhisheng Huang, Karen Kwast Awareness, Negation and Logical Omniscience
 LP-90-17 Paul Dekker Existential Disclosure, Implicit Arguments in Dynamic Semantics

Mathematical Logic and Foundations

- ML-90-01 Harold Schellinx Isomorphisms and Non-Isomorphisms of Graph Models
 ML-90-02 Jaap van Oosten A Semantical Proof of De Jongh's Theorem
 ML-90-03 Yde Venema Relational Games
 ML-90-04 Maarten de Rijke Unary Interpretability Logic
 ML-90-05 Domenico Zambella Sequences with Simple Initial Segments
 ML-90-06 Jaap van Oosten Extension of Lifschitz' Realizability to Higher Order Arithmetic, and a Solution to a Problem of F. Richman
 ML-90-07 Maarten de Rijke A Note on the Interpretability Logic of Finitely Axiomatized Theories
 ML-90-08 Harold Schellinx Some Syntactical Observations on Linear Logic
 ML-90-09 Dick de Jongh, Duccio Pianigiani Solution of a Problem of David Guaspari
 ML-90-10 Michiel van Lambalgen Randomness in Set Theory
 ML-90-11 Paul C. Gilmore The Consistency of an Extended NaDSet


Computation and Complexity Theory

- CT-90-01 John Tromp, Peter van Emde Boas Associative Storage Modification Machines
 CT-90-02 Sieger van Denneheuvel, Gerard R. Renardel de Lavalette A Normal Form for PCSJ Expressions
 CT-90-03 Ricard Gavaldà, Leen Torenvliet, Osamu Watanabe, José L. Balcázar Generalized Kolmogorov Complexity in Relativized Separations
 CT-90-04 Harry Buhman, Edith Spaan, Leen Torenvliet Bounded Reductions
 CT-90-05 Sieger van Denneheuvel, Karen Kwast Efficient Normalization of Database and Constraint Expressions
 CT-90-06 Michiel Smid, Peter van Emde Boas Dynamic Data Structures on Multiple Storage Media, a Tutorial
 CT-90-07 Kees Doets Greatest Fixed Points of Logic Programs
 CT-90-08 Fred de Geus, Ernest Rotterdam, Sieger van Denneheuvel, Peter van Emde Boas Physiological Modelling using RL
 CT-90-09 Roel de Vrijer Unique Normal Forms for Combinatory Logic with Parallel Conditional, a case study in conditional rewriting
 X-90-01 A.S. Troelstra Remarks on Intuitionism and the Philosophy of Mathematics, Revised Version
 X-90-02 Maarten de Rijke Some Chapters on Interpretability Logic
 X-90-03 L.D. Beklemishev On the Complexity of Arithmetical Interpretations of Modal Formulae
 X-90-04 Annual Report 1989
 X-90-05 Valentin Shehtman Derived Sets in Euclidean Spaces and Modal Logic
 X-90-06 Valentin Goranko, Solomon Passy Using the Universal Modality: Gains and Questions
 X-90-07 V.Yu. Shavrukov The Lindenbaum Fixed Point Algebra is Undecidable
 X-90-08 L.D. Beklemishev Provability Logics for Natural Turing Progressions of Arithmetical Theories
 X-90-09 V.Yu. Shavrukov On Rosser's Provability Predicate
 X-90-10 Sieger van Denneheuvel, Peter van Emde Boas An Overview of the Rule Language RL/1
 X-90-11 Alessandra Carbone Provable Fixed points in $\text{ID}_0 + \Omega_1$, revised version
 X-90-12 Maarten de Rijke Bi-Unary Interpretability Logic
 X-90-13 K.N. Ignatiev Dzhaparidze's Polymodal Logic: Arithmetical Completeness, Fixed Point Property, Craig's Property
 X-90-14 L.A. Chagrova Undecidable Problems in Correspondence Theory
 X-90-15 A.S. Troelstra Lectures on Linear Logic

1991

Logic, Semantics and Philosophy of Language

- LP-91-01 Wiebe van der Hoek, Maarten de Rijke Generalized Quantifiers and Modal Logic
 LP-91-02 Frank Veltman Defaults in Update Semantics
 LP-91-03 Willem Groeneveld Dynamic Semantics and Circular Propositions
 LP-91-04 Makoto Kanazawa The Lambek Calculus enriched with Additional Connectives
 LP-91-05 Zhisheng Huang, Peter van Emde Boas The Schoenmakers Paradox: Its Solution in a Belief Dependence Framework
 LP-91-06 Zhisheng Huang, Peter van Emde Boas Belief Dependence, Revision and Persistence
 LP-91-07 Henk Verkuyl, Jaap van der Does The Semantics of Plural Noun Phrases
 LP-91-08 Víctor Sánchez Valencia Categorical Grammar and Natural Reasoning
 LP-91-09 Arthur Nieuwendijk Semantics and Comparative Logic
 LP-91-10 Johan van Benthem Logic and the Flow of Information
 ML-91-01 Yde Venema Cylindric Modal Logic
 ML-91-02 Alessandro Berarducci, Rineke Verbrugge On the Metamathematics of Weak Theories
 ML-91-03 Domenico Zambella On the Proofs of Arithmetical Completeness for Interpretability Logic
 ML-91-04 Raymond Hoofman, Harold Schellinx Collapsing Graph Models by Preorders
 ML-91-05 A.S. Troelstra History of Constructivism in the Twentieth Century
 ML-91-06 Inge Bethke Finite Type Structures within Combinatory Algebras
 ML-91-07 Yde Venema Modal Derivation Rules
 ML-91-08 Inge Bethke Going Stable in Graph Models
 ML-91-09 V.Yu. Shavrukov A Note on the Diagonalizable Algebras of PA and ZF
 ML-91-10 Maarten de Rijke, Yde Venema Sahlqvist's Theorem for Boolean Algebras with Operators
 ML-91-11 Rineke Verbrugge Feasible Interpretability
 ML-91-12 Johan van Benthem Modal Frame Classes, revisited
 CT-91-01 Ming Li, Paul M.B. Vitányi Kolmogorov Complexity Arguments in Combinatorics
 CT-91-02 Ming Li, John Tromp, Paul M.B. Vitányi How to Share Concurrent Wait-Free Variables
 CT-91-03 Ming Li, Paul M.B. Vitányi Average Case Complexity under the Universal Distribution Equals Worst Case Complexity
 CT-91-04 Sieger van Denneheuvel, Karen Kwast Weak Equivalence
 CT-91-05 Sieger van Denneheuvel, Karen Kwast Weak Equivalence for Constraint Sets
 CT-91-06 Edith Spaan Census Techniques on Relativized Space Classes
 CT-91-07 Karen L. Kwast The Incomplete Database
 CT-91-08 Kees Doets Levationis Laus
 CT-91-09 Ming Li, Paul M.B. Vitányi Combinatorial Properties of Finite Sequences with high Kolmogorov Complexity



Institute for Logic, Language and Computation

Plantage Muidersgracht 24

1018TV Amsterdam

Telephone 020-525.6051, Fax: 020-525.5101

**ALGORITHMIC PROBLEMS CONCERNING FIRST-
ORDER DEFINABILITY OF MODAL FORMULAS ON
THE CLASS OF ALL FINITE FRAMES**

A.V. Chagrov, L.A. Chagrova

Tver State University

Tver, Russia

ALGORITHMIC PROBLEMS CONCERNING FIRST-ORDER DEFINABILITY OF MODAL FORMULAS ON THE CLASS OF ALL FINITE FRAMES

CHAGROV A.V., CHAGROVA L.A.

TVER STATE UNIVERSITY
ZHELYABOV STR., 33
TVER 170013, RUSSIA

• Introduction

The problem of describing the "behaviour" of modal, or other propositional intensional formulas on the class of finite frames seems rather natural. Almost all modal logics resulting from formalization of substantial ideas have turned out to have the finite model property. Moreover, different semantic constructions in the finite case are effectively intertranslatable (there is no incompleteness effect). In this paper we consider the question, for any modal formula, *how to recognize whether it has a first-order equivalent on the class of finite frames*. Our main result is that no effective algorithmic answer to this question is possible.

Earlier, in [5], algorithmic undecidability was proved for first-order definability without restrictions on frame cardinality. The proof there uses essentially undecidable calculi and, so, it cannot be transferred directly to finite frames. But another construction, used in [4] to show the undecidability of finite frame consequence between modal formulas (a research problem first stated in [2]) turned out to be a suitable replacement. Thereby the proof increases considerably in size, since we have to manipulate first-order definability of rather large formulas (cf. the forthcoming paper [6] concerning superintuitionistic logics). One indirect purpose of the present work is to give a simplest possible variant for modal formulas. For this purpose it turned out convenient to use finite GL-frames, having a transitive and irreflexive alternative relation. Note that, in the infinite case, it is almost pointless to use this frame class, as it is not first-order definable.

This paper is organized as follows. In Section 1, the relevant result from [4] about the undecidability of semantic consequence for finite GL-frames is proved, as well as some generalizations thereof. The main technical lemma of this Section, which shows how to reduce the Halting Problem for Minsky Machines to the finite frame consequence problem, will be used in Section 4 to obtain the main results of the present paper. In Section 2, some technical facts are proved about first-order definability of modal formulas. This contains the messy complications we mentioned. Readers interested only in the general idea of our main results may omit the details of constructing the relevant first-order equivalents. In Section 3, formulas closely connected with the constructions of Sections 1 and 2 are presented, which have no first-order equivalents on the class of finite frames. The results of Sections 1–3 are combined in Section 4 to obtain short proofs of the undecidability of various problems connected with first-order definability on the class of finite frames. In the course of this exposition, two notions of first-order definability will be considered, a "local" one and a "global" one. In particular, in Section 3 the problem raised in [1] concerning the difference between global and local definability on the class of all transitive frames is decided.

0 Some Definitions

The usual definitions and results from Modal Logic are employed, including its connections with standard logic. In particular, any Kripke frame can be considered both as a semantic structure for modal formulas and as a model for the first-order language with equality of a single binary predicate. We say that a modal formula ϕ and a first-order sentence ψ are *globally equivalent* on a frame class \mathbf{K} if, for any frame F in \mathbf{K} , $F \models \phi$ iff $F \models \psi$. A

modal formula ϕ is *globally first-order definable* on the class \mathbf{K} if there exists a first-order sentence globally equivalent to it on \mathbf{K} . A modal formula ϕ and a first-order formula ψ with one free variable are *locally equivalent* on \mathbf{K} if, for any frame F in \mathbf{K} and any world d , $\langle F, d \rangle \models \phi$ iff $F \models \psi[d]$. A modal formula ϕ is *locally first-order definable* on the frame class class \mathbf{K} iff there is a first-order formula with one free variable which is locally equivalent to it on \mathbf{K} . It is clear that local first-order definability implies global first-order definability on any frame class. In this paper, we are mainly interested in the class \mathbf{K} of finite frames (but our results remain true for finite transitive irreflexive ones). Henceforth, all frames considered will be finite. If we want to allow that a class contains infinite frames, then we will mention this specifically. Here is some useful notation:

$$\begin{aligned} \phi \models_{\text{fin}} \psi & : \text{for any frame } F, F \models \phi \text{ implies } F \models \psi, \text{ and} \\ \phi \models_{\text{fin}}^{\text{loc}} \psi & : \text{for any } F \text{ and world } d, \langle F, d \rangle \models \phi \text{ implies } \langle F, d \rangle \models \psi. \end{aligned}$$

The relations \models_{fin} and $\models_{\text{fin}}^{\text{loc}}$ are different. An example is provided by:

$$\text{LF} \wedge \diamond \top \models_{\text{fin}} \perp \quad \text{but } \text{LF} \wedge \diamond \top \not\models_{\text{fin}}^{\text{loc}} \perp ,$$

where $\text{LF} = \Box(\Box p \rightarrow p) \rightarrow \Box p$ is the formula that axiomatizes Löb's Logic GL. This principle is non-first-order in general, but "Transitivity and Irreflexivity" is a first-order equivalent (both global and local) for it over the finite frames. We will often "translate" first-order formulas into English: e.g., the preceding term "Irreflexivity" is our translation of the formula $\forall x \neg xRx$. Finally, we will freely use various modal and first-order abbreviations: such as $\Box\phi$ for $\Box\phi \& \phi$, $x=y=z$ for $x=y \wedge y=z$, $xRyRz$ for $xRy \wedge yRz$, xR^3y for $\exists u \exists v (xRuRvRy)$. Other useful notations will be explained in due course.

1 Reduction of Minsky Machine Computability to Modal Consequence on Finite Frames

A 'Minsky Machine' is a two-tape effective machine operating on two integers s_1 and s_2 . A *Minsky Machine Program* is a finite set of instructions I of the forms:

- (1) $q_\alpha \rightarrow q_\beta T_1 T_0$: in state q_α , add 1 to s_1 , and go to q_β ;
- (2) $q_\alpha \rightarrow q_\beta T_0 T_1$: in state q_α , add 1 to s_2 , and go to q_β ;
- (3) $q_\alpha \rightarrow q_\beta T_{-1} T_0 (q_\gamma T_0 T_0)$: in state q_α , subtract 1 from s_1 if $s_1 \neq 0$, and go to q_β , otherwise go to q_γ ;
- (4) $q_\alpha \rightarrow q_\beta T_0 T_{-1} (q_\gamma T_0 T_0)$: in state q_α , subtract 1 from s_2 if $s_2 \neq 0$, and go to q_β , otherwise go to q_γ .

A *Minsky Machine Configuration* is an ordered triple (i, j, k) of natural numbers, where i is a state number, $j = s_1$, and $k = s_2$. We write $P : (\alpha, m, n) \rightarrow (\beta, k, l)$ to express that the program P starting at configuration (α, m, n) can reach configuration (β, k, l) . Henceforth, we fix the symbols q_α and q_β for the initial and final states of the machine. Finally, we introduce two conventions which do not influence standard facts about Minsky machines, but are useful for our constructions: (i) all machines considered are deterministic, i.e., they do not contain different instructions with the same left parts, (ii) "blocking" states do not occur in our machines, i.e., if some nonfinal state is in the right part of some machine instruction, then there is an instruction with that state in the left part.

In this chapter we use two variants of undecidable halting problems of Minsky machines:

- There is a Minsky Machine P such that no algorithm recognizes for any configuration whether P eventually halts (in a final state), starting from that configuration.
- There is a configuration (α, m, n) such that no algorithm recognizes for any Minsky machine program whether it eventually halts, starting from (α, m, n) .

Next, we introduce some important sets of formulas. Although we do not need their semantic "sense" right now, it is useful to have a picture in mind, in accordance with which they are defined. The irreflexive transitive frame of Figure 1 is such picture; where the part of the frame surrounded by the dotted line is not needed in this Section. Here is some




Figure 1

Next, we introduce some important sets of formulas. Although we do not need their semantic "sense" right now, it is useful to have a picture in mind, in accordance with which they are defined. The irreflexive transitive frame of Figure 1 is such picture; where the part of the frame surrounded by the dotted line is not needed in this Section. Here is some explanation of its key features. The frame is constructed from a machine program P and an initial configuration (α, m, n) . The presence of the worlds $h(s_i)$, where $s_i = (\alpha_i, m_i, n_i)$, expresses the fact that s_i is the configuration after the first $i-1$ steps of a computation starting from (α, m, n) . Now set:

$$F = \Box^2 \perp \rightarrow \Box p \vee \Box \neg p, \quad F' = \Box F.$$

These formulas are falsifiable in this frame by the same valuation, F at world f , and F' at f' (and only there). For as soon as F is falsifiable, say by a valuation V , then

$$V(p, f_0^1) = 1, \quad V(p, f_1^1) = 0 \quad \text{or} \quad V(p, f_0^1) = 0, \quad V(p, f_1^1) = 1.$$

We continue to introduce modal formulas, emphasizing their connection with the worlds in which they are true:

- $F_1^1 = \Box \perp \wedge p, \quad F_1^1 = \Box \perp \wedge \neg p,$
 $F_1^j = \Box j \perp \wedge \Box^{j-1} F_1^1 \wedge \Box^{j-1} F_{i-1}^1 \quad (i \in \{0, 1\}, 2 \leq j \leq 7)$
- $A_0^1 = \Diamond F_0^{i+3} \wedge \Diamond F_1^{i+3} \wedge \Box^{i+4} \perp \quad (0 \leq i \leq 3),$
 $A_j^i = \Box j A_0^j \wedge \neg \Box^{j+1} A_0^i \wedge \bigwedge_{k=1}^4 \neg \Box A_0^k \quad (0 \leq i \leq 3, j > 0),$
- $S(\delta, A_k^1, A_l^2) = \bigwedge_{i=0}^{\delta} \Diamond A_i^1 \wedge \neg \Diamond A_{\delta}^0 \wedge \Diamond A_k^1 \wedge \neg \Diamond \Diamond A_k^1 \wedge \Diamond A_l^2 \wedge \neg \Diamond \Diamond A_l^2$
 (where $\delta, k, l \geq 0$).

The formula $S(\delta, A_k^1, A_l^2)$ corresponds to the configuration (δ, k, l) . But for describing instructions we need formulas speaking about arbitrary configurations. Therefore, we set:

$$\begin{array}{ll} Q_1 & (\Diamond A_0^1 \vee A_0^1) \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^2 \wedge \neg \Diamond A_0^3 \wedge p_1 \wedge \neg \Diamond p_1, \\ Q_2 & \Diamond A_0^1 \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^2 \wedge \neg \Diamond A_0^3 \wedge p_1 \wedge \neg \Diamond p_1, \\ R_1 & (\Diamond A_0^2 \vee A_0^2) \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^1 \wedge \neg \Diamond A_0^3 \wedge p_2 \wedge \neg \Diamond p_2, \\ R_2 & \Diamond A_0^2 \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^1 \wedge \neg \Diamond A_0^3 \wedge p_2 \wedge \neg \Diamond p_2, \\ T_1 & \Diamond A_0^3 \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^1 \wedge \neg \Diamond A_0^2 \wedge p_3 \wedge \neg \Diamond p_3, \\ T_2 & \Diamond A_0^3 \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^1 \wedge \neg \Diamond A_0^2 \wedge p_3 \wedge \neg \Diamond p_3, \\ S(\delta, Q_i, R_j) & \bigwedge_{k=0}^{\delta} \Diamond A_k^0 \wedge \neg \Diamond A_{\delta+1}^0 \wedge \Diamond Q_i \wedge \neg \Diamond \Diamond Q_i \wedge \Diamond R_j \wedge \neg \Diamond \Diamond R_j, \\ S(\delta, Q_1, A_0^2) & \bigwedge_{k=0}^{\delta} \Diamond A_k^0 \wedge \neg \Diamond A_{\delta+1}^0 \wedge \Diamond Q_1 \wedge \neg \Diamond \Diamond Q_1 \wedge \Diamond A_0^2 \wedge \neg \Diamond \Diamond A_0^2, \\ S(\delta, A_0^1, R_1) & \bigwedge_{k=0}^{\delta} \Diamond A_k^0 \wedge \neg \Diamond A_{\delta+1}^0 \wedge \Diamond A_0^1 \wedge \neg \Diamond \Diamond A_0^1 \wedge \Diamond R_1 \wedge \neg \Diamond \Diamond R_1 \end{array}$$

(where $\delta \geq 0, i, j \in \{1, 2\}$).

We shall write $\varphi \equiv \psi$ if $\varphi \leftrightarrow \psi$ is true in all finite transitive and irreflexive frames, i.e. $GL \vdash \varphi \leftrightarrow \psi$. $\varphi \equiv \psi$ implies that φ and ψ are always semantically interchangeable in models of GL .

Lemma 1.1 Let φ^* be the result of the substitution in φ of $\diamond^k A_0^1$ for p_1 , $\diamond^L A_0^2$ for p_2 and $\diamond A_0^3$ for p_3 . Then

- i) $Q_1^* \equiv A_k^1, Q_2^* \equiv A_{k+1}^1;$
- ii) $R_1^* \equiv A_1^2, R_2^* \equiv A_{1+1}^2;$
- iii) $T_1^* \equiv A_m^3, T_2^* \equiv A_{m+1}^3;$
- iv) $(S(\delta, Q_i, R_j))^* \equiv S(\delta, A_{k+(i-1)}^1, A_{1+(j-1)}^2) \quad (i, j \in \{1, 2\});$
- v) $(S(\delta, Q_1, A_0^2))^* \equiv S(\delta, A_k^1, A_0^2);$
- vi) $(S(\delta, A_0^1, R_1))^* \equiv S(\delta, A_0^1, A_1^2).$

Proof. Immediate. □

The formulas that simulate instructions use subformulas of the form T_i for the "calculation of some number of steps". For instruction I set:

- if I is of the form $q_\delta \rightarrow q_\delta T_1 T_0$, then
 $AxI = \neg F \wedge \diamond(S(\delta, Q_1, R_1) \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1) \wedge \diamond T_2 \rightarrow \diamond(S(\delta, Q_2, R_1) \wedge \diamond T_2 \wedge \neg \diamond \diamond T_2);$
- if I is of the form $q_\delta \rightarrow q_\delta T_0 T_1$, then
 $AxI = \neg F \wedge \diamond(S(\delta, Q_1, R_1) \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1) \wedge \diamond T_2 \rightarrow \diamond(S(\delta, Q_1, R_2) \wedge \diamond T_2 \wedge \neg \diamond \diamond T_2);$
- if I is of the form $q_\delta \rightarrow q_\delta T_{-1} T_0(q_\epsilon, T_0 T_0)$, then
 $AxI = (\neg F \wedge \diamond(S(\delta, Q_2, R_1) \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1) \wedge \diamond T_2 \rightarrow \diamond(S(\delta, Q_1, R_1) \wedge \diamond T_2 \wedge \neg \diamond \diamond T_2))$
 $\wedge (\neg F \wedge \diamond(S(\delta, A_0^1, R_1) \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1) \wedge \diamond T_2 \rightarrow \diamond(S(\epsilon, A_0^1, R_1) \wedge \diamond T_2 \wedge \neg \diamond \diamond T_2));$
- if I is of the form $q_\delta \rightarrow q_\delta T_0 T_{-1}(q_\epsilon, T_0 T_0)$, then
 $AxI = (\neg F \wedge \diamond(S(\delta, Q_1, R_2) \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1) \wedge \diamond T_2 \rightarrow \diamond(S(\delta, Q_1, R_1) \wedge \diamond T_2 \wedge \neg \diamond \diamond T_2))$
 $\wedge (\neg F \wedge \diamond(S(\delta, Q_1, A_0^2) \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1) \wedge \diamond T_2 \rightarrow \diamond(S(\epsilon, Q_1, A_0^2) \wedge \diamond T_2 \wedge \neg \diamond \diamond T_2)).$

Next, for a Minsky program P set

$$AxP = \bigwedge_{I \in P} AxI,$$

and then define the formula $A(P)$ as follows:

$$A(P) = LF \wedge AxP \wedge LinT \wedge UT,$$

where

$$LinT = \neg(\neg F \wedge \diamond(\diamond A_0^3 \wedge \neg \diamond A_0^0 \wedge \neg \diamond A_0^1 \wedge \neg \diamond A_0^2 \wedge \square r \wedge \neg q) \wedge \diamond(\diamond A_0^3 \wedge \neg \diamond A_0^0 \wedge \neg \diamond A_0^1 \wedge \neg \diamond A_0^2 \wedge \square q \wedge \neg r));$$

$$UT = \neg(\neg F \wedge \diamond(q \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1 \wedge \diamond A_0^0 \wedge \diamond A_0^1 \wedge \diamond A_0^2) \wedge \diamond(\neg q \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1 \wedge \diamond A_0^0 \wedge \diamond A_0^1 \wedge \diamond A_0^2)).$$

Finally, define the formula $B(\alpha, m, n)$ as follows :

$$B(\alpha, m, n) = \neg F \wedge \diamond S(\alpha, A_m^1, A_n^2) \wedge \diamond A_3^1 \wedge \diamond A_3^1 \rightarrow \neg \diamond(S(\beta, Q_1, R_1) \wedge \diamond T_1 \wedge \neg \diamond \diamond T_1).$$

Now, the main technical result of this Section is the following

Lemma 1.2 $A(P) \models_{fin} B(\alpha, m, n)$ iff the program P , starting at configuration (α, m, n) cannot reach a final state (with number β).

Proof. "If". This is the direction for which our specific finite frames were developed. Let the Minsky program P continue indefinitely, starting from the configuration (α, m, n) . We show that then, $A(P) \models_{fin} B(\alpha, m, n)$, by a reductio ad absurdum. Suppose that, for some finite frame F

- (1) $F \models A(P)$,
- (2) $F \not\models B(\alpha, m, n)$.

Condition (2) means that, for some valuation V , in some world a from F , we have $V(B(\alpha, m, n), a) = 0$ (we abbreviate this as: $a, B(\alpha, m, n)$). I.e.,

- (3) $a \not\models F$,
- (4) $a \models \Diamond(S(\alpha, A_m^1, A_n^2) \wedge \Diamond A_1^3 \wedge \neg \Diamond \Diamond A_1^3)$,
- (5) $a \models \Diamond(S(\beta, Q_1, R_1) \wedge \Diamond T_1 \wedge \neg \Diamond \Diamond T_1)$.

Condition (4) implies, that there is a world b in F such that aRb and

- (6) $b \models S(\alpha, A_m^1, A_n^2)$,
- (7) $b \models \Diamond A_1^3$,
- (8) $b \not\models \Diamond \Diamond A_1^3$.

Condition (5) implies, that there is a world c in F such that aRc and

- (9) $c \models \Diamond S(\beta, Q_1, R_1)$,
- (10) $c \models \Diamond T_1$,
- (11) $c \not\models \Diamond \Diamond T_1$.

Condition (7) implies that there is a world (denote it by a_1^3) such that bRa_1^3 and $a_1^3 \models A_1^3$, i.e.,

- (12) $a_1^3 \models \Diamond A_0^3 \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^1 \wedge \neg \Diamond A_0^2$,
- (13) $a_1^3 \not\models \Diamond \Diamond A_0^3$.

Condition (10) implies that there must be a world x such that sRx and $x \models T_1$, i.e.,

- (14) $x \models p_3 \wedge \neg \Diamond p_3$.

Now we need a further auxiliary result.

Lemma 1.3 For any frame F , if $F \models LF \wedge LinT$ and the formula F is falsifiable at one of its worlds h by some valuation, then the set of worlds of this frame which are accessible from h , in which the formula $\Diamond A_0^3 \wedge \neg \Diamond A_0^0 \wedge \neg \Diamond A_0^1 \wedge \neg \Diamond A_0^2$ is true, is strictly linearly ordered by the relation of accessibility.

Proof. This is a standard exercise. □

We continue the proof of Lemma 1.2. By Lemma 1.3, conditions (1), (3) above imply that the set of worlds in F that are accessible from a at which the conditions (12), (14) hold, form a chain strictly linearly ordered by the alternative relation in which a_1^3 is the R-greatest element, because of (13). Denote this chain by

$$a_k^3 R a_{k-1}^3 R a_{k-2}^3 R \dots R a_2^3 R a_1^3.$$

We can characterize its elements as follows by the formulas $A_i^3 : a_i^3 \models A_i^3$ iff $i = j$. Now, (14) implies, that $x = a_s^3$ for some s , $1 \leq s \leq k$, so that we have

$$(15) \quad a_s^3 \models T_1 \quad (\text{from (12), (14)}),$$

$$(16) \quad c \models \Diamond A_s^3,$$

$$(17) \quad c \not\models \Diamond \Diamond A_s^3 \quad (\text{from (11)}).$$

Now that we have succeeded in identifying x with some a_s^3 , and with the help of the conjunctive members of $A(P)$, which are true in F by (1), we shall get "step by step" from a_1^3 to x . This turns out possible exactly thanks to the finiteness of F :

Let P , starting from (α, m, n) , one application of some instruction at a time, give the successive configurations $(\alpha_2, m_2, n_2), (\alpha_3, m_3, n_3), \dots$; where we identify (α, m, n) with (α_1, m_1, n_1) . Note, that by the given conditions, $\alpha_i \neq \beta$ for any $i \in \omega$. In this sequence of configurations, we shall only be interested in finite initial segments of length s :

$$(\alpha_1, m_1, n_1), (\alpha_2, m_2, n_2), \dots, (\alpha_s, m_s, n_s).$$

Note that for each i , $1 \leq i \leq k-1$,

$$a \models \begin{aligned} & \Diamond(S(\alpha_i, A_{m;i}^1, A_{n;i}^2) \wedge \Diamond A_i^3 \wedge \neg \Diamond \Diamond A_i^3) \rightarrow \\ & \Diamond(S(\alpha_{i+1}, A_{m;i+1}^1, A_{n;i+1}^2) \wedge \Diamond A_{i+1}^3 \wedge \neg \Diamond \Diamond A_{i+1}^3). \end{aligned}$$

Indeed, from (1), we have by substitution in a suitable conjunct of AxP :

$$a \models \begin{aligned} & \neg F \wedge \Diamond(S(\alpha_i, A_{m;i}^1, A_{n;i}^2) \wedge \Diamond A_i^3 \wedge \neg \Diamond \Diamond A_i^3) \wedge \Diamond A_{i+1}^3 \rightarrow \\ & \Diamond(S(\alpha_{i+1}, A_{m;i+1}^1, A_{n;i+1}^2) \wedge \Diamond A_{i+1}^3 \wedge \neg \Diamond \Diamond A_{i+1}^3), \end{aligned}$$

and (3) and aRa_{i+1}^3 together with $a_{i+1}^3 \models A_{i+1}^3$ yield that $a \models \neg F \wedge \Diamond A_{i+1}^3$, which gives the desired result. Applying (4) and "Modus Ponens" successively one obtains then, for any i , $1 \leq i \leq k-1$, that

$$a \models \Diamond(S(\alpha_{i+1}, A_{m;i+1}^1, A_{n;i+1}^2) \wedge \Diamond A_{i+1}^3 \wedge \neg \Diamond \Diamond A_{i+1}^3)$$

and hence in particular,

$$a \models \Diamond(S(\alpha_s, A_{m;s}^1, A_{n;s}^2) \wedge \Diamond A_s^3 \wedge \neg \Diamond \Diamond A_s^3).$$

The last condition implies that there is a world d such that aRd and

$$(18) \quad d \models S(\alpha_s, A_{m;s}^1, A_{n;s}^2),$$

$$(19) \quad d \models \Diamond A_s^3,$$

$$(20) \quad d \not\models \Diamond \Diamond A_s^3.$$

From (19), (20), using Lemma 1.3 and the fact that $a_1^3 \models A_1^3$, dRa_s^3 while not dRa_{s+1}^3 , because of which there is a unique world among a_1^3, \dots, a_k^3 in which the formula T_1 holds, it follows with the help of (12), (14) and $x = a_s^3$ that

$$(21) \quad d \models \Diamond T_1 \wedge \neg \Diamond \Diamond T_1.$$

Now, we have collected all necessary conditions. From (9), (10), (11) we obtain

$$(22) \quad c \models \Diamond T_1 \wedge \neg \Diamond \Diamond T_1 \wedge \Diamond A_0^0 \wedge \Diamond A_0^1 \wedge \Diamond A_0^2,$$

and from (19), (21),

$$(23) \quad d \models \Diamond T_1 \wedge \neg \Diamond \Diamond T_1 \wedge \Diamond A_0^0 \wedge \Diamond A_0^1 \wedge \Diamond A_0^2.$$

The worlds c and d are different, because (9) implies that $c \models \bigwedge_{i=0}^{\beta} \Diamond A_i^0 \wedge \neg \Diamond A_{\beta+1}^0$, and from (18) we get $d \models \bigwedge_{i=0}^{\alpha_s} \Diamond A_i^0 \wedge \neg \Diamond A_{\alpha_s+1}^0$, by the condition that $\alpha_s \neq \beta$. This difference allows us to define the valuation V for the propositional variable q in such a way that $c \models q$, $d \not\models q$. Together with (22), (23), and also (3), this gives a $\not\models UT$ which contradicts (1).

This contradiction shows that $A(P) \models_{\text{fin}} B(\alpha, m, n)$: and thus, we have proved the "If" direction of Lemma 1.2.

"Only if". Let us now assume that the program P can reach a final state starting from the configuration (α, m, n) . We will show that $A(P) \not\models_{\text{fin}} B(\alpha, m, n)$, by constructing a suitable finite frame. Let

$$s_1 = (\alpha, m, n) = (\alpha_1, m_1, n_1), s_2 = (\alpha_2, m_2, n_2), \dots, s_k = (\alpha_k, m_k, n_k)$$

be the list of all configurations, successively traversed by program P , starting at (α, m, n) . Set

$$u = \max(\alpha_1, \dots, \alpha_k), v = \max(m_1, \dots, m_k), w = \max(n_1, \dots, n_k).$$

Consider the frame F sketched in Figure 1. We remind the reader that the part of F which is "enclosed by the dotted line" remains disregarded for the present.

Lemma 1.4 $F \not\models A(P)$.

Proof. This is a routine verification. For analogous scrupulous verifications see [5], [3]. \square

Now introduce a valuation on F such that

$$\begin{aligned} x \models p &\iff x = f_0^1; & x \models p_1 &\iff x = a_{m_k}^1; \\ x \models p_2 &\iff x = a_{n_k}^2; & x \models p_3 &\iff x = a_k^3. \end{aligned}$$

Lemma 1.5 This valuation has the effect that $f \not\models B(\alpha, m, n)$.

Proof. This is again a routine verification. \square

From Lemmas 1.4 and 1.5 we obtain the desired result, and Lemma 1.2 is proved. \square

Because $A(P)$ and the formulas $B(\alpha, m, n)$ are in fact constructed effectively from the program P and the configuration (α, m, n) , the undecidability of the earlier two halting problems gives the following two results.

Theorem 1.6 *There is a formula φ such that the problem of recognizing, given any formula ψ , whether $\varphi \models_{fin} \psi$, is algorithmically undecidable.*

Theorem 1.7 *There is a formula ψ such that the problem of recognizing, given any formula φ , whether $\varphi \models_{fin} \psi$, is algorithmically undecidable.*

From these theorems we obtain

Corollary 1.8 *There is no algorithm which recognizes, given any two modal formulas φ and ψ , whether $\varphi \models_{fin} \psi$.*

Taking into account the simple fact that $\{\langle \varphi, \psi \rangle \mid \varphi \not\models_{fin} \psi\}$ is recursively enumerable, we obtain from 1.6 – 1.8 this further

Corollary 1.9 *a) The set $\{\langle \varphi, \psi \rangle \mid \varphi \models_{fin} \psi\}$ is not recursively enumerable.
b) There is a formula φ such that $\{\psi \mid \varphi \models_{fin} \psi\}$ is not recursively enumerable.
c) There is a formula ψ such that $\{\varphi \mid \varphi \models_{fin} \psi\}$ is not recursively enumerable.*

From the proofs of statements 1.6 – 1.9 we obtain the following analogues for \models_{fin}^{loc} .

Theorem 1.10 *There is a formula φ such that the problem of recognizing, given any modal formula ψ , whether $\varphi \models_{fin}^{loc} \psi$, is algorithmically undecidable.*

Theorem 1.11 *There is a formula ψ such that the problem of recognizing, given any modal formula φ , whether $\varphi \models_{fin}^{loc} \psi$, is algorithmically undecidable.*

Corollary 1.12 *There is no algorithm which recognizes, given any two modal formulas φ and ψ , whether $\varphi \models_{fin}^{loc} \psi$.*

Corollary 1.13 *a) The set $\{\langle \varphi, \psi \rangle \mid \varphi \models_{fin}^{loc} \psi\}$ is not recursively enumerable.
b) There is a formula φ such that $\{\psi \mid \varphi \models_{fin}^{loc} \psi\}$ is not recursive enumerable.
c) There is a formula ψ such that $\{\varphi \mid \varphi \models_{fin}^{loc} \psi\}$ is not recursively enumerable.*

Now change the definition of $A(P)$ by adding a conjunct $\Diamond T$. In this case the above proofs will work for \models_{fin}^{loc} , but no longer for \models_{fin} , because after such a change, we will have $A(P) \wedge \Diamond T \models_{fin} B(\alpha, m, n)$, independently of the choice of $P, (\alpha, m, n)$. Thus, one obtains

Theorem 1.14 *a) $\{\langle \varphi, \psi \rangle \mid \varphi \models_{fin} \psi, \varphi \not\models_{fin}^{loc} \psi\}$ is not recursive.
b) There is a formula φ such that $\{\psi \mid \varphi \models_{fin} \psi, \varphi \not\models_{fin}^{loc} \psi\}$ is not recursive.
c) There is a formula ψ such that $\{\varphi \mid \varphi \models_{fin} \psi, \varphi \not\models_{fin}^{loc} \psi\}$ is not recursive.*

We do not know if it is possible to replace "recursive" in the above statements by "recursively enumerable".

2 A First-Order Definable Variant On Finite Frames For Formulas Describing Minsky Machines

To obtain the main result of this paper we need an additional property for the formulas $A(P)$ from the previous chapter – namely their first-order definability on finite frames. Although the above construction may have produced formulas of this kind, we shall complicate the organization of $A(P)$ to simplify the proof of its first-order definability. It will have to be checked, of course, that the new version will still do the job of the original formula $A(P)$. Now we shall merely add conjuncts that are true in the frame of Figure 1, as may be verified immediately, and this is harmless. The new $A(P)$ is defined as follows:

$$A(P) = \quad LF \wedge UF \wedge \text{Last}_2 \wedge UF_0^1 \wedge \dots \wedge F_0^6 \wedge UA_0^0 \wedge UA_0^1 \wedge UA_0^2 \wedge UA_0^3 \\ \wedge \text{Lin}A_0^1 \wedge \text{Lin}A_0^2 \wedge \text{Lin}A_0^3 \wedge UT \wedge AxP,$$

$$\text{where} \quad \text{Last}_2 = \quad \Box(q \wedge r \rightarrow \Diamond T) \vee \Box(q \wedge \neg r \rightarrow \Diamond T) \vee \Box(\neg q \wedge r \rightarrow \Diamond T), \\ UF = \quad \Box(q \rightarrow F) \vee \Box(\neg q \rightarrow F),$$

and for $X \notin \{F, S\}$,

$$UX = \quad (\Box(X \rightarrow \Box(Y_1 \vee \dots \vee Y_m)) \wedge (\Box(q \rightarrow X) \vee \Box(\neg q \rightarrow X))) \vee F,$$

where X, Y_1, \dots, Y_m are formulas corresponding to different worlds x, y_1, \dots, y_m in Figure 1, such that xRy_1, \dots, xRy_m and whenever xRz , then $z \in \{y_1, \dots, y_m\}$,

$$\text{and finally,} \quad \text{Lin}A_0^i = \quad \Box(\Diamond A_0^i \wedge \bigwedge_{j=0}^3 \neg \Diamond A_0^j \wedge \Box r \rightarrow q) \\ \vee \Box(\Diamond A_0^i \wedge \bigwedge_{j=0}^3 \neg \Diamond A_0^j \wedge \Box q \rightarrow r) \vee F.$$

Now we use the new variant of $A(P)$. For this formula, the old Lemma 1.2 still holds. Moreover, we have:

Lemma 2.1 *The new formula $A(P)$ is first-order definable on finite frames (both globally and locally).*

Proof. We shall prove the first-order definability step by step, for which we introduce the following notation. For a formula of the form $A_1 \wedge \dots \wedge A_{j-1} \wedge \dots \wedge A_m$, A_i^* will denote $A_1 \wedge \dots \wedge A_i$. For example, $(UF_0^2)^* = LF \wedge UF \wedge \text{Last}_2 \wedge UF_0^1 \wedge UF_0^2$ and $(AxP)^* = A(P)$ for the new formula $A(P)$. We shall construct a first-order local equivalent for the conjunction of the first i conjuncts of $A(P)$, increasing i till we get to the full $A(P)$. The first-order equivalent on finite frames ("first-order equivalent", for short) for formulas $(A)^*$ will be denoted by $\text{foe}(A)^*$. For instance, we have

$$\text{foe}(LF)^* = \text{foe}(LF) = \text{"transitivity and irreflexivity"}.$$

In all conjuncts of $A(P)$, except the first, the formula F occurs. We describe the conditions of its refutation. The following auxiliary predicate will be useful:

$$l(x) = n \quad \Leftrightarrow \quad \text{"a chain of worlds of length } n \text{ is accessible from } x \\ \text{and chains of length } n+1 \text{ are not accessible"}.$$

Lemma 2.2 *The formula F is refuted at world x of frame F iff there are worlds y, z_1, z_2 in F such that y is accessible from x and $l(y) = 1$, $l(z_1) = l(z_2) = 0$, while yRz_1 , yRz_2 , $z_1 \neq z_2$. This assertion presupposes that the refuting valuation has $z_1 \models p$, $z_2 \not\models p$.*

Proof. This may be verified immediately. \square

For convenience, we denote the right-hand condition in Lemma 2.2 by $CR(x, y, z_1, z_2)$.

Remark. Lemma 2.2 implies that the formulas F and $\forall y \forall z_1 \forall z_2 \neg CR(x, y, z_1, z_2)$ are locally equivalent. \square

Now we set $foe(UF)^* = foe(LF) \wedge \forall x_1 \forall x_2 ((xRx_1 \vee x=x_1) \wedge (xRx_2 \vee x=x_2) \wedge \exists y \exists z_1 \exists z_2 CR(x_2, y, z_1, z_2) \rightarrow x=x_1 \wedge x=x_2)$.

This formula expresses the uniqueness of the world in which F is refuted.

Lemma 2.3 $(UF)^*$ and $foe(UF)^*$ are locally equivalent.

Proof. Let $(F)^*$ be refuted at world a of frame F . We show that $F \not\models foe(UF)^*(a/x)$. $\langle F, a \rangle \not\models (UF)^*$ means that there is a valuation V such that $F, V, a \not\models (UF)^*$ (later on, we shall omit V as usual). Then $a \not\models LF$ or $a \models LF$, but $a \not\models UF$. We may suppose that $\langle F, x \rangle \models foe(LF)$, because otherwise, the desired result is obtained at once. Then we have $a \not\models UF$, i.e.,

$$(1) \quad a \not\models \Box(p \rightarrow F), \quad (2) \quad a \not\models \Box(\neg p \rightarrow F).$$

From (1) we obtain that in F there is a world a_1 such that aRa_1 or $a = a_1$ such that

$$(3) \quad a_1 \models p, \quad (4) \quad a_1 \not\models F.$$

From (2) we obtain that in F there is also a world a_2 such that aRa_2 or $a = a_2$ with

$$(5) \quad a_2 \not\models p, \quad (6) \quad a_2 \not\models F.$$

From (4) and (6), using Lemma 2.2 we obtain that in F there are worlds $b', c_1', c_2', b'', c_1'', c_2''$ such that the formulas $CR(a_1, b', c_1', c_2')$ and $CR(a_2, b'', c_1'', c_2'')$, and hence the formulas $\exists y \exists z_1 \exists z_2 CR(a_1, y, z_1, z_2)$, $\exists y \exists z_1 \exists z_2 CR(a_2, y, z_1, z_2)$ are true. From this, by (3) and (5) it follows, that $a \neq a_1$ or $a \neq a_2$. Collecting all facts obtained so far, one obtains that

$$F \not\models (aRa_1 \vee a=a_1) \wedge (aRa_2 \vee a=a_2) \wedge \exists y \exists z_1 \exists z_2 CR(a_1, y, z_1, z_2) \wedge \exists y \exists z_1 \exists z_2 CR(a_2, y, z_1, z_2) \rightarrow a=a_1 \wedge a=a_2,$$

which refutes the second conjunct of $foe(UF)^*(a/x)$, i.e. $F \not\models foe(UF)^*(a/x)$.

Next, let, for some world a of some frame F , $F \not\models foe(UF)^*(a/x)$. We show that $(UF)^*$ is refuted at a by some valuation on F . First, we may suppose that $F \models foe(LF)(a/x)$, because otherwise, we can refute LF at the world a . Now the cone of F generated by the world a is transitive and irreflexive, and there are a_1, a_2 in F such that aRa_1 or $a = a_1$ and aRa_2 or $a = a_2$, and $a \neq a_1$ or $a \neq a_2$, and moreover, there are $b', c_1', b'', c_1'', c_2''$ such that in F the formulas $CR(a_1, b', c_1', c_2')$, $CR(a_2, b'', c_1'', c_2'')$ hold. Because of $l(b') = l(b'') = 1$, we have

$$(7) \quad b' \models \Box^2 \perp, \quad (8) \quad b'' \models \Box^2 \perp.$$

Two cases arise here: $\{c_1', c_2'\} \cap \{c_1'', c_2''\} = \emptyset$ and $\{c_1', c_2'\} \cap \{c_1'', c_2''\} \neq \emptyset$. In the former case, we can choose a valuation such that:

$$c_1' \models p, c_2' \not\models p, c_1'' \models p, c_2'' \not\models p.$$

In the latter case, say $c_1' = c_2''$, we can choose a valuation such that:

$$c_1' \models p, c_2' \not\models p, c_1'' \not\models p.$$

Thus, in both cases we have

$$(9) \quad b' \not\models \Box p \vee \Box \neg p, \quad (10) \quad b'' \not\models \Box p \vee \Box \neg p.$$

From (7), (9) and (8), (10) we obtain $b' \not\models F$, $b'' \not\models F$, and hence $a \not\models F$, $a_2 \not\models F$. Recall that at least two worlds from a, a_1, a_2 are different, say $a \neq a_2$. Then we can suppose that

$$a \models q, a_2 \not\models q,$$

whence $a \not\models \Box(q \rightarrow F) \vee \Box(\neg q \rightarrow F)$, and so $a \not\models (UF)^*$. Thus, Lemma 2.3 is proved. \square

Now, we define $\text{foe}(\text{Last}_2)$:

$$\text{foe}(\text{Last}_2) = \forall y \forall z \forall u (xRy \wedge xRz \wedge xRu \wedge y \neq u \wedge y \neq z \wedge u \neq z \rightarrow l(y) \neq 0 \vee l(z) \neq 0 \vee l(u) \neq 0).$$

Lemma 2.4 *The formulas Last_2 and $\text{foe}(\text{Last}_2)$ are locally equivalent.*

Proof. This is a standard exercise. \square

This statement allows us to define $\text{foe}(\text{Last}_2)^*$ as follows:

$$\text{foe}(\text{Last}_2)^* = \text{foe}(UF)^* \wedge \text{foe}(\text{Last}_2),$$

Thus, as a consequence of Lemmas 2.3. and 2.4 we have the following statement:

Lemma 2.5 *The formulas $(\text{Last}_2)^*$ and $\text{foe}(\text{Last}_2)^*$ are locally equivalent.*

Now we define $\text{foe}(UF_0^2)^*$:

$$\text{foe}(UF_0^2)^* = \text{foe}(\text{Last}_2)^* \wedge \forall y \forall z_1 \forall z_2 \forall u_1 \forall u_2 (CR(x, y, z_1, z_2) \wedge xRu_1 \wedge xRu_2 \wedge \neg u_1Rz_2 \wedge \neg u_2Rz_2) \vee (u_1Rz_2 \wedge u_2Rz_2 \wedge \neg u_1Rz_1 \wedge \neg u_2Rz_1) \wedge l(u_1) = l(u_2) = 1 \rightarrow u_1 = u_2).$$

Applied to the frame of Figure 1 the conjunct added to $\text{foe}(\text{Last}_2)^*$ means that the world from which the world f_0^1 (or f_1^1) is accessible exactly by one step, but f_1^1 (respectively, f_0^1) is not accessible, is unique – in this particular frame, it is f_0^2 (respectively, f_1^2). The conjuncts to be added later to UF_0^3, \dots, UF_0^6 have an analogous sense.

Lemma 2.6 *The formulas $(UF_0^2)^*$ and $\text{foe}(UF_0^2)^*$ are locally equivalent.*

Our next step is the following definition, for all $i, 3 \leq i \leq 6$:

$$\text{foe}(UF_0^i)^* = \text{foe}(UF_0^{i-1})^* \wedge \forall y \forall z_1 \forall z_2 \forall u_1 \forall u_2 (CR(x, y, z_1, z_2) \wedge ((u_1R^{i-1}z_1 \wedge u_2R^{i-1}z_1 \wedge \neg u_1Rz_2 \wedge \neg u_2Rz_2) \vee (u_1R^{i-1}z_2 \wedge u_2R^{i-1}z_2 \wedge \neg u_1Rz_1 \wedge \neg u_2Rz_1)) \wedge l(u_1) = l(u_2) = i-1 \rightarrow u_1 = u_2).$$

Lemma 2.7 *The formulas $(UF_0^i)^*$ and $\text{foe}(UF_0^i)^*$ are locally equivalent for $3 \leq i \leq 6$.*

Proof. By a stepwise argument for each successive i ($3 \leq i \leq 6$). \square

Let now $\text{foe}(UA_0^0)$ be the negation of a formula expressing the following situation: worlds are accessible from world x which form a subframe generated by worlds y, u_1, u_2 . This is sketched in Figure 2. Set

$$\text{foe}(UA_0^0)^* = \text{foe}(UF_0^6)^* \wedge \text{foe}(UA_0^0).$$

The formulas $\text{foe}(UA_0^1)$, $\text{foe}(UA_0^2)$, $\text{foe}(UA_0^3)$ are defined analogously, using the frames in Figures 3, 4, 5, respectively, instead of Figure 2. Now define

$$\text{foe}(UA_0^i)^* = \text{foe}(UA_0^{i-1})^* \wedge \text{foe}(UA_0^i), \quad 1 \leq i \leq 3.$$

Lemma 2.8 *The formulas $(UA_0^i)^*$ and $\text{foe}(UA_0^i)^*$ are locally equivalent, for $0 \leq i \leq 3$.*

Proof. Again, this is constructed successively for each i , $0 \leq i \leq 3$. \square

Now, if $\text{foe}(UA_0^3)$ is true in some world x of some frame, then the generated subframes sketched in Figures 2–5 cannot be accessible from this world, whereas the generated subframes sketched in Figures 6–9 can be. We shall say that, if one of these generalized subframes is accessible from world x in a frame, then world v_i (see Figures 6–9) is the 'i-th marked world'. Note that the formulas $(UA_0^i)^*$ and $\text{foe}(UA_0^i)^*$ state, in particular, that in a frame at most one i-th marked world is accessible from some world. For later convenience, we introduce the following predicates $\text{ch}_i(u)$:

- $\text{ch}_i(u)$: the i-th marked world is accessible from u or is equal to u ,
and the other marked worlds are not accessible from u , $0 \leq i \leq 2$;
- $\text{ch}_3(u)$: the third marked world is accessible from u ,
while other marked worlds are not accessible.

Let $\text{foe}(\text{Lin} \diamond A_0^i)$ express the following: a set of worlds accessible from x and having the property ch_i is linearly ordered. Set

$$\begin{aligned} \text{foe}(\text{Lin} \diamond A_0^0)^* &= \text{foe}(UA_0^3)^* \wedge \text{foe}(\text{Lin} \diamond A_0^0), \\ \text{foe}(\text{Lin} \diamond A_0^i)^* &= \text{foe}(\text{Lin} \diamond A_0^{i-1})^* \wedge \text{foe}(\text{Lin} \diamond A_0^i), \quad 1 \leq i \leq 3. \end{aligned}$$

Lemma 2.9 *The formulas $(\text{Lin} \diamond A_0^i)^*$ and $\text{foe}(\text{Lin} \diamond A_0^i)^*$ are locally equivalent, for each i , $0 \leq i \leq 3$.*

Proof. The proof is again by successive construction for each i , $0 \leq i \leq 3$. \square

Next, for $(UT)^*$ we define the following first-order equivalent:

$$\begin{aligned} \text{foe}(UT)^* &= (\text{foe}(\text{Lin} \diamond A_0^3)^* \wedge \forall y \forall z_1 \forall z_2 \forall u \forall v (CR(x, y, z_1, z_2) \wedge xRu \wedge xRv \\ &\wedge \exists w_0 (uRw_0 \wedge vRw_0 \wedge \text{ch}_0(w_0)) \wedge \exists w_1 (uRw_1 \wedge vRw_1 \wedge \text{ch}_1(w_1)) \\ &\wedge \exists w_2 (uRw_2 \wedge vRw_2 \wedge \text{ch}_2(w_2)) \wedge \exists w_3 (uRw_3 \wedge vRw_3 \wedge \neg uR^2w_3 \\ &\wedge \neg vR^2w_3 \wedge \text{ch}_3(w_3)) \rightarrow u=v). \end{aligned}$$

Lemma 2.10 *The formulas $(UT)^*$ and $\text{foe}(UT)^*$ are locally equivalent.*

Proof. Via a calculation using all previous Lemmas of this Section. \square




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9

Next, we get to the conjuncts simulating the instructions of our Minsky Machine. For future use, we make the following abbreviations (where u, w_1, w_2 are always free variables):

$$\begin{aligned} \text{fodS}(\delta, Q_1, R_1) = & \\ \exists t_1 \exists t_2 \dots \exists t_{\delta+1} (uRt_1 Rt_2 \dots Rt_{\delta+1} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+1})) & \\ \wedge \neg \exists t_1 \exists t_2 \dots \exists t_{\delta+2} (uRt_1 Rt_2 \dots Rt_{\delta+2} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+2})) \wedge uRw_1 & \\ \wedge \neg uR^2 w_1 \wedge \text{ch}_1(w_1) \wedge uRw_2 \wedge \neg uR^2 w_2 \wedge \text{ch}_2(w_2), & \end{aligned}$$

$$\begin{aligned} \text{fodS}(\delta, Q_2, R_1) = & \\ \exists t_1 \exists t_2 \dots \wedge \exists t_{\delta+1} (uRt_1 Rt_2 \dots Rt_{\delta+1} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+1})) & \\ \wedge \neg \exists t_1 \exists t_2 \dots \exists t_{\delta+2} (uRt_1 Rt_2 \dots Rt_{\delta+2} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+2})) \wedge uR^2 w_1 & \\ \wedge \neg uR^3 w_1 \wedge \text{ch}_1(w_1) \wedge uRw_2 \wedge \neg uR^2 w_2 \wedge \text{ch}_2(w_2), & \end{aligned}$$

$$\begin{aligned} \text{fodS}(\delta, Q_1, R_2) = & \\ \exists t_1 \exists t_2 \dots \exists t_{\delta+1} (uRt_1 Rt_2 \dots Rt_{\delta+1} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+1})) \wedge & \\ \wedge \neg \exists t_1 \exists t_2 \dots \exists t_{\delta+2} (uRt_1 Rt_2 \dots Rt_{\delta+2} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+2})) \wedge uRw_1 & \\ \wedge \neg uR^2 w_1 \wedge \text{ch}_1(w_1) \wedge uR^2 w_2 \wedge \neg uR^3 w_2 \wedge \text{ch}_2(w_2), & \end{aligned}$$

$$\begin{aligned} \text{fodS}(\delta, A_0^1, R_1) = & \\ \exists t_1 \exists t_2 \dots \exists t_{\delta+1} (uRt_1 Rt_2 \dots Rt_{\delta+1} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+1})) & \\ \wedge \neg \exists \tau_1 \exists t_2 \dots \exists t_{\delta+2} (uRt_1 Rt_2 \dots Rt_{\delta+2} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+2})) \wedge uRw_1 & \\ \wedge \text{ch}_1(w_1) \wedge \forall w' (uR^2 w' \rightarrow \neg \text{ch}_1(w')) \wedge uRw_2 \wedge \neg uR^2 w_2 \wedge \text{ch}_2(w_2), & \end{aligned}$$

$$\begin{aligned} \text{fodS}(\delta, Q_1, A_0^2) = & \\ \exists t_1 \exists t_2 \dots \exists t_{\delta+1} (uRt_1 Rt_2 \dots Rt_{\delta+1} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+1})) & \\ \wedge \neg \exists \tau_1 \exists t_2 \dots \exists t_{\delta+2} (uRt_1 Rt_2 \dots Rt_{\delta+2} \wedge \text{ch}_0(t_1) \wedge \text{ch}_0(t_2) \wedge \dots \wedge \text{ch}_0(t_{\delta+2})) \wedge uRw_1 & \\ \wedge \neg uR^2 w_1 \wedge \text{ch}_1(w_1) \wedge uRw_2 \wedge \text{ch}_2(w_2) \wedge \forall w' (uR^2 w' \rightarrow \neg \text{ch}_2(w')). & \end{aligned}$$

Now we can continue our definitions. For any instruction I we write:

- if $I = q_\delta \rightarrow q_\delta T_1 T_0$, then $\text{foe}((UT)^* \wedge AxI) = \text{foe}(UT)^* \wedge \forall y \forall z_1 \forall z_2 \forall w_1 \forall w_2 \forall w_3 (CR(x, y, z_1, z_2) \wedge \exists u (xRu \wedge \text{fodS}(\delta, Q_1, R_1) \wedge uRw_3 \wedge \neg uR^2 w_3 \wedge \text{ch}_3(w_3)) \wedge \exists w'_3 (xRw'_3 \wedge w'_3 R w_3 \wedge \text{ch}_3(w'_3)) \rightarrow \exists u (xRu \wedge \text{fodS}(\delta, Q_2, R_1) \wedge \exists w'_3 (uRw'_3 R w_3 \wedge \neg w'_3 R^2 w_3 \wedge \text{ch}_3(w'_3))))$,
- if $I = q_\delta \rightarrow q_\delta T_0 T_1$, then the definition of $\text{foe}((UT)^* \wedge AxI)$ can be obtained from the previous one by substituting $\text{fodS}(\delta, Q_1, R_2)$ for $\text{fodS}(\delta, Q_2, R_1)$,
- if $I = q_\delta \rightarrow q_\delta T_{-1} T_0 (q_\varepsilon T_0 T_0)$, then $\text{foe}((UT)^* \wedge AxI) = \text{foe}(UT)^* \wedge \forall y \forall z_1 \forall z_2 \forall w_1 \forall w_2 \forall w_3 (CR(x, y, z_1, z_2) \wedge \exists u (xRu \wedge \text{fodS}(\delta, Q_2, R_1) \wedge uRw_3 \wedge \neg uR^2 w_3 \wedge \text{ch}_3(w_3)) \wedge \exists w'_3 (xRw'_3 \wedge w'_3 R w_3 \wedge \text{ch}_3(w'_3)) \rightarrow \exists u (xRu \wedge \text{fodS}(\delta, Q_1, R_1) \wedge \exists w'_3 (uRw'_3 R w_3 \wedge \neg w'_3 R^2 w_3 \wedge \text{ch}_3(w'_3)))) \wedge \forall y \forall z_1 \forall z_2 \forall w_1 \forall w_2 \forall w_3 (CR(x, y, z_1, z_2) \wedge \exists u (xRu \wedge \text{fodS}(\delta, A_0^1, R_1) \wedge uRw_3 \wedge \neg uR^2 w_3 \wedge \text{ch}_3(w_3)) \wedge \exists w'_3 (xRw'_3 \wedge w'_3 R w_3 \wedge \text{ch}_3(w'_3)) \rightarrow \exists u (xRu \wedge \text{fodS}(\delta, A_0^1, R_1) \wedge \exists w'_3 (uRw'_3 R w_3 \wedge \neg w'_3 R^2 w_3 \wedge \text{ch}_3(w'_3))))$,
- if $I = q_\delta \rightarrow q_\delta T_0 T_{-1} (q_\varepsilon T_0 T_0)$, then the definition of $\text{foe}((UT)^* \wedge AxI)$ can be obtained from the previous one by substituting $\text{fodS}(\delta, Q_1, R_2)$ for $\text{fodS}(\delta, Q_2, R_1)$, and $\text{fodS}(\delta, Q_1, A_0^2)$ instead of $\text{fodS}(\delta, A_0^1, R_1)$.

Lemma 2.11 *The formulas $(UT)^* \wedge AxI$ and $\text{foe}((UT)^* \wedge AxI)$ are locally equivalent, for any instruction I .*

Finally, we define

$$\text{foe}(AxP)^* = \bigwedge_{I \in P} \text{foe}((UT)^* \wedge AxI).$$

Evidently, local equivalence of φ_1 and ψ_1 , φ_2 and ψ_2 implies local equivalence of $\varphi_1 \wedge \varphi_2$ and $\psi_1 \wedge \psi_2$, and hence Lemma 2.11 implies:

Lemma 2.12 *The formulas $(AxP)^*$ and $\text{foe}(AxP)^*$ are locally equivalent.*

Because $(AxP)^*$ is $A(P)$, Lemma 2.1. is then obtained from Lemma 2.12. Now change $A(P)$, adding a conjunct $\diamond T$. From Lemma 2.12, we also get

Lemma 2.13 *The formulas $A(P) \wedge \diamond T$ and $\text{foe}(AxP)^* \wedge \exists yxRy$ are locally equivalent, and so $A(P) \wedge \diamond T$ is locally first-order definable.*

Note that a global first-order equivalent of $A(P) \wedge \diamond T$ is \perp or, for example, $\forall x(x \neq x)$.

It only remains to be verified that our new first-order version of $A(P)$ works as before. Consider the proof of the main technical Lemma 1.2 from Section 1. Its "if" part depended on the conjuncts LF , $LinT$, UT , AxP , but these still occur in the new variant of $A(P)$. For its "only if" part, it was important that $A(P)$ should be true in the frame of Figure 1, and that $A(P) \wedge \diamond T$ be true in world f of this frame. For the latter purpose, because of Lemmas 2.12 and 2.13, it suffices to have the two formulas $\forall x \text{foe}(AxP)^*$ and $\text{foe}(AxP)^* \wedge \exists yxRy(x/f)$ true in this frame. And this is obvious from their definitions (which were in fact inspired by Figure 1) plus the proofs of Lemmas 2.12 and 2.13. Thus, we obtain the following results:

Proposition 2.14 *For the variant of the formula $A(P)$ defined in this chapter, Lemma 1.2 holds.*

Proposition 2.15 *For the variant of the formula $A(P)$ defined in this chapter, Lemma 1.2 also holds with \vDash_{fin}^{loc} instead of \vDash_{fin} .*

3 Modal Formulas Without First-Order Equivalents On Finite Frames

In this Section, we consider the part of Figure 1 surrounded by the dotted line. Essentially, the results formulated here are small modifications of results obtained by K. Doets in [8]. First we formulate some key facts which we shall need.

Recall the definition of *n-equivalence* of models, notation: $A \equiv_n B$. We use game-theoretical terminology as in [9], [8] ([7] has this notion in a different but equivalent form). Note that we have only equality and one binary relation in the signature (or language) of our models. Now the *n-game* on A and B , $G(A, B, n)$, has two players, I and II, which move alternatively. I is allowed the first move; each player is allowed n moves altogether. A move consists in the selection of an element in either W_1 or W_2 (where W_1 and W_2 are the universes of A and B , respectively). However, if player I chooses an element in W_1 (W_2), then player II has to counter in W_2 (W_1). Therefore, a move of player I and the following counter-move of player II form an ordered pair in $W_1 \times W_2$. When the game is over, the set of ordered pairs of moves is at most an n -element relation $h \subseteq W_1 \times W_2$. II has won the play by definition if h is a partial isomorphism between W_1 and W_2 . Now, A and B are called '*n-equivalent*' if II has a winning strategy for $G(A, B, n)$. Here is the basic logical property of \equiv_n . If $A \equiv_n B$ then, for any first-order formula φ of quantifier depth at most n ,

$$A \vDash \varphi \text{ iff } B \vDash \varphi.$$

Following [8], let PZ_k be the nontransitive frame of Figure 10. The painted circles represent reflexive worlds.

Lemma 3.1 [8]. *If $k, m \geq 2^n$, then $PZ_k \equiv_n PZ_m$.*




Figure 10




Figure 11




Figure 12

If we now take the McKinsey Formula $\Box\Diamond p \rightarrow \Diamond\Box p$, then it is noted, once more in [10], that this formula is true in PZ_k iff k is odd. If one then supposes that this formula has a first-order equivalent (global or local), say φ of quantifier depth n , then we get that both $PZ_{2^n} \models \varphi$ and $PZ_{2^{n+1}} \models \varphi$, although $PZ_{2^n} \not\models \Box\Diamond p \rightarrow \Diamond\Box p$, $PZ_{2^{n+1}} \models \Box\Diamond p \rightarrow \Diamond\Box p$: which is a contradiction. Thus, the McKinsey Formula is not first-order definable (locally or globally) on finite frames ([8]). We shall modify the formula and frames employed in this argument in a manner suitable for our purposes. Call the frame in Figure 11 $FENCE_k$.

Lemma 3.2 *If $k, m \geq 2^n$, then $FENCE_k \equiv_n FENCE_m$.*

Proof. Use the winning strategy for player II from the proof of Lemma 3.1. \square

If we take the frames from Figure 12, denoting them by $FENCE^*_k$ for future reference, then Lemma 3.2 implies that, if $k, m \geq 2^n$, then $FENCE^*_k \equiv_n FENCE^*_m$. Indeed the strategy from the proof of Lemma 3.1 is still suitable, with the following stipulation added:

"if player I chooses f in one model, then II chooses f in the other model".

Next, we use a modified McKinsey Formula of the form

$$\alpha = \underline{\Diamond\Diamond T \wedge \Box(\underline{\Diamond T} \rightarrow \Diamond p)} \rightarrow \underline{\Diamond(\underline{\Diamond T \wedge \Box p})}.$$

The underlined parts of α represent changes to the original McKinsey Formula. It is easy to see that α is refutable in $FENCE^*_k$ iff k is odd, and the refutation can occur only in the world f . Thus α is not first-order definable on the class of finite frames, and not even in the class of finite transitive irreflexive frames. These observations establish the following

Proposition 3.3 *The formula $(\Box(\Box p \rightarrow p) \rightarrow \Box p) \wedge \alpha$ is not first-order definable on the class of finite frames.*

Proposition 3.4 *The formula $(\Box(\Box p \rightarrow p) \rightarrow \Box p) \wedge \alpha \wedge \Diamond T$ is not locally first-order definable on the finite frames, although is globally first-order definable on them.*

Proof. A global first-order equivalent of this formula is, for example, $\forall x x \neq x$. The absence of a local equivalent is proved as above, but now using the frames $FENCE^*_k$, in which f is the 'real world'. \square

Remark. Consider the modal formula $\beta = \Box^3 \perp \wedge (\Box(\Box p \rightarrow p) \rightarrow \Box p) \wedge \alpha \wedge \Diamond T$.

It may be proved analogously that β is globally first-order definable on the class of all transitive frames, but it is not locally first-order definable on the class of all transitive frames. This answers a question by van Benthem ([1], p.129). \square

The frame sketched in Figure 1 is obtained from $FENCE_t$ by adding some subframe. Let us call it $F_t(P, \alpha, m, n)$.

Lemma 3.5 *If $t_1, t_2 \geq 2^l$, then $F_{t_1}(P, \alpha, m, n) \equiv_l F_{t_2}(P, \alpha, m, n)$.*

Proof. Player II needs the strategy from the proof of Lemma 3.2 with one additional rule: "if I chooses some element out of $FENCE_{t_i}$ in $F_{t_i}(P, \alpha, m, n)$, then II chooses the same element in $F_{t_{1-i}}(P, \alpha, m, n)$ ". \square

Now we explain the modified McKinsey Formula, i.e., the underlined parts of formula α . The part $\underline{\Diamond\Diamond T \wedge}$ has been added to make sure that a possible refutation could occur only at world f in the frame $FENCE^*_k$. The parts $\underline{\Diamond T \rightarrow}$ and $\underline{\Diamond T \wedge}$ were added to give some information just about the worlds d_1, \dots, d_k in $FENCE^*_k$. Define the following further modification of the McKinsey Formula (changes are again underlined):

$$\gamma = \wedge F \wedge \Box(\underline{\Diamond\Diamond B} \rightarrow \underline{\Diamond(\underline{\Diamond B \wedge q})}) \rightarrow \underline{\Diamond(\underline{\Diamond\Diamond B} \wedge \Box(\underline{\Diamond B} \rightarrow q))},$$

(here q is used instead of p , because p was already used in F), where

$$B = \Diamond F_0^7 \wedge \Diamond F_1^7 \wedge \Box^8 \perp.$$

From Lemmas 1.4, 1.5, we then obtain the following fact:

Lemma 3.6. *The formula $A(P) \wedge (B(\alpha, m, n) \vee \delta)$ is refutable in the frame $F_t(P, \alpha, m, n)$ iff t is odd, and the refutation can take place only in the world f .*

As a consequence of Lemmas 3.5 and 3.6, we obtain one final

Proposition 3.7 *If a program P , starting at a configuration (α, m, n) halts in a final state with the number β , then :*

- i) *the formula $A(P) \wedge (B(\alpha, m, n) \vee \delta)$ is not first-order definable on finite frames (either locally or globally);*
- ii) *the formula $A(P) \wedge (B(\alpha, m, n) \vee \delta) \wedge \Diamond T$ is globally first-order definable on finite frames, but is not locally first-order definable on them.*

4 Main Results

Now we can formulate the main results of this paper. Here, $A(P)$ is defined as in Section 2.

Lemma 4.1 *The following three conditions are equivalent for any program P and configuration (α, m, n) :*

- i) *program P , starting from configuration (α, m, n) , cannot reach a final state (with number β),*
- ii) *the formula $A(P) \wedge (B(\alpha, m, n) \vee \delta)$ is locally first-order definable on finite frames,*
- iii) *the formula $A(P) \wedge (B(\alpha, m, n) \vee \delta)$ is globally first-order definable on finite frames.*

Proof. i) \Rightarrow ii). If i) holds, then by Lemma 1.2, Proposition 2.14 and the Remark preceding Theorem 1.10, in any frame with a designated world where $A(P)$ is true, $B(\alpha, m, n)$ is true as well. So the formula $A(P) \wedge (B(\alpha, m, n) \vee \delta)$ is locally equivalent to the formula $A(P)$, which is locally first-order definable by Lemma 2.1.

ii) \Rightarrow iii). This direction is trivial.

iii) \Rightarrow i). Let i) be false. Then, by Proposition 3.7.i), iii) is not true either. \square

Because the Halting Problem for Minsky Machines is undecidable, and $A(P)$ and $B(\alpha, m, n)$ have been constructed effectively from P and (α, m, n) , Lemma 4.1 now implies two further results:

Theorem 4.2 *The problem of recognizing, given any modal formula, whether it is locally first-order definable on finite frames, is algorithmically undecidable.*

Theorem 4.3 *The problem of recognizing, given any modal formula, whether it is globally first-order definable on finite frames, is algorithmically undecidable.*

Theorems 4.2 and 4.3 are in a sense independent. Neither follows directly from the other, witness the following observation:

Lemma 4.4 *The following conditions are equivalent for any program P and configuration (α, m, n) :*

- i) *program P , starting at configuration (α, m, n) , halts in a final state (with number β);*
- ii) *the formula $A(P) \wedge (B(\alpha, m, n) \vee \delta) \wedge \Diamond T$ is globally first-order definable on finite frames, but is not locally first-order definable on finite frames.*

Proof. i) \Rightarrow ii). By Proposition 3.7.ii). ii) \Rightarrow i). Analogous to the proof of i) \Rightarrow ii), but now using Lemma 2.13 instead of Lemma 2.1. \square

From Lemma 4.4, we obtain our next result:

Theorem 4.5 *The set of modal formulas which are globally, but not locally first-order definable on finite frames, is algorithmically undecidable.*

We conclude with one natural question. The formula LF is first-order definable on finite frames, but is not first-order definable even on all countable frames.

Theorem 4.6 i) *The set of modal formulas which are first-order definable on the class of finite frames, but not first-order definable on the class of all frames, is algorithmically undecidable.*
 ii) *The set of modal formulas which are first-order definable on the class of finite frames, but not first-order definable on the class of all countable frames, is algorithmically undecidable.*

Proof An argument for i) and ii) can be given simultaneously. Return to the proof of Lemma 4.1. The proof of the implication iii) \Rightarrow i) showed in fact that, if i) is not true, then $A(P) \wedge (B(\alpha, m, n) \vee \delta)$ is not a formula, which is first-order definable on finite frames, but not first-order definable over all (countable) frames. In the case of the proof i) \Rightarrow ii), we observe that the formula $A(P)$ is not first-order definable in the class of all (countable) frames. This is proved via the usual counter-example to first-order definability for LF, using linear frames in which all conjuncts of $A(P)$ except LF are true. \square

Remark The fact that Theorem 4.6.i) and Theorem 4.6.ii) are independent follows from the undecidability (announced in [5]) of a set of formulas which is first-order definable on countable frames, but not first-order definable in general. \square

Question *Whether any of the sets of formulas mentioned in Theorems 4.2, 4.3, 4.5, 4.6, or their complements, are recursive enumerable.*

REFERENCES

- [1] J. F. A. K. van Benthem, *Modal Logic and Classical Logic*, Bibliopolis, Napoli, 1985.
- [2] -----, *Notes on Modal Definability*, *Notre Dame Journal of Formal Logic*, vol. 30 (1989), pp. 20-35.
- [3] A. V. Chagrov, *Undecidable Properties of Extensions of Provability Logic*. I, II, *Algebra i Logika*, vol. 29 (1990), pp. 350-367, 613-623; English translation, *Algebra and Logic*, vol. 29 (1990) (to appear).
- [4] -----, *Undecidability of Finite Semantic Consequence*, *Proceeding of 10th USSR Conference on Mathematical logic*, Alma-Ata, 1990, p. 162.
- [5] L. A. Chagrova, *An Undecidable Problem in Correspondence Theory*, *The Journal of Symbolic Logic*, vol. 56 (1991), 4, pp. 1261-1272.
- [6] -----, *An Undecidable Problem in Correspondence Theory for Finite Frames*, (in preparation).
- [7] C. C. Chang, H. J. Keisler, *Model Theory*, North-Holland, Amsterdam, 1973.
- [8] H. C. Doets, *Completeness and Definability, Applications of the Ehrenfeucht Game in Second-order and Intensional Logic*, Dissertation, Mathematical Institute, University of Amsterdam, 1987.
- [9] A. Ehrenfeucht, *An Application of Games to the Completeness Problem for Formalized Theories*, *Fundamenta Mathematicae*, vol. 49 (1961), pp. 128-141.
- [10] R. I. Goldblatt, *First-Order Definability in Modal Logic*, *The Journal of Symbolic Logic*, vol. 40 (1975), pp. 35-40.

The ILLC Prepublication Series

- CT-91-10 John Tromp, Paul Vitányi A Randomized Algorithm for Two-Process Wait-Free Test-and-Set
 CT-91-11 Lane A. Hemachandra, Édith Spaan Quasi-Injective Reductions
 CT-91-12 Krzysztof R. Apt, Dino Pedreschi Reasoning about Termination of Prolog Programs
- Computational Linguistics*
 CL-91-01 J.C. Scholtes Kohonen Feature Maps in Natural Language Processing
 CL-91-02 J.C. Scholtes Neural Nets and their Relevance for Information Retrieval
 CL-91-03 Hub Prüst, Remko Scha, Martin van den Berg A Formal Discourse Grammar tackling Verb Phrase Anaphora
- Other Prepublications*
 X-91-01 Alexander Chagrov, Michael Zakharyashev The Disjunction Property of Intermediate Propositional Logics
 X-91-02 Alexander Chagrov, Michael Zakharyashev On the Undecidability of the Disjunction Property of Intermediate Propositional Logics
 X-91-03 V. Yu. Shavrukov Subalgebras of Diagonalizable Algebras of Theories containing Arithmetic
 X-91-04 K.N. Ignatiev Partial Conservativity and Modal Logics
 X-91-05 Johan van Benthem Temporal Logic
 X-91-06 Annual Report 1990
 X-91-07 A.S. Troelstra Lectures on Linear Logic, Errata and Supplement
 X-91-08 Giorgie Dzhaparidze Logic of Tolerance
 X-91-09 L.D. Beklemishev On Bimodal Provability Logics for Π_1 -axiomatized Extensions of Arithmetical Theories
 X-91-10 Michiel van Lambalgen Independence, Randomness and the Axiom of Choice
 X-91-11 Michael Zakharyashev Canonical Formulas for K4. Part I: Basic Results
 X-91-12 Herman Hendriks Flexibele Categoriale Syntaxis en Semantiek: de proefschriften van Frans Zwarts en Michael Moortgat
 X-91-13 Max I. Kanovich The Multiplicative Fragment of Linear Logic is NP-Complete
 X-91-14 Max I. Kanovich The Horn Fragment of Linear Logic is NP-Complete
 X-91-15 V. Yu. Shavrukov Subalgebras of Diagonalizable Algebras of Theories containing Arithmetic, revised version
 X-91-16 V.G. Kanovei Undecidable Hypotheses in Edward Nelson's Internal Set Theory
 X-91-17 Michiel van Lambalgen Independence, Randomness and the Axiom of Choice, Revised Version
 X-91-18 Giovanna Cepparello New Semantics for Predicate Modal Logic: an Analysis from a standard point of view
 X-91-19 Papers presented at the Provability Interpretability Arithmetic Conference, 24-31 Aug. 1991, Dept. of Phil., Utrecht University
1992
Logic, Semantics and Philosophy of Language
 LP-92-01 Víctor Sánchez Valencia Lambek Grammar: an Information-based Categorical Grammar
 LP-92-02 Patrick Blackburn Modal Logic and Attribute Value Structures
 LP-92-03 Szabolcs Mikulás The Completeness of the Lambek Calculus with respect to Relational Semantics
 LP-92-04 Paul Dekker An Update Semantics for Dynamic Predicate Logic
 LP-92-05 David I. Beaver The Kinematics of Presupposition
 LP-92-06 Patrick Blackburn, Edith Spaan A Modal Perspective on the Computational Complexity of Attribute Value Grammar
 LP-92-07 Jeroen Groenendijk, Martin Stokhof A Note on Interrogatives and Adverbs of Quantification
 LP-92-08 Maarten de Rijke A System of Dynamic Modal Logic
 LP-92-09 Johan van Benthem Quantifiers in the world of Types
 LP-92-10 Maarten de Rijke Meeting Some Neighbours (a dynamic modal logic meets theories of change and knowledge representation)
 LP-92-11 Johan van Benthem A note on Dynamic Arrow Logic
 LP-92-12 Heinrich Wansing Sequent Calculi for Normal Modal Propositional Logics
 LP-92-13 Dag Westerståhl Iterated Quantifiers
 LP-92-14 Jeroen Groenendijk, Martin Stokhof Interrogatives and Adverbs of Quantification
Mathematical Logic and Foundations
 ML-92-01 A.S. Troelstra Comparing the theory of Representations and Constructive Mathematics
 ML-92-02 Dmitriy P. Skvortsov, Valentin B. Shehtman Maximal Kripke-type Semantics for Modal and Superintuitionistic Predicate Logics
 ML-92-03 Zoran Marković On the Structure of Kripke Models of Heyting Arithmetic
 ML-92-04 Dimitar Vakarelov A Modal Theory of Arrows, Arrow Logics I
 ML-92-05 Domenico Zambella Shavrukov's Theorem on the Subalgebras of Diagonalizable Algebras for Theories containing $\Delta_0 + EXP$
 ML-92-06 D.M. Gabbay, Valentin B. Shehtman Undecidability of Modal and Intermediate First-Order Logics with Two Individual Variables
 ML-92-07 Harold Schellinx How to Broaden your Horizon
 ML-92-08 Raymond Hoofman Information Systems as Coalgebras
 ML-92-09 A.S. Troelstra Realizability
 ML-92-10 V.Yu. Shavrukov A Smart Child of Peano's
- Computation and Complexity Theory*
 CT-92-01 Erik de Haas, Peter van Emde Boas Object Oriented Application Flow Graphs and their Semantics
 CT-92-02 Karen L. Kwast, Sieger van Denneheuvel Weak Equivalence: Theory and Applications
 CT-92-03 Krzysztof R. Apt, Kees Doets A new Definition of SLDNF-resolution
- Other Prepublications*
 X-92-01 Heinrich Wansing The Logic of Information Structures
 X-92-02 Konstantin N. Ignatiev The Closed Fragment of Dzhaparidze's Polymodal Logic and the Logic of Σ_1 conservativity
 X-92-03 Willem Groeneveld Dynamic Semantics and Circular Propositions, revised version
 X-92-04 Johan van Benthem Modeling the Kinematics of Meaning
 X-92-05 Erik de Haas, Peter van Emde Boas Object Oriented Application Flow Graphs and their Semantics, revised version
1993
Logic, Semantics and Philosophy of Language
 LP-93-01 Martijn Spaan Parallel Quantification
 LP-93-02 Makoto Kanazawa Dynamic Generalized Quantifiers and Monotonicity
Mathematical Logic and Foundations
 ML-93-01 Maciej Kandulski Commutative Lambek Categorical Grammars
 ML-93-02 Johan van Benthem, Natasha Alechina Modal Quantification over Structured Domains
 ML-93-03 Mati Pentus The Conjoinability Relation in Lambek Calculus and Linear Logic
 ML-93-04 Andreja Prijatelj Bounded Contraction and Many-Valued Semantics
 ML-93-05 Raymond Hoofman, Harold Schellinx Models of the Untyped λ -calculus in Semi Cartesian Closed Categories
 ML-93-06 J. Zashev Categorical Generalization of Algebraic Recursion Theory
 ML-93-07 A.V. Chagrov, L.A. Chagrova Algorithmic Problems Concerning First-Order Definability of Modal Formulas on the Class of All Finite Frames
- Computation and Complexity Theory*
 CT-93-01 Marianne Kalsbeek The Vanilla Meta-Interpreter for Definite Logic Programs and Ambivalent Syntax
 CT-93-02 Sophie Fischer A Note on the Complexity of Local Search problems
Other Prepublications
 X-93-01 Paul Dekker Existential Disclosure, revised version