




Institute for Logic, Language and Computation

EMBEDDINGS OF HEYTING ALGEBRAS

Dick de Jongh
Albert Visser

ILLC Prepublication Series
for Mathematical Logic and Foundations ML-93-14



University of Amsterdam

The ILLC Prepublication Series

1990 *Logic, Semantics and Philosophy of Language*

- LP-90-01 Jaap van der Does A Generalized Quantifier Logic for Naked Infinitives
 LP-90-02 Jeroen Groenendijk, Martin Stokhof Dynamic Montague Grammar
 LP-90-03 Renate Bartsch Concept Formation and Concept Composition
 LP-90-04 Aarne Ranta Intuitionistic Categorical Grammar
 LP-90-05 Patrick Blackburn Nominal Tense Logic
 LP-90-06 Gennaro Chierchia The Variability of Impersonal Subjects
 LP-90-07 Gennaro Chierchia Anaphora and Dynamic Logic
 LP-90-08 Herman Hendriks Flexible Montague Grammar
 LP-90-09 Paul Dekker The Scope of Negation in Discourse, towards a Flexible Dynamic Montague grammar
 LP-90-10 Theo M.V. Janssen Models for Discourse Markers
 LP-90-11 Johan van Benthem General Dynamics
 LP-90-12 Serge Lapierre A Functional Partial Semantics for Intensional Logic
 LP-90-13 Zhisheng Huang Logics for Belief Dependence
 LP-90-14 Jeroen Groenendijk, Martin Stokhof Two Theories of Dynamic Semantics
 LP-90-15 Maarten de Rijke The Modal Logic of Inequality
 LP-90-16 Zhisheng Huang, Karen Kwast Awareness, Negation and Logical Omniscience
 LP-90-17 Paul Dekker Existential Disclosure, Implicit Arguments in Dynamic Semantics

Mathematical Logic and Foundations

- ML-90-01 Harold Schellinx Isomorphisms and Non-Isomorphisms of Graph Models
 ML-90-02 Jaap van Oosten A Semantical Proof of De Jongh's Theorem
 ML-90-03 Yde Venema Relational Games
 ML-90-04 Maarten de Rijke Unary Interpretability Logic
 ML-90-05 Domenico Zambella Sequences with Simple Initial Segments
 ML-90-06 Jaap van Oosten Extension of Lifschitz' Realizability to Higher Order Arithmetic, and a Solution to a Problem of F. Richman
 ML-90-07 Maarten de Rijke A Note on the Interpretability Logic of Finitely Axiomatized Theories
 ML-90-08 Harold Schellinx Some Syntactical Observations on Linear Logic
 ML-90-09 Dick de Jongh, Duccio Pianigiani Solution of a Problem of David Guaspari
 ML-90-10 Michiel van Lambalgen Randomness in Set Theory
 ML-90-11 Paul C. Gilmore The Consistency of an Extended NaDSet

Computation and Complexity Theory

- CT-90-01 John Tromp, Peter van Emde Boas Associative Storage Modification Machines
 CT-90-02 Sieger van Denneheuvel, Gerard R. Renardel de Lavalette A Normal Form for PCSJ Expressions
 CT-90-03 Ricard Gavaldà, Leen Torenvliet, Osamu Watanabe, José L. Balcázar Generalized Kolmogorov Complexity in Relativized Separations

- CT-90-04 Harry Buhrman, Edith Spaan, Leen Torenvliet Bounded Reductions
 CT-90-05 Sieger van Denneheuvel, Karen Kwast Efficient Normalization of Database and Constraint Expressions
 CT-90-06 Michiel Smid, Peter van Emde Boas Dynamic Data Structures on Multiple Storage Media, a Tutorial
 CT-90-07 Kees Doets Greatest Fixed Points of Logic Programs
 CT-90-08 Fred de Geus, Ernest Rotterdam, Sieger van Denneheuvel, Peter van Emde Boas Physiological Modelling using RL
 CT-90-09 Roel de Vrijer Unique Normal Forms for Combinatory Logic with Parallel

Other Prepublications

- X-90-01 A.S. Troelstra Conditional, a case study in conditional rewriting
 X-90-02 Maarten de Rijke Remarks on Intuitionism and the Philosophy of Mathematics, Revised Version
 X-90-03 L.D. Beklemishev Some Chapters on Interpretability Logic
 X-90-04 On the Complexity of Arithmetical Interpretations of Modal Formulae
 X-90-05 Valentin Shehtman Annual Report 1989
 X-90-06 Valentin Goranko, Solomon Passy Derived Sets in Euclidean Spaces and Modal Logic
 X-90-07 V.Yu. Shavrukov Using the Universal Modality: Gains and Questions
 X-90-08 L.D. Beklemishev The Lindenbaum Fixed Point Algebra is Undecidable
 X-90-09 V.Yu. Shavrukov Provability Logics for Natural Turing Progressions of Arithmetical Theories
 X-90-10 Sieger van Denneheuvel, Peter van Emde Boas On Rosser's Provability Predicate
 X-90-11 Alessandra Carbone An Overview of the Rule Language RL/1
 X-90-12 Maarten de Rijke Provable Fixed points in $\Lambda_0 + \Omega_1$, revised version
 X-90-13 K.N. Ignatiev Bi-Unary Interpretability Logic
 X-90-14 L.A. Chagrova Dzhaparidze's Polymodal Logic: Arithmetical Completeness, Fixed Point Property, Craig's Property
 X-90-15 A.S. Troelstra Undecidable Problems in Correspondence Theory
 Lectures on Linear Logic

1991 *Logic, Semantics and Philosophy of Language*


- LP-91-01 Wiebe van der Hoek, Maarten de Rijke Generalized Quantifiers and Modal Logic
 LP-91-02 Frank Veltman Defaults in Update Semantics
 LP-91-03 Willem Groeneveld Dynamic Semantics and Circular Propositions
 LP-91-04 Makoto Kanazawa The Lambek Calculus enriched with Additional Connectives
 LP-91-05 Zhisheng Huang, Peter van Emde Boas The Schoenmakers Paradox: Its Solution in a Belief Dependence Framework
 LP-91-06 Zhisheng Huang, Peter van Emde Boas Belief Dependence, Revision and Persistence
 LP-91-07 Henk Verkuyl, Jaap van der Does The Semantics of Plural Noun Phrases
 LP-91-08 Víctor Sánchez Valencia Categorical Grammar and Natural Reasoning
 LP-91-09 Arthur Nieuwendijk Semantics and Comparative Logic
 LP-91-10 Johan van Benthem Logic and the Flow of Information

Mathematical Logic and Foundations

- ML-91-01 Yde Venema Cylindric Modal Logic
 ML-91-02 Alessandro Berarducci, Rineke Verbrugge On the Metamathematics of Weak Theories
 ML-91-03 Domenico Zambella On the Proofs of Arithmetical Completeness for Interpretability Logic
 ML-91-04 Raymond Hoofman, Harold Schellinx Collapsing Graph Models by Preorders
 ML-91-05 A.S. Troelstra History of Constructivism in the Twentieth Century
 ML-91-06 Inge Bethke Finite Type Structures within Combinatory Algebras
 ML-91-07 Yde Venema Modal Derivation Rules
 ML-91-08 Inge Bethke Going Stable in Graph Models
 ML-91-09 V.Yu. Shavrukov A Note on the Diagonalizable Algebras of PA and ZF
 ML-91-10 Maarten de Rijke, Yde Venema Sahlqvist's Theorem for Boolean Algebras with Operators
 ML-91-11 Rineke Verbrugge Feasible Interpretability
 ML-91-12 Johan van Benthem Modal Frame Classes, revisited

Computation and Complexity Theory

- CT-91-01 Ming Li, Paul M.B. Vitányi Kolmogorov Complexity Arguments in Combinatorics
 CT-91-02 Ming Li, John Tromp, Paul M.B. Vitányi How to Share Concurrent Wait-Free Variables
 CT-91-03 Ming Li, Paul M.B. Vitányi Average Case Complexity under the Universal Distribution Equals Worst Case Complexity
 CT-91-04 Sieger van Denneheuvel, Karen Kwast Weak Equivalence
 CT-91-05 Sieger van Denneheuvel, Karen Kwast Weak Equivalence for Constraint Sets
 CT-91-06 Edith Spaan Census Techniques on Relativized Space Classes
 CT-91-07 Karen L. Kwast The Incomplete Database
 CT-91-08 Kees Doets Levationis Laus
 CT-91-09 Ming Li, Paul M.B. Vitányi Combinatorial Properties of Finite Sequences with high Kolmogorov Complexity
 CT-91-10 John Tromp, Paul Vitányi A Randomized Algorithm for Two-Process Wait-Free Test-and-Set
 CT-91-11 Lane A. Hemachandra, Edith Spaan Quasi-Injective Reductions
 CT-91-12 Krzysztof R. Apt, Dino Pedreschi Reasoning about Termination of Prolog Programs
 CL-91-01 J.C. Scholtes Computational Linguistics Kohonen Feature Maps in Natural Language Processing
 CL-91-02 J.C. Scholtes Neural Nets and their Relevance for Information Retrieval
 CL-91-03 Hub Prüst, Remko Scha, Martin van den Berg A Formal Discourse Grammar tackling Verb Phrase Anaphora



Institute for Logic, Language and Computation

Plantage Muidergracht 24

1018TV Amsterdam

Telephone 020-525.6051, Fax: 020-525.5101

EMBEDDINGS OF HEYTING ALGEBRAS

Dick de Jongh

Department of Mathematics and Computer Science
University of Amsterdam

Albert Visser

Department of Philosophy
University of Utrecht
Heidelberglaan 8, Postbus 80126, 3508TC Utrecht

Embeddings of Heyting Algebras

*Dick de Jongh &
Albert Visser*

ABSTRACT: In this paper we study embeddings of Heyting Algebras. It is pointed out that such embeddings are naturally connected with Derived Rules. We compare the Heyting Algebras embeddable in the Heyting Algebra of the Intuitionistic Propositional Calculus (IPC), i.e. the free Heyting Algebra on countably infinitely many generators, and those embeddable in the Heyting Algebra of Heyting's Arithmetic (HA). A partial result is obtained. We show that every recursively enumerable prime Heyting Algebra is embeddable in the Heyting Algebra of HA^* , a 'natural' extension of HA.

1 Introduction

This paper sprung from an interest in the Heyting Algebras of Constructive Arithmetical Theories. This interest was in its turn inspired by an interest in the Propositional Derived Rules of constructive Arithmetical Theories. We study and compare four specific Heyting Algebras in some detail:

- The free Heyting Algebra on countably infinitely many generators, in other words: the Heyting Algebra \mathfrak{H}_{IPC} of the Intuitionistic Propositional Calculus (IPC).
- The Heyting Algebra \mathfrak{H}_{HA} of Heyting's Arithmetic (HA).
- The Heyting Algebra \mathfrak{S}_{HA} of $B\Sigma_1$ -sentences in HA (here $B\Sigma_1$ is the set of Boolean (or perhaps more appropriately: Brouwerian) combinations of Σ_1 -sentences).
- The Heyting Algebra \mathfrak{H}_{HA^*} of HA^* , an arithmetical theory studied in Visser[82].

We ask ourselves which RE Heyting Algebras can be embedded in our target algebras. As we will see the answer to this question also determines what the Propositional Derived Rules for the various theories are. A complete answer has only been obtained for \mathfrak{H}_{HA^*} . All RE algebras of which one could reasonably expect it, i.e. those satisfying the property of primeness (corresponding to satisfying the disjunction property), are embeddable in \mathfrak{H}_{HA^*} , and in consequence, only rules directly derivable in intuitionistic logic are rules under which HA^* is closed. This property of HA^* is a nice one —and in a surprising manner enables one to prove some properties of HA itself— but it does not seem to hold for more usual theories. Neither in \mathfrak{H}_{HA} , nor in \mathfrak{H}_{IPC} itself can all these algebras be embedded, since both these theories validate many

additional rules not derivable in intuitionistic logic, the best known being:

$$\neg A \rightarrow (B \vee C) / (\neg A \rightarrow B) \vee (\neg A \rightarrow C). \quad (\text{Independence of Premiss Rule})$$

It is even an interesting open question whether in \mathfrak{H}_{HA} and in $\mathfrak{H}_{\text{IPC}}$ the same Heyting algebras can be embedded. This is where \mathfrak{S}_{HA} comes in: it is possible to show that in \mathfrak{S}_{HA} and in $\mathfrak{H}_{\text{IPC}}$ at least the same finitely generated Heyting algebras can be embedded. It follows that rules validated by formal arithmetic when one restricts oneself to substitutions of propositional combinations of Σ_1 -sentences, and rules validated by the propositional calculus are the same.

We state some sample results:

- Any RE prime Heyting Algebra \mathfrak{H} can be embedded in $\mathfrak{H}_{\text{HA}^*}$. (5.1)
- There are Σ_1 -sentences A and B such that the subalgebra of $\mathfrak{H}_{\text{HA}^*}$ generated by A and B is RE, non-recursive. (7)
- Let \mathfrak{H} be a Heyting Algebra on finitely many generators, which is embeddable in $\mathfrak{H}_{\text{IPC}}$. Then \mathfrak{H} is the Heyting Algebra of a finitely axiomatizable IPC-theory. (2.3)
- Let \mathfrak{H} be a Heyting Algebra on finitely many generators. Then \mathfrak{H} is embeddable in \mathfrak{S}_{HA} iff \mathfrak{H} is embeddable in $\mathfrak{H}_{\text{IPC}}$. (6.2)

The paper is organized as follows. In section 1 we define Heyting algebras in the presentation most useful to our purposes. In section 2 we introduce the notion of embedding and a connected notion of propositional formulas exactly provable for sentences of a theory. Propositional formulas with no iterations of implications on the left (NNIL formulas) turn out to play an important role. In section 3 and 4 necessary facts about HA and IPC, and HA^* respectively, are given. In section 5 the above mentioned ‘RE universality’ of HA^* is proved. In section 6 \mathfrak{S}_{HA} is treated. Finally, in section 7 it is shown that, in consequence of the previous results, there is a Heyting algebra on two generators which is RE, but non-recursive, that can be embedded in $\mathfrak{H}_{\text{HA}^*}$, whereas such an algebra could never be embeddable in \mathfrak{S}_{HA} or $\mathfrak{H}_{\text{IPC}}$.

1.1 Acknowledgements: Most of the results of this paper were obtained during an exceptionally inspiring visit of both authors to the Katedra Logiky of Prague University and the Institute of Computer and Information Science of the Czech Academy of Sciences. We thank the Prague logicians for their wonderful hospitality.

Some of the main methods employed in this paper were invented by Volodya Shavrukov (see Shavrukov[93]) and further developed and simplified by Domenico Zambella (see Zambella[92]). The work of Shavrukov and Zambella concerns embeddings of RE Diagonalizable Algebras into Diagonalizable Algebras of Classical Arithmetical Theories.

A major tool of the present paper is also Pitts's Uniform Interpolation Theorem (see Pitts[92]).

1.2 The classical case: Before going on, let's briefly look at the Boolean Algebras of classical Arithmetical Theories. The Boolean Algebras of all consistent RE arithmetical theories extending Q are isomorphic to the free Boolean Algebra on \aleph_0 generators, i.e. to the Boolean Algebra of the Classical Propositional Calculus, say: $\mathfrak{B}_{\text{CPC}}$. As far as we can trace this result is folklore. It follows from three observations. First: the Boolean Algebras of all consistent RE arithmetical theories extending Q are countably infinite and (by Rosser's Theorem, see also 1.4) atomless. Second: $\mathfrak{B}_{\text{CPC}}$ is countably infinite and atomless. Third: all countably infinite atomless Boolean Algebras are isomorphic.

It is not difficult to show that every countable Boolean Algebra can be embedded into $\mathfrak{B}_{\text{CPC}}$.

1.3 Heyting Algebras: A Heyting Algebra \mathfrak{H} is a structure $\langle H, \wedge, \vee, \perp, \rightarrow \rangle$, where $\langle H, \wedge, \vee, \perp \rangle$ is a lattice with bottom \perp . We demand that \mathfrak{H} is non-trivial, i.e. that H contains at least two elements. Let $x \leq y$ be defined by $x \vee y = y$. \rightarrow is a binary operation satisfying: $x \wedge y \leq z \Leftrightarrow x \leq (y \rightarrow z)$. It is easily seen that if a partial order can be extended to a Heyting Algebra such an extension is unique. Heyting Algebras can be shown to be distributive lattices. Conversely every finite distributive lattice determines a Heyting Algebra.

There are many good sources for Heyting Algebras. We just mention van Troelstra & van Dalen[88b].

We will write:

- $\top := \perp \rightarrow \perp$,
- $\neg x := x \rightarrow \perp$,
- $x \leftrightarrow y := (x \rightarrow y) \wedge (y \rightarrow x)$,
- $\mathfrak{H} \models A(\mathbf{x}) :\Leftrightarrow A(\mathbf{x}) = \top$,

where A is a polynomial in $\wedge, \vee, \perp, \rightarrow$ and \mathbf{x} is a sequence of elements of \mathfrak{H} .

Note that \mathfrak{H} can be recovered from \models , since $A(\mathbf{x}) = B(\mathbf{y}) \Leftrightarrow \mathfrak{H} \models A(\mathbf{x}) \leftrightarrow B(\mathbf{y})$.

Define:

- $\mathfrak{f} : \mathfrak{H} \cong \mathfrak{R} :\Leftrightarrow \mathfrak{f}$ is an embedding of \mathfrak{H} into \mathfrak{R}
- $\mathfrak{H} \cong \mathfrak{R} :\Leftrightarrow \mathfrak{f} : \mathfrak{H} \cong \mathfrak{R}$ for some \mathfrak{f}
- $\mathfrak{H} \equiv \mathfrak{R} :\Leftrightarrow \mathfrak{H} \cong \mathfrak{R}$ and $\mathfrak{R} \cong \mathfrak{H}$

Clearly \cong is a preorder on Heyting Algebras with induced equivalence relation \equiv .

1.3.1 Example: Equivalent Heyting Algebras need not be isomorphic.

It is easily seen that any linear order with endpoints determines a Heyting Algebra. E.g. we find: $x \rightarrow y := \top$ if $x \leq y$, $x \rightarrow y := y$ if $y < x$. Moreover an embedding of linear orderings determines an embedding of Heyting Algebras. Consider the algebras given by the real interval $[0,1]$ and by $[0,1/2] \cup \{1\}$. On the one hand these algebras are equivalent, on the other they are not isomorphic. \circ

Let T be any consistent theory in constructive propositional logic or in constructive predicate logic. We take \mathfrak{H}_T to be the obvious Heyting Algebra given by the T -provable equivalence classes. Sometimes we will consider only equivalence classes of a subset X of the language of T , which is closed under the propositional connectives. In this case we write: $\mathfrak{H}_T(X)$.

We can go from theory to algebra. Obviously it is sometimes natural to go back and recover theories from algebras. We introduce some notions relevant to this motion, which is executed by choosing a set of generators.

- A *numbered* Heyting Algebra \mathbf{H} is a pair $\langle \mathfrak{f}, \mathfrak{H} \rangle$, where
 - (i) \mathfrak{f} is a function (not necessarily injective) from either $n = \{0, \dots, n-1\}$ or ω to $H_{\mathfrak{H}}$;

- (ii) \mathfrak{S} is generated by the range of \mathfrak{f} .
- A numbered Heyting Algebra is *finitely based* if $\text{dom}(\mathfrak{f})$ is finite.
- $\|\mathfrak{S}\| := \{ \langle \mathfrak{f}, \mathfrak{R} \rangle \mid \langle \mathfrak{f}, \mathfrak{R} \rangle \text{ is finitely based and } \mathfrak{R} \leq \mathfrak{S} \}$
- \mathfrak{L}_ν is the language of IPC if $\nu = \omega$, and the language of IPC restricted to p_0, \dots, p_{n-1} if $\nu = n$. We often write \mathfrak{L} for \mathfrak{L}_ω .
- For $A \in \mathfrak{L}_{\text{dom}(\mathfrak{f})}$: $\mathbf{H} \models A \Leftrightarrow \mathfrak{S} \models A[\mathfrak{f}]$, where $A[\mathfrak{f}]$ is the result of substituting $\mathfrak{f}i$ for p_i in A (for each relevant i). It is pleasant to use \models also when A contains p_j for $j \notin \text{dom}(\mathfrak{f})$. In this case we substitute \top for p_j .
- $\text{Th}(\mathbf{H}) := \{ A \in \mathfrak{L}_{\text{dom}(\mathfrak{f})} \mid \mathbf{H} \models A \}$.

1.3.2 Fact: Let $\mathbf{H} = \langle \mathfrak{f}, \mathfrak{S} \rangle$ be a numbered Heyting Algebra. Then \mathfrak{S} is isomorphic to $\mathfrak{S}_{\text{Th}(\mathbf{H})}$.

Proof: Trivial. □

Define:

- A numbered Heyting Algebra \mathbf{H} is RE if $\text{Th}(\mathbf{H})$ is RE.
- A Heyting Algebra is RE if it can be extended to an RE numbered Heyting Algebra.

Note that the Heyting Algebra of an RE theory is RE.

Let T be any theory and let \mathfrak{f} be a function from the propositional variables to the language of T . We write $A[\mathfrak{f}]$ for the result of substituting the $\mathfrak{f}(p_i)$ for p_i in A . Define:

- $A \models_T B \Leftrightarrow \forall \mathfrak{f} \ T \vdash A[\mathfrak{f}] \Rightarrow T \vdash B[\mathfrak{f}]$.

We say that the inference form A to B is an *IPC-derived rule for T* . Since all derived rules we will consider in this paper are IPC-derived we will suppress the ‘IPC’.

IPC-derived rules are studied in detail by V.V. Rybakov. A good reference is Rybakov[92], where it is shown that the IPC-derived rules for IPC are decidable.

1.3.3 Fact

- i) $A \models_T B \Leftrightarrow \forall \mathbf{H} \in \|\mathfrak{S}_T\| \ (\mathbf{H} \models A \Rightarrow \mathbf{H} \models B)$
- ii) $\|\mathfrak{S}_T\| \subseteq \|\mathfrak{S}_U\|$ and $A \models_U B \Rightarrow A \models_T B$.
- iii) $\mathfrak{S} \leq \mathfrak{R} \Rightarrow \|\mathfrak{S}\| \subseteq \|\mathfrak{R}\|$.

Proof: Obvious. \square

1.4 The density of Heyting Algebras of Arithmetical Theories

Evidently many properties of Heyting Algebras are not captured by embeddability results (see example 1.3.1). Such properties are not the main subject of this paper, yet they at least merit a brief comment here. Moreover many properties of the Boolean Algebra of Classical Arithmetical Theories can be generalized to the constructive case. We briefly illustrate this for the property of density.

Let $i\text{-Q}$ be the constructive version of Robinson's Arithmetic.

1.4.1 Fact: The Heyting Algebra of a consistent RE extension of $i\text{-Q}$ is dense, i.e. between every two points there is a third one.

Proof: Fix a consistent RE extension of $i\text{-Q}$, say T . Let \square stand for (the formalization of) provability in T . Consider any two sentences A and B such that $\square(A \rightarrow B)$ and not $\square(B \rightarrow A)$.

Interpolated Remark: The usual proof of this theorem for the classical case would be as follows. Take the Rosser sentence R of $T+B+\neg A$. I.e. something like:

$$T \vdash R \leftrightarrow \square((B \wedge \neg A) \rightarrow \neg R) \leq \square((B \wedge \neg A) \rightarrow R),$$

holds. Here \leq is the witness comparison relation, which is defined between formulas having an outer existential quantifier. There are two witness comparison relations, which are defined as follows:

- $(\exists x D_x \leq \exists y E_y) := \exists x (D_x \wedge \forall y < x \neg E_y)$,
- $(\exists x D_x < \exists y E_y) := \exists x (D_x \wedge \forall y \leq x \neg E_y)$.

The element between A and B will be: $C := (A \vee (B \wedge R))$. In constructive logic one cannot even conclude from the data that $T+B+\neg A$ is consistent. The correct constructive proof is just a slight variation on the classical argument. \circ End of Remark

Define by the fixed point theorem a sentence R such that (verifiably in T):

$$R \leftrightarrow \square((B \wedge R) \rightarrow A) \leq \square(B \rightarrow (A \vee R)).$$

Let $S := \square(B \rightarrow (A \vee R)) < \square((B \wedge R) \rightarrow A)$ and $C := (A \vee (B \wedge R))$. Clearly $\square(A \rightarrow C)$ and

$\Box(C \rightarrow B)$.

We have:

$$\begin{aligned}\Box(C \rightarrow A) &\rightarrow \Box((B \wedge R) \rightarrow A) \\ &\rightarrow R \vee S.\end{aligned}$$

On the other hand:

$$\begin{aligned}\Box((B \wedge R) \rightarrow A) \wedge R &\rightarrow \Box((B \wedge R) \rightarrow A) \wedge \Box R \\ &\rightarrow \Box(B \rightarrow A).\end{aligned}$$

And:

$$\begin{aligned}\Box((B \wedge R) \rightarrow A) \wedge S &\rightarrow \Box((B \wedge R) \rightarrow A) \wedge \Box(B \rightarrow (A \vee R)) \\ &\rightarrow \Box(B \rightarrow A).\end{aligned}$$

Combining we find: $\Box(C \rightarrow A) \rightarrow \Box(B \rightarrow A)$. Ergo not $\Box(C \rightarrow A)$.

Also we have:

$$\begin{aligned}\Box(B \rightarrow C) &\rightarrow \Box(B \rightarrow (A \vee R)) \\ &\rightarrow R \vee S.\end{aligned}$$

On the other hand:

$$\begin{aligned}\Box(B \rightarrow (A \vee R)) \wedge S &\rightarrow \Box(B \rightarrow (A \vee R)) \wedge \Box S \\ &\rightarrow \Box(B \rightarrow (A \vee R)) \wedge \Box \neg R \\ &\rightarrow \Box(B \rightarrow A).\end{aligned}$$

And:

$$\begin{aligned}\Box(B \rightarrow (A \vee R)) \wedge R &\rightarrow \Box(B \rightarrow (A \vee R)) \wedge \Box((B \wedge R) \rightarrow A) \\ &\rightarrow \Box(B \rightarrow A).\end{aligned}$$

Combining we find: $\Box(B \rightarrow C) \rightarrow \Box(B \rightarrow A)$. Ergo not $\Box(C \rightarrow A)$. \square

2 Embeddings of Heyting Algebras in Free Heyting Algebras

Every Heyting Algebra on countably many generators is the homomorphic image of \mathfrak{H}_{IPC} . In other words: it is the Heyting Algebra of some theory in IPC. On the other hand not every Heyting Algebra on countably many generators can be embedded into \mathfrak{H}_{IPC} . First of all \mathfrak{H}_{IPC} is prime, i.e.: $\mathfrak{H}_{IPC} \models (x \vee y) \Rightarrow (\mathfrak{H}_{IPC} \models x \text{ or } \mathfrak{H}_{IPC} \models y)$, or, in other words: IPC has the disjunction property. Clearly subalgebras inherit primeness. In this section we illustrate that many countable prime Heyting Algebras are not embeddable in \mathfrak{H}_{IPC} . We provide some information about the Heyting Algebras on finitely many generators that are embeddable in \mathfrak{H}_{IPC} . The problem of giving a neat characteri-

zation of the algebras embeddable in \mathfrak{S}_{IPC} is still open.

Whenever ‘ \vdash ’ is used without exhibiting a theory we intend provability in IPC.

2.1 Example: There are many non-trivial derived rules for IPC. For example:

- $\vdash (\neg\neg A \rightarrow A) \rightarrow (A \vee \neg A) \Rightarrow \vdash \neg\neg A \vee \neg A$ (De Jongh[82])
- $\vdash \neg A \rightarrow (B \vee C) \Rightarrow \vdash (\neg A \rightarrow B) \vee (\neg A \rightarrow C)$ (Independence of Premiss Rule)

This means that every embeddable algebra \mathfrak{S} will satisfy:

- $\mathfrak{S} \models (\neg\neg A \rightarrow A) \rightarrow (A \vee \neg A) \Rightarrow \mathfrak{S} \models \neg\neg A \vee \neg A$
- $\mathfrak{S} \models \neg A \rightarrow (B \vee C) \Rightarrow \mathfrak{S} \models (\neg A \rightarrow B) \vee (\neg A \rightarrow C)$.

2.2 Example: We give an infinitary derived rule. Let $F_n(p)$ be an enumeration of the formulas presenting the non-top elements of the Rieger Nishimura Lattice. (For information about this lattice, see e.g.: Troelstra & van Dalen[88a], p49.) We have:

- (For all $n \vdash F_n(A) \rightarrow B$) $\Rightarrow \vdash B$.

It follows that in an embedded Heyting Algebra for any x there can be no element between the $F_n(x)$ and the top.

Proof: Suppose for all $n \vdash F_n(A) \rightarrow B$. Let p be a propositional variable not in A and B . It follows that for all $n: \vdash F_n(p) \rightarrow ((p \leftrightarrow A) \rightarrow B)$. By Pitts[92] there is a uniform pre-interpolant of $((p \leftrightarrow A) \rightarrow B)$ w.r.t. to the variables in this formula unequal to p . This means that there is a formula C with just p free such that for any formula D containing no variables of A or B we have:

$$\vdash D \rightarrow ((p \leftrightarrow A) \rightarrow B) \Leftrightarrow \vdash D \rightarrow C.$$

(Following Pitts we could write the formula C as: $\forall \mathbf{q}((p \leftrightarrow A) \rightarrow B)$, where \mathbf{q} represents the propositional variables in A, B .) It follows that for every $n: \vdash F_n(p) \rightarrow C$. Ergo (since C only contains p): $\vdash C$ and hence $\vdash ((p \leftrightarrow A) \rightarrow B)$. Substituting A for p we find: $\vdash B$. □

2.3 Theorem: Every Heyting Algebra on finitely many generators that is embeddable in \mathfrak{S}_{IPC} is the Heyting Algebra of a finitely axiomatizable IPC theory.

Proof: Suppose the generators of the algebra go to A_1, \dots, A_n . We have:

$$\vdash B(A_1, \dots, A_n) \Leftrightarrow \vdash ((p_1 \leftrightarrow A_1) \wedge \dots \wedge (p_n \leftrightarrow A_n)) \rightarrow B(p_1, \dots, p_n).$$

We suppose that $\{p_1, \dots, p_n\} \cap \text{VAR}(A_i) = \emptyset$ and $\text{VAR}(B) \subseteq \{p_1, \dots, p_n\}$. Now let C be the Pittsean post-interpolant of $((p_1 \leftrightarrow A_1) \wedge \dots \wedge (p_n \leftrightarrow A_n))$ w.r.t. the variables in the A_i . So, if these variables are \mathbf{q} , we could write C as: $\exists \mathbf{q}((p_1 \leftrightarrow A_1) \wedge \dots \wedge (p_n \leftrightarrow A_n))$. The only variables of C are the p_i and we have: $\vdash B(A_1, \dots, A_n) \Leftrightarrow \vdash C \rightarrow B$. \square

Which formulas C are axioms of Heyting Algebras on finitely many generators that are embeddable in $\mathfrak{S}_{\text{IPC}}$? We call such C *IPC-exactly provable*. In this paper we will abbreviate *IPC-exactly provable* by *exact*. So $C(p_1, \dots, p_n)$ is exact if there are A_1, \dots, A_n such that for all $B(p_1, \dots, p_n)$: $\vdash B(A_1, \dots, A_n) \Leftrightarrow \vdash C \rightarrow B$. The notion of exactly provable formula was introduced in De Jongh[82].

Clearly by the above the exactly provable formulas are precisely those which are provably equivalent to Pitts' formulas of the form $\exists \mathbf{q}((p_1 \leftrightarrow A_1) \wedge \dots \wedge (p_n \leftrightarrow A_n))$, where \mathbf{q} contains precisely the variables occurring in the A_i and where none of the p_j is in \mathbf{q} .

We say that A is *prime* if $\mathfrak{S}_{\text{IPC}+A}$ is prime, i.e. $\text{IPC}+A$ is consistent and $\text{IPC}+A$ has the disjunction property:

- for all $B, C \in \mathfrak{L}$ $\vdash A \rightarrow (B \vee C) \Rightarrow \vdash A \rightarrow B$ or $\vdash A \rightarrow C$.

In an alternative formulation, adhering to the convention that the empty disjunction is \perp , IPC is prime if for every finite set of formulas X :

$$\vdash A \rightarrow \bigvee X \Rightarrow \exists B \in X \vdash A \rightarrow B.$$

We give some properties of exact formulas and provide some special classes of such formulas. Our primary aim is to show that the prime NNIL-formulas are all exact. *NNIL-formulas* are formulas with No Nestings of Implications to the Left. Let's define NNIL more precisely. Let $\text{Sub}(A)$ be the set of subformulas of A . We have:

- A is in NNIL iff for all $(B \rightarrow C) \in \text{Sub}(A)$: B does not contain \rightarrow .

NNIL-formulas were studied in Visser[85] and Renardel[86]. The lemmas we give, can, however, also be used to establish exactness for more formulas than our target class. The result on NNIL will be used in the proof of 6.2.

2.4 Observation: If A is exact, then A is prime.

2.5 Observation: Suppose p does not occur in A . Then $(p \rightarrow A) \wedge B$ is exact if $B[p:=p \wedge A]$ is.

Proof: Let \mathfrak{f} be the embedding for $B[p:=p \wedge A]$. We show that $g := [p:=p \wedge A] \circ \mathfrak{f}$ is the embedding for $(p \rightarrow A) \wedge B$. We have for C with variables from p, A, B :

$$\begin{aligned} \vdash Cg &\Leftrightarrow \vdash C[p:=p \wedge A]\mathfrak{f} \\ &\Leftrightarrow \vdash B[p:=p \wedge A] \rightarrow C[p:=p \wedge A] \\ &\Leftrightarrow \vdash ((p \rightarrow A) \wedge B) \rightarrow C. \end{aligned}$$

(The last equivalence from left to right is because $\vdash (p \rightarrow A) \Leftrightarrow (p \Leftrightarrow (p \wedge A))$. From right to left is by substituting $p \wedge A$ for p .) \square

We say that a formula is *confined* if it is a conjunction of formulas of the form $p \rightarrow B$. A formula is *strictly confined* if it is confined and if for any two distinct conjuncts the antecedent variables are different. (We consider \top as the empty conjunction, so \top is strictly confined).

2.6 Corollary: Any confined formula is exact.

Proof: Suppose A is confined. First rewrite A to a strictly confined formula A' by merging different conjuncts $p \rightarrow B$ and $p \rightarrow C$ to $p \rightarrow (B \wedge C)$. Suppose A' is of the form $(p \rightarrow D) \wedge E$. This formula is equivalent to $A'' := ((p \rightarrow D[p:=\top]) \wedge E)$. According to observation 3 A'' is exact if $A^* := E[p:=(p \wedge D[p:=\top])]$ is. Clearly A^* is again a strictly confined formula with less conjuncts than A' . Repeat the procedure till all conjuncts are eliminated and we end up with \top . \top is exact by the identity substitution. \square

Note that it follows that confined formulas are prime.

2.7 Observation Suppose p does not occur in A . Then $(p \wedge A)$ is exact if A is.

Proof: Suppose \mathfrak{f} is the embedding for A . Take $g := [p:=\top] \circ \mathfrak{f}$. Then g is the embedding for $(p \wedge A)$. Let the variables of B be among the variables of $(p \wedge A)$, we have:

$$\begin{aligned} \vdash Bg &\Leftrightarrow \vdash B[p:=\top]\mathfrak{f} \\ &\Leftrightarrow \vdash A \rightarrow B[p:=\top] \\ &\Leftrightarrow \vdash (p \wedge A) \rightarrow B. \end{aligned} \quad \square$$

2.8 Theorem: Every prime NNIL-formula is exact.

2.8.1 Lemma: Suppose p does not occur in A . Then A is prime if $(p \wedge A)$ is.

Proof: Suppose $(p \wedge A)$ is prime. Let X be a finite set of formulas and suppose $\vdash A \rightarrow \bigvee X$. Without loss of generality we may assume that p does not occur in X . It follows that $\vdash (p \wedge A) \rightarrow \bigvee X$ and hence $\vdash (p \wedge A) \rightarrow B$ for some $B \in X$. By substituting \top for p we find: $\vdash A \rightarrow B$. \square

Proof of 2.8: Let A be a NNIL-formula. We will reduce A to a formula A_0 . The formula A_0 satisfies one of the following properties: (i) A_0 is confined or (ii) A_0 is a prime NNIL-formula and has strictly less propositional variables than A . Moreover we have: if A_0 is exact, then A is exact. In the first case we are done, in the second case we repeat the procedure.

Step 1: We first remove \top and \perp from A by the obvious procedure. This only fails when we end up with either \top or \perp . We cannot end up with \perp , since A was supposed to be prime and hence non-refutable. If we end up with \top , then A is exact by the identity substitution. If we do not end up with \top go on to step 2.

Step 2: Write A in disjunctive normal form (treating the implications as atoms). Since A is prime, it is equivalent with one of its disjuncts, say A' . A' is a conjunction of atoms and implications. If the number of atoms is zero go on to step 3. Otherwise write A' in the form $p \wedge C$. Clearly $p \wedge C$ is equivalent to $p \wedge (C[p:=\top])$. Put $A_0 := C[p:=\top]$.

Step 3: A' is a conjunction of implications. Reduce subformulas of the form $(B \wedge C) \rightarrow D$ to $(B \rightarrow (C \rightarrow D))$ and subformulas of the form $(B \vee C) \rightarrow D$ to $(B \rightarrow D) \wedge (C \rightarrow D)$. Repeat the procedure till no such subformulas are left. Let A_0 be the result. Since A' was in NNIL, clearly A_0 is confined. \square

3 Some useful facts about IPC and HA

In this section we provide some technical preliminaries to the result of section 5.

We suppose the reader is familiar with Kripke models for IPC (see Troelstra & van Dalen[88a], or Smorynski[73]). To fix notations: a *Kripke model* is a structure $\mathbb{K} = \langle K, \leq, \Vdash \rangle$, where K is a non-empty set of nodes, \leq is a partial ordering, \Vdash is the atomic forcing relation: it is a relation between nodes and propositional atoms, satisfying: $k \leq k'$ and $k \Vdash p \Rightarrow k' \Vdash p$. The relation \Vdash can be extended to the full language of IPC in the standard way. We write $\mathbb{K} \Vdash A$ for: $\forall k \in K \ k \Vdash A$. A *rooted Kripke model* \mathbb{K} is a structure $\langle K, k_0, \leq, \Vdash \rangle$, where $\langle K, \leq, \Vdash \rangle$ is a Kripke model and where $k_0 \in K$ is the bottom element w.r.t. \leq . For any $k \in K$ $\mathbb{K}[k]$ is the model $\langle K', k, \leq', \Vdash' \rangle$, where $K' := \{k' \mid k \leq k'\}$ and where \leq' and \Vdash' are the restrictions of \leq respectively \Vdash to K' . (We will often simply write \leq and \Vdash for \leq' and \Vdash' .)

3.1 The Henkin construction: A set X is *adequate* if it is finite, closed under subformulas and contains \perp . A set Γ is X -saturated if:

- (i) $\Gamma \subseteq X$, (ii) $\Gamma \not\vdash \perp$, (iii) $\Gamma \vdash A, A \in X \Rightarrow A \in \Gamma$,
- (iv) $\Gamma \vdash (B \vee C), (B \vee C) \in X \Rightarrow B \in \Gamma$ or $C \in \Gamma$.

The Henkin model \mathbb{H}_X has as nodes the X -saturated sets and as accessibility relation \sqsubseteq . The atomic forcing in the nodes is given by: $\Gamma \Vdash p \Leftrightarrow p \in \Gamma$. We have by a standard argument: for $A \in X$: $\Gamma \Vdash A \Leftrightarrow A \in \Gamma$.

3.2 Definitions

i) Let \mathbf{K} be a set of Kripke models. $M(\mathbf{K})$ is the model with nodes $\langle k, \mathbb{K} \rangle$ for $k \in \mathbb{K} \in \mathbf{K}$ and ordering: $\langle k, \mathbb{K} \rangle \leq \langle m, \mathbb{M} \rangle \Leftrightarrow \mathbb{K} = \mathbb{M}$ and $k \leq_{\mathbb{K}} m$. As atomic forcing we take:

- $\langle k, \mathbb{K} \rangle \Vdash p \Leftrightarrow k \Vdash_{\mathbb{K}} p$.

(In practice we will forget the second components of the new nodes, pretending the domains to be disjoint already.)

ii) Let \mathbb{K} be a Kripke model. $B(\mathbb{K})$ is the rooted model obtained by adding a new bottom b to \mathbb{K} and by taking: $b \Vdash p \Leftrightarrow \mathbb{K} \Vdash p$. We write $\text{Glue}(\mathbf{K}) := BM(\mathbf{K})$.

3.3 Push Down Lemma: Let X be adequate. Suppose Δ is X -saturated and $\mathbb{K} \Vdash \Delta$. Then $\text{Glue}(\mathbb{H}_X[\Delta], \mathbb{K}) \Vdash \Delta$.

Proof: We show by induction on $A \in X$ that $b \models A \Leftrightarrow A \in \Delta$. The cases of atoms, conjunction and disjunction are trivial. If $(B \rightarrow C) \in X$ and $b \models (B \rightarrow C)$, then $\Delta \models (B \rightarrow C)$ and hence $(B \rightarrow C) \in \Delta$. Conversely suppose $(B \rightarrow C) \in \Delta$. If $b \not\models B$, we are easily done. If $b \models B$, then $B \in \Delta$, hence $C \in \Delta$ and by the Induction Hypothesis: $b \models C$. \square

We say that Δ is prime if it is consistent and:

for every $(C \vee D) \in \mathcal{L}$: $\Delta \vdash (C \vee D) \Rightarrow \Delta \vdash C$ or $\Delta \vdash D$.

3.4 Theorem: Suppose X is adequate and Δ is X -saturated. then Δ is prime.

Proof: Δ is consistent by definition. Suppose $\Delta \vdash C \vee D$ and $\Delta \not\vdash C$ and $\Delta \not\vdash D$. Suppose $\mathbb{K} \models \Delta$, $\mathbb{K} \not\models C$, $\mathbb{M} \models \Delta$ and $\mathbb{M} \not\models D$. Consider $\text{Glue}(\mathbb{H}_X(\Delta), \mathbb{K}, \mathbb{M})$. By 3.3 we have: $b \models \Delta$. On the other hand by persistence: $b \not\models C$ and $b \not\models D$. Contradiction. \square

3.5 A big model: Construct a Henkin model by taking as nodes $\langle \Gamma, X \rangle$, where X is adequate and Γ is X -saturated. Take $\langle \Gamma, X \rangle \leq \langle \Delta, Y \rangle :\Leftrightarrow \Gamma \subseteq \Delta$ and $X \subseteq Y$. Also: $\langle \Gamma, X \rangle \models p :\Leftrightarrow p \in \Gamma$. Then for all $A \in \mathcal{L}$: $\langle \Gamma, X \rangle \models A \Leftrightarrow \Gamma \vdash A$. The proof, which uses 3.4, is left to the industrious reader. \circ

3.6 Formalization in HA: We first formalize Kripke completeness for finite models in Peano Arithmetic (PA). Noting that the model existence theorem yields a multi-exponential bound E on the size of the Henkin model we formulate the result as follows: $\text{PA} \vdash \forall A((\forall \mathbb{K} \leq E(A) \mathbb{K} \models A) \rightarrow \text{IPC} \vdash A)$. Noting that the formula proved is Π_2 , we see that by a theorem due to Kreisel: $\text{HA} \vdash \forall A((\forall \mathbb{K} \leq E(A) \mathbb{K} \models A) \rightarrow \text{IPC} \vdash A)$. Since the converse is readily verifiable in HA we find:

$$\text{HA} \vdash \forall A((\forall \mathbb{K} \leq E(A) \mathbb{K} \models A) \leftrightarrow \text{IPC} \vdash A).$$

So IPC-provability is decidable in HA.

In intuitionistic theories even subsets of the singleton set are not decidable. We, however, assume that the finite sets that we are using, e.g. in the construction of the Henkin model, are *coded as numbers* and hence provably finite and decidable. Under this convention whether a finite set is X -saturated or not becomes decidable, given the decidability of IPC-provability.

We leave it to the reader to verify 3.3 and 3.4 in HA (assuming \mathbf{K} to be a finite set of finite models, etc.). Note that the reductio reasoning in 3.4 is harmless because of decidability. \circ

3.7 Theorem: Let X be a prime, RE set of IPC-formulas, closed under IPC-consequence. Without loss of generality we may assume that X is given by a recursive increasing sequence of finite approximations X_i . We assume that $X_0 = \emptyset$. Say, this sequence is presented by the Δ_1 -formula $\xi(i, x)$. Then we can represent X by a sequence $\langle Y_i, Z_i \rangle$, where:

- i) $X = \bigcup Y_i$,
- ii) $i < j \Rightarrow (Y_i \subseteq Y_j \text{ and } Z_i \subseteq Z_j)$
- iii) Z_i is adequate
- iv) Y_i is Z_i -saturated

Our sequence can be represented by a Δ_1 -formula $\sigma(i, y, z)$ such that HA verifies the functionality of the sequence, plus (ii), (iii), (iv). Let Y be given as: $\{B \mid \exists i, y, z (\sigma(i, y, z) \wedge B \in y)\}$. It follows by 3.4-3.6 that HA verifies that:

Y is prime, that $Y \subseteq X$ and that $(X \text{ is prime}) \rightarrow Y = X$.

Proof: We reason informally, but constructively, and leave verifiability in HA to the industrious reader. Remember that our sets are really finite, decidable sets represented by numbers. Fix an increasing sequence U_i of adequate sets such that for every A we can effectively find an i such that $\text{Sub}(A) \subseteq U_i$. We define weakly monotonic functions $f, g: \omega \rightarrow \omega$ and take $Z_i := U_{f_i}$ and $Y_i := \{B \in Z_i \mid X_{g_i} \vdash B\}$.

- $f_0 := 0, g_0 := 0$
- Consider $U_{f_{n+1}}$. In case $\{B \in U_{f_{n+1}} \mid X_{g_n} \vdash B\}$ is $U_{f_{n+1}}$ -saturated, put $f(n+1) := f_n + 1, g(n+1) := n + 1$. Otherwise $f(n+1) := f_n, g(n+1) := g_n$.

f and g are recursive, since, by 3.6, IPC-provability is (verifiably) decidable. By the formalization of 3.4, every Y_i is prime (in case $i=0$, this uses the fact that IPC is prime).

Suppose X is prime, then both f and g tend to infinity (and hence $Y = X$). Evidently it is sufficient to show that after every stage the first clause of the definition of f and g will become active. Consider stage n . If at stage $n+1$ $\{B \in U_{f_{n+1}} \mid X_{g_n} \vdash B\}$ is $U_{f_{n+1}}$ -

saturated, we are done. Suppose not. Since U_{fn+1} is finite we can exhaustively enumerate the U_{fn+1} -disjunctions $E \vee F$ proved by X_{n+1} . Since X is prime we can find for any such $E \vee F$ an i such that $X_i \vdash E$ or $X_i \vdash F$. Let n_1 be the maximum of these i 's. By assumption $\{B \in U_{fn+1} \mid X_{n_1} \vdash B\}$ properly extends $\{B \in U_{fn+1} \mid X_{n+1} \vdash B\}$. If $\{B \in U_{fn+1} \mid X_{n_1} \vdash B\}$ is U_{fn+1} -saturated then either f and g have moved between $n+1$ and n_1 or will move at n_1 . If not we repeat the procedure. Thus we obtain a sequence $n+1 := n_0 < n_1 < \dots < n_N < \dots$. Note that $\{B \in U_{fn+1} \mid X_{n_{i+1}} \vdash B\}$ properly extends $\{B \in U_{fn+1} \mid X_{n_i} \vdash B\}$ so $N < |U_{fn+1}|$. So eventually we reach a U_{fn+1} -saturated stage and our functions did move or move now. \square

4 What is HA^* ?

In this section we introduce the theory HA^* . This theory was introduced in Visser[82]. HA^* is to Beeson's fp-realizability (Beeson[75]) as Troelstra's $HA+ECT_0$ is to Kleene's r-realizability. This means that for a suitable variant of fp-realizability HA^* is the set of sentences such that their fp-translations are provable in HA . The natural way to define HA^* is by a fixed point construction as: HA plus the *Completeness Principle for HA^** . (Here it is essential that the construction is verifiable in HA , see below.) The Completeness Principle can be viewed as an arithmetically interpreted modal principle. The Completeness Principle viewed modally is:

$$C \quad \vdash A \rightarrow \Box A$$

The Completeness Principle for a specific theory T is:

$$C[T] \quad \vdash A \rightarrow \Box_T A.$$

Here \Box_T stands for the formalization of provability in T . So we have:

$$HA^* = HA + C[HA^*].$$

We briefly review some of the results of Visser[82].

- Let \mathfrak{A} be the smallest class closed under atoms and all connectives except implication, satisfying: $A \in \Sigma_1$ and $B \in \mathfrak{A} \Rightarrow (A \rightarrow B) \in \mathfrak{A}$. Note that modulo provable equivalence in HA all formulas of the classical arithmetical hierarchy in their standard form are in \mathfrak{A} . HA^* is conservative w.r.t. \mathfrak{A} over HA .
- There are infinitely many incomparable T with $T=HA+C_T$. However if $T=HA+C_T$ verifiably in HA , then $T=HA^*$.
- Let KLS:=Kreisel-Lacombe-Shoenfield's Theorem on the continuity of the effective

operations. We have: $HA^* \vdash KLS \rightarrow \Box_{HA^*} \perp$. This immediately gives Beeson's result that $HA \not\vdash KLS$ (see Beeson[75]).

Consider the Löb conditions.

- L1 $\vdash A \Rightarrow \vdash \Box A$
- L2 $\vdash \Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)$
- L3 $\vdash \Box A \rightarrow \Box \Box A$
- L4 $\vdash \Box(\Box A \rightarrow A) \rightarrow \Box A$

i-K is given by IPC+L1,L2. i-L is i-K+L3,L4. We write i-K{P} for the extension of i-K with some principle P. Note that i-L{C} is valid for provability interpretations in HA^* .

A principle closely connected to C is the Strong Löb Principle:

$$SL \quad \vdash (\Box A \rightarrow A) \rightarrow A$$

As a special case of SL we have: $\vdash \neg \Box \perp$.

4.2 Fact: i-L{C} is interderivable with i-K{SL}.

Proof: L4 is immediate from SL. "i-K{SL} \vdash C":

$$\begin{aligned} \vdash A &\rightarrow (\Box(A \wedge \Box A) \rightarrow (A \wedge \Box A)) \\ &\rightarrow A \wedge \Box A \\ &\rightarrow \Box A. \end{aligned}$$

"i-L{C} \vdash SL":

$$\begin{aligned} \vdash (\Box A \rightarrow A) &\rightarrow (\Box A \rightarrow A) \wedge \Box(\Box A \rightarrow A) \\ &\rightarrow (\Box A \rightarrow A) \wedge \Box A \\ &\rightarrow A. \end{aligned} \quad \square$$

i-L{C} is a kind of Kindergarten Theory in which all the well-known syntactical results of Provability Logic have extremely simple versions. We add the proofs for completeness. 4.3-4.6 are not essential for the rest of the paper.

4.3 Substitution Lemma: In $i-L\{C\}$ we have a very powerful substitution principle:

$$S^{++} \vdash (A \leftrightarrow B) \rightarrow (CA \leftrightarrow CB)$$

Proof: By a simple induction on C . □

We say that p occurs only modalized in A if all occurrences of p are in the scope of \Box .

4.4 Uniqueness of Fixed Points in $i-L\{C\}$: Suppose p occurs only modalized in Ap and q does not occur in Ap . We have in $i-L\{C\}$:

$$\begin{aligned} \vdash (p \leftrightarrow Ap) \wedge (q \leftrightarrow Aq) &\rightarrow (\Box(p \leftrightarrow q) \rightarrow (Ap \leftrightarrow Aq)) \\ &\rightarrow (p \leftrightarrow q) \\ &\rightarrow (p \leftrightarrow q). \end{aligned} \quad \square$$

4.5 Explicit Fixed Points in $i-L\{C\}$: Suppose p occurs only modalized in Ap . We show that Ap has fixed point $A\top$. We have:

$$\begin{aligned} \vdash A\top &\rightarrow (A\top \leftrightarrow \top) \\ &\rightarrow AA\top. \\ \vdash AA\top &\rightarrow (\Box A\top \rightarrow \Box(A\top \leftrightarrow \top)) \\ &\rightarrow A\top) \\ &\rightarrow A\top. \end{aligned} \quad \square$$

A formula of the modal language is closed if it contains no propositional variables. We define: $\Box^0 \perp := \perp$, $\Box^{n+1} \perp := \Box \Box^n \perp$, $\Box^\omega \perp := \top$.

4.6 The Closed Fragment of $i-L\{C\}$: Every closed formula A is $i-L\{C\}$ -provably equivalent to a formula of the form $\Box^\alpha \perp$ for $\alpha \leq \omega$.

Proof: The proof is by induction on A . We have:

$$\begin{aligned} \vdash \top &\leftrightarrow \Box^\omega \perp, \vdash \perp \leftrightarrow \Box^0 \perp \\ \vdash \Box^\alpha \perp \wedge \Box^\beta \perp &\leftrightarrow \Box^{\min(\alpha, \beta)} \perp \\ \vdash \Box^\alpha \perp \vee \Box^\beta \perp &\leftrightarrow \Box^{\max(\alpha, \beta)} \perp \\ \vdash \Box^\alpha \perp \rightarrow \Box^\beta \perp &\leftrightarrow \Box^{\alpha \rightarrow \beta} \perp, \text{ where } (\alpha \rightarrow \beta) := \top \text{ if } \alpha \leq \beta \text{ and } (\alpha \rightarrow \beta) := \beta \text{ if } \beta < \alpha. \\ \text{Note that } \min(\alpha, \gamma) \leq \beta &\Leftrightarrow \gamma \leq (\alpha \rightarrow \beta). \end{aligned}$$

$$\vdash \Box\Box^\alpha \perp \leftrightarrow \Box^{1+\alpha} \perp \quad \square$$

4.7 Open Problems

- i) Is $i\text{-L}\{C\}$ the provability logic of HA^* ?
- ii) Prove or refute: $HA \vdash \neg\neg KLS$.

5 A Shavrukov Style Embedding Theorem for HA^*

Shavrukov proved that every RE Diagonalizable Algebra satisfying an appropriate Disjunction Property is embeddable in the Diagonalizable Algebra of Peano Arithmetic. It is clear from section 3 that there is no analogous result for Heyting Algebras and Heyting Arithmetic. In this section we show that an analogue can be obtained for the theory HA^* .

5.1 Theorem: Any RE prime Heyting Algebra \mathfrak{H} can be embedded in \mathfrak{H}_{HA^*} . Moreover the equivalence classes in the range of the embedding all contain a Σ_1 -sentence.

Before proving the theorem we briefly look at an illustrative example to give the reader some feeling of how it is possible that an embedded algebra can completely consist of equivalence classes of Σ_1 -sentences.

5.1.1 Example: Consider the algebra \mathfrak{H} , IPC-axiomatized by $\neg\neg p \rightarrow p$. To be precise: $\mathfrak{H} = \mathfrak{H}_{IPC+(\neg\neg p \rightarrow p)}(\mathcal{L}_1)$. We have:

\mathfrak{H} can be embedded into \mathfrak{H}_{IPC} by e.g. $[p := \neg p]$;

\mathfrak{H} can be embedded into \mathfrak{H}_{HA} by e.g. $[p := \neg\Box_{HA} \perp]$.

On the other hand \mathfrak{H} cannot be embedded into \mathfrak{H}_{HA} by sending p to a Σ_1 -sentence, since for any Σ_1 -sentence B , we have:

$$\begin{aligned} HA \vdash \neg\neg B \rightarrow B &\Rightarrow HA \vdash B \vee \neg B \\ &\Rightarrow HA \vdash B \text{ or } HA \vdash \neg B. \end{aligned}$$

(The first implication is proved by applying the Friedman translation for $\neg B$ to $(\neg\neg B \rightarrow B)$. See e.g. Visser[85].)

We turn to HA^* . Let R be the ordinary Σ_1 Rosser sentence for HA^* . I.e.:

$$R \leftrightarrow \Box_{HA^*} \neg R \leq \Box_{HA^*} R.$$

Let $S := \Box_{HA^*}R < \Box_{HA^*}\neg R$. We have by the ordinary Rosser property:

$$HA^* \not\vdash R \text{ and } HA^* \not\vdash S.$$

On the other hand we have:

$$HA^* \vdash \neg R \leftrightarrow S \text{ and } HA^* \vdash \neg S \leftrightarrow R.$$

We prove the first equivalence. “ \leftarrow ” Trivially $HA^* \vdash S \rightarrow \neg R$. “ \rightarrow ” Reason in HA^* . Suppose $\neg R$ and $\Box_{HA^*}S$. It follows from the second assumption that $\Box_{HA^*}\neg R$ and hence that $R \vee S$. Combining $R \vee S$ with our first assumption, we get: S . By SL we may drop the assumption $\Box_{HA^*}S$.

Using the above facts it is easy to see that the subalgebra of \mathfrak{S}_{HA^*} generated by R is given by the non-equivalent Σ_1 -sentences: $\perp, R, S, \Box_{HA^*}\perp, \top$. This algebra is clearly isomorphic to \mathfrak{S} . ○

In 5.6.1 we show that it is definitely *not* the case in HA^* , that the Σ_1 -sentences are closed under the Boolean operations (modulo provable equivalence).

Proof of 5.1: Let \Box stand for \Box_{HA^*} and Proof for Proof_{HA^*} .

Consider the following Kripke model \mathbb{H} , which is a variant of the Big Model of 3.5. Its nodes are of the form $\langle i, U, V \rangle$, where:

- $i \in \{0, 1\}$
- V is an adequate set of formulas,
- $U \subseteq V$, U is V -saturated.

Define \preceq and \models as follows:

- $\langle i, U, V \rangle \preceq \langle j, W, T \rangle : \Leftrightarrow i \leq j \text{ and } U \subseteq W \text{ and } V \subseteq T \text{ and } (i=1 \Rightarrow V=T)$
- $\langle i, U, V \rangle \models p : \Leftrightarrow p \in U$.

Using 3.3 and 3.6 it is easy to see (in HA) that:

$$\text{for any formula } A: \langle 0, U, V \rangle \models A \Leftrightarrow U \vdash A,$$

$$\text{for } A \in V: \langle 1, U, V \rangle \models A \Leftrightarrow U \vdash A.$$

Note that it follows that the relation $k \models A$ is decidable.

Let $\langle Y_i, Z_i \rangle$ be an enumeration of a propositional theory presenting \mathfrak{S} , satisfying the properties promised in 3.7. We define a Solovay function \mathfrak{h} from ω to the nodes of \mathbb{H} .

ξx , the *state* of \mathfrak{h} at x , is defined as $(\mathfrak{h}x)_0$. $\xi 0$ will be set at 0. Till a certain catastrophic Event happens, the state will remain 0 and \mathfrak{h} will run upward through nodes $\langle 0, Y_i, Z_i \rangle$. As soon as (and *if*) the Event happens, the state will definitively move to 1 and our function runs upwards through nodes of the form $\langle 1, U, V \rangle$. Define by the Recursion Theorem \mathfrak{h} as follows:

- $[A] :\leftrightarrow \exists x \mathfrak{h}x \neq A$
- $\mathfrak{h}0 := \langle 0, Y_0, Z_0 \rangle$
- $\mathfrak{h}(n+1) := k$ if (*) $\text{Proof}(n, [A])$, $\mathfrak{h}n \neq A$,
 k is a 1-node, $\mathfrak{h}n \leq k$, k maximal such that $k \neq A$
- $\mathfrak{h}(n+1) := \langle 0, Y_{n+1}, Z_{n+1} \rangle$ if case (*) does not obtain and $\xi n = 0$
- $\mathfrak{h}(n+1) := \mathfrak{h}n$ if case (*) does not obtain and $\xi n = 1$.

Since \models is (provably in HA) decidable, it follows that \mathfrak{h} is a well defined recursive function.

Note that the catastrophic Event is the first time that (*) obtains. Before the Event the function enumerates nodes representing better and better approximations of ξ . After the event it behaves like an ordinary Solovay function traveling upwards through a converse wellfounded (w.r.t. $<$) part of the model.

5.2 Lemma

$$\text{HA} \vdash x \leq y \rightarrow \mathfrak{h}x \leq \mathfrak{h}y$$

$$\text{HA} \vdash (x \leq y \wedge \mathfrak{h}x \neq A) \rightarrow \mathfrak{h}y \neq A$$

Proof: Obvious. □

5.3 Lemma: $\text{HA} \vdash \xi x \neq 0 \rightarrow \Box \exists y \mathfrak{h}x < \mathfrak{h}y$.

Proof: Reason in HA. Suppose $\xi x \neq 0$. \mathfrak{h} must have arrived at $\mathfrak{h}x$ by case (*). So for some A and for some $p < x$: $\text{Proof}_{\text{HA}}^*(p, [A])$, $\mathfrak{h}(p+1) = \mathfrak{h}x \neq A$. By Σ -completeness we have: $\Box \mathfrak{h}x \neq A$. Combining this with $\Box \exists y \mathfrak{h}y \neq A$, we obtain, using 5.2, the desired result. □

5.4 Lemma: $\xi n = 0$ for any n .

Proof: Suppose $\varepsilon_n \neq 0$. By 5.3: $\Box \exists y \ \check{h}_n < \check{h}_y$. Remember that HA^* is Π_2 -conservative over HA . Thus HA^* will certainly satisfy Σ -reflection. It follows that for some m : $\check{h}_n < \check{h}_m$. Repeating the argument we can construct an infinite strictly ascending chain above \check{h}_n . This contradicts $\varepsilon_n \neq 0$. \square

5.5 Lemma: $[\cdot]$ commutes modulo HA^* -provability with the propositional connectives.

Proof: Reason in HA^* . Clearly $[\perp] \leftrightarrow \perp$ and $[\top] \leftrightarrow \top$.

Suppose $[A \wedge B]$, then for some x : $\check{h}_x \models A \wedge B$. It follows that $\check{h}_x \models A$ and $\check{h}_x \models B$ and hence $[A] \wedge [B]$. Conversely suppose $[A] \wedge [B]$. Say $\check{h}_y \models A$ and $\check{h}_z \models B$. Let $u := \max(y, z)$, then by 5.2: $\check{h}_u \models A$ and $\check{h}_u \models B$ and thus $\check{h}_u \models A \wedge B$. We may conclude: $[A \wedge B]$.

Suppose $[A \vee B]$, then for some x : $\check{h}_x \models A \vee B$. It follows that $\check{h}_x \models A$ or $\check{h}_x \models B$ and hence $[A] \vee [B]$. Conversely suppose $[A] \vee [B]$. Suppose e.g. $\check{h}_y \models A$. It is immediate that also $\check{h}_y \models A \vee B$ and so $[A \vee B]$. Similarly in case $\check{h}_z \models B$.

Suppose $[A \rightarrow B]$ and $[A]$. Then for some x and y : $\check{h}_x \models A \rightarrow B$ and $\check{h}_y \models A$. Take $u := \max(x, y)$. Clearly $\check{h}_u \models A \rightarrow B$ and $\check{h}_u \models A$. Ergo: $\check{h}_u \models B$ and thus $[B]$. Conversely suppose $[A] \rightarrow [B]$. We show $[A \rightarrow B]$ using the SL. So we may also assume $\Box [A \rightarrow B]$. Suppose $\text{Proof}(p, [A \rightarrow B])$. In case $\check{h}(p) \models (A \rightarrow B)$ we have $[A \rightarrow B]$. Suppose $\check{h}(p) \not\models (A \rightarrow B)$. In this case $\check{h}(p+1)$ is a maximal $k \geq \check{h}_p$ such that $k \not\models (A \rightarrow B)$. It follows that $k \models A$ and $k \not\models B$. From $\check{h}(p+1) = k \models A$, we have: $[A]$, and hence by assumption: $[B]$. But $[B]$ immediately implies: $[A \rightarrow B]$. So in both cases we find $[A \rightarrow B]$. By the SL we may drop the assumption that $\Box [A \rightarrow B]$. \square

We finish our proof of 5.1, by showing that: $A \in X \Leftrightarrow HA^* \vdash [A]$.

Suppose $A \in X$. Then for some n : $A \in Y_n$. By 5.4: $\varepsilon_n = 0$ and hence $\check{h}_n = \langle 0, Y_n, Z_n \rangle$. Ergo $\check{h}_n \models A$ and so $HA^* \vdash \check{h}_n \models A$ and thus $HA^* \vdash [A]$. Conversely suppose $HA^* \vdash [A]$. Say m codes a proof of $[A]$. Suppose $Y_m \not\models A$. Since $\check{h}_m = \langle 0, Y_m, Z_m \rangle$ it follows that $\check{h}_m \not\models A$. So clause (*) would become active and the catastrophic Event would take place. But 5.4

tells us this cannot happen at a standard stage. \square

5.6 Remarks on the proof: (i) The present proof combines the proof strategy from Zambella[92] with an idea from Visser[85] (on how to handle implication using the SL). In fact our proof follows Zambella's quite closely modulo some inessential stylistic differences (like our use of a kind of Henkin model).

ii) The proof cannot be extended in any obvious way to give a completeness theorem for the provability logic of HA^* , since nodes of our Henkin model where we have $\Box \perp$ also satisfy Excluded Third. But of course HA^* does not prove Excluded Third from $\Box \perp$.

iii) An attractive alternative formulation of the proof is to take, on the one hand, as nodes of the Henkin model the more traditional pairs $\langle U, V \rangle$, but to work, on the other hand, with two accessibility relations:

- $\langle U, V \rangle \leq_0 \langle W, T \rangle \quad :\Leftrightarrow U \subseteq W \text{ and } V \subseteq T$
- $\langle U, V \rangle \leq_1 \langle W, T \rangle \quad :\Leftrightarrow U \subseteq W \text{ and } V = T$

Corresponding to these different accessibility relations we have forcing relations \models_0 and \models_1 . We define a suitably adapted Solovay function simultaneously with an auxiliary state function. Which accessibility relation and which forcing relation is relevant, will depend on the state. We leave it to the reader to work out more details.

iv) The $[A]$'s are Σ_1 . So our embedding is into the Σ_1 -formulas modulo HA^* -provable equivalence. The surprising property of the $[A]$'s is that they are closed under implication (modulo HA^* -provable equivalence). It is not true in general that the Σ -sentences of HA^* are closed under implication. This is immediate from the following well-known fact (which is a simple adaptation of Kripke's result on flexible sentences to the constructive case).

5.6.1 Fact: Let T be any consistent extension of HA . Then there is a Σ_1 -sentence Ω , such that for no Σ_1 -sentence S : $T \vdash \neg \Omega \leftrightarrow S$.

Proof: Let T be a consistent extension of HA . Take Ω such that:

$$HA \vdash \Omega \leftrightarrow \text{True}_{\Sigma}(\epsilon S. \Box_T(\neg \Omega \leftrightarrow S)).$$

Here True_Σ is the usual truth-predicate for Σ_1 -sentences and $\epsilon S. \Box_T(\neg\Omega \leftrightarrow S)$ is the first S such that $\Box_T(\neg\Omega \leftrightarrow S)$ that we find if we run through the T-proofs. Clearly $\Omega \in \Sigma_1$. Suppose for some $S' \in \Sigma_1$: $T \vdash \neg\Omega \leftrightarrow S'$. Let S be the first such S' that we encounter, when running through the T-proofs. We have: $HA \vdash S = \epsilon S'. \Box_T(\neg\Omega \leftrightarrow S')$ and hence: $HA \vdash \Omega \leftrightarrow \text{True}_\Sigma(S)$. We may conclude that $HA \vdash \Omega \leftrightarrow S$. On the other hand $T \vdash \neg\Omega \leftrightarrow S$. Ergo $T \vdash \perp$. Quod non. \square

5.7 Corollary: $A \models_{HA} B \Leftrightarrow IPC \vdash (A \rightarrow B)$.

Proof: ' \Leftarrow ' is trivial. ' \Rightarrow ' Suppose $IPC \not\vdash (A \rightarrow B)$. Then there is a finite Kripke model \mathbb{K} such that $\mathbb{K} \models A$ and $\mathbb{K} \not\models B$. Let \mathfrak{H} be the Heyting Algebra of upwards closed sets of \mathbb{K} . Obviously \mathfrak{H} is finite and hence RE. Embedding \mathfrak{H} into \mathfrak{H}_{HA}^* gives us an interpretation \mathfrak{f} such that $HA^* \vdash A[\mathfrak{f}]$ and $HA^* \not\vdash B[\mathfrak{f}]$. \square

6 On HA

Are the same Heyting Algebras embeddable in \mathfrak{H}_{HA} and in \mathfrak{H}_{IPC} ? The answer to this question is unknown. We conjecture: *yes*. In this section we show that if we restrict ourselves to Heyting Algebras on finitely many generators embeddable in $\mathfrak{S}_{HA} := \mathfrak{H}_{HA}(B\Sigma_1)$, then the answer is *yes*.

We first prove the uniform version of De Jongh's Completeness Theorem for IPC w.r.t. interpretations in HA using the result of 5.

6.1 Theorem: \mathfrak{H}_{IPC} is embeddable in \mathfrak{S}_{HA} .

Proof: Let $[\cdot]$ give the embedding of 5.1 of \mathfrak{H}_{IPC} into \mathfrak{H}_{HA}^* . Let \mathfrak{f} be given by: $\mathfrak{f}p := [p]$. We have:

$$\begin{aligned}
IPC \vdash A &\Rightarrow HA \vdash A[\mathfrak{f}] \\
&\Rightarrow HA^* \vdash A[\mathfrak{f}] \\
&\Rightarrow HA^* \vdash [A] \\
&\Rightarrow IPC \vdash A. \quad \square
\end{aligned}$$

6.2 Theorem: Let \mathfrak{H} be a Heyting Algebra on finitely many generators. Then \mathfrak{H} is embeddable in \mathfrak{S}_{HA} iff \mathfrak{H} is embeddable in $\mathfrak{H}_{\text{IPC}}$. It follows immediately that the IPC-derived rules for IPC are equal to the IPC-derived rules for HA w.r.t. substitutions involving only $\text{B}\Sigma_1$ -sentences.

To prove 6.2 we borrow three facts from Visser[85].

6.2.1 Fact: For each IPC-formula A , there is a formula A^* in NNIL such that:

- i) All propositional variables of A^* occur in A ,
- ii) For all $B \in \text{NNIL}$: $\text{IPC} \vdash B \rightarrow A \Leftrightarrow \text{IPC} \vdash B \rightarrow A^*$.

Note that 6.2.1(ii) tells us in terms of $\mathfrak{H}_{\text{IPC}}$ that $\{B \in \text{NNIL} \mid B \leq A\}$ both has and contains a supremum A^* . Thus A^* is the *greatest lower NNIL-approximant* of A .

6.2.2 Fact: Let \mathfrak{f} assign Σ_1 -sentences to the propositional variables. Then for any propositional A : $\text{HA} \vdash A[\mathfrak{f}] \Rightarrow \text{HA} \vdash A^*[\mathfrak{f}]$.

6.2.3 Fact: The number of NNIL-formulas in p_1, \dots, p_m modulo IPC-provable equivalence is finite.

Proof of 6.2: Let \mathfrak{H} be a Heyting Algebra on finitely many generators.

Suppose \mathfrak{H} is embeddable in $\mathfrak{H}_{\text{IPC}}$. By 6.1 $\mathfrak{H}_{\text{IPC}}$ is embeddable in \mathfrak{S}_{HA} and hence \mathfrak{H} is embeddable in \mathfrak{S}_{HA} .

Suppose \mathfrak{H} is embeddable in \mathfrak{S}_{HA} . Let the generators of \mathfrak{H} be A_1, \dots, A_n . These generators are in their turn Boolean combinations of Σ_1 -sentences, say, S_1, \dots, S_m . So $A_i = B_i(S_1, \dots, S_m)$ for some propositional B_i . Let \mathfrak{R} be the subalgebra of \mathfrak{S}_{HA} generated by S_1, \dots, S_m . Since \mathfrak{H} is embedded in \mathfrak{R} by assigning B_i to p_i , it is sufficient to show that \mathfrak{R} is embeddable in $\mathfrak{H}_{\text{IPC}}$. Let C^* be the greatest lower NNIL-approximant of C promised by 6.2.1. We find by 6.2.2:

$$\text{HA} \vdash C(S_1, \dots, S_m) \Rightarrow \text{HA} \vdash C^*(S_1, \dots, S_m).$$

So $\mathfrak{R} \models C \Rightarrow \mathfrak{R} \models C^*$. Since the set of NNIL-formulas in p_1, \dots, p_m is finite (modulo IPC-provable equivalence) by 6.2.3, there are only finitely many possible C^* . Let C^+

be the conjunction of the C^* . We find for D in p_1, \dots, p_m , $\mathfrak{R} \models D \Leftrightarrow \text{IPC} \vdash C^+ \rightarrow D$. Clearly C^+ is a prime NNIL-formula. By 2.8 C^+ is exact. Ergo \mathfrak{R} is embeddable in $\mathfrak{S}_{\text{IPC}}$. \square

6.3 Open question: Is \mathfrak{S}_{HA} isomorphic to $\mathfrak{S}_{\text{IPC}}$? We conjecture: *no*.

7 An RE, non-recursive Heyting Algebra on two generators

We show that there is an RE, non-recursive Heyting Algebra in two generators. It follows by 5.1 that there are Σ_1 -sentences A and B such that the subalgebra of $\mathfrak{S}_{\text{HA}}^*$ generated by A and B is RE, non-recursive. In contrast we know by 2.3 that every finitely generated Heyting Algebra embeddable in $\mathfrak{S}_{\text{IPC}}$ is decidable and similarly, by 6.2, for \mathfrak{S}_{HA} . It is still consistent with everything we know that there are arithmetical sentences A and B such that the subalgebra of \mathfrak{S}_{HA} generated by A and B is RE, non-recursive.

It is sufficient to produce an infinite decidable set X of IPC-formulas in p, q such that:

- for every finite $X_0 \subseteq X$ and every $A \in X/X_0$: $X_0 \not\models A$
- every finite $X_0 \subseteq X$ has the disjunction property

The desired algebra is obtained by taking an RE and non-recursive subset of X as axiomatization.

In de Jongh[80] infinite sequences are produced of finite rooted Kripke models \mathbb{L}_i , of formulas A_i (ψ_i^* in de Jongh[80], p107) and of formulas B_i (ψ_i in de Jongh[80], p107) such that:

- $\mathbb{L}_i \models A_j \Leftrightarrow i \neq j$
- $\mathbb{L}_i \models B_j \Leftrightarrow i = j$
- A_i is of the form $B_i \rightarrow C$ for some C
- It is decidable whether a formula is of the form A_i

(This result is originally due to Jankov, see Jankov[68])

We take X to be the set of A_i . Consider a finite $X_0 \subseteq X$ and $A \in X/X_0$. Suppose $A = A_i$. Then clearly $\mathbb{L}_i \models X_0$ and $\mathbb{L}_i \not\models A_i$. Hence $X_0 \not\models A_i$.

To prove the Disjunction Property, consider any finite $X_0 \subseteq X$. Suppose $X_0 \vdash E \vee F$, but

$X_0 \not\models E$ and $X_0 \not\models F$. Let $\mathbb{K} \models X_0$ and $\mathbb{K} \not\models E$ and $\mathbb{M} \models X_0$ and $\mathbb{M} \not\models F$. Let j be such that A_j is not in X_0 . We have: $\mathbb{L}_j \models X_0$ and $\mathbb{L}_j \not\models B_i$ for A_i in X_0 . Consider $\text{Glue}(\mathbb{K}, \mathbb{M}, \mathbb{L}_j)$. Clearly $\mathfrak{b} \not\models E$ and $\mathfrak{b} \not\models F$. Consider any $A_i \in X_0$. $\mathfrak{b} \not\models B_i$, since $\mathbb{L}_j \not\models B_i$. Since A_i is of the form $B_i \rightarrow C$ and \mathbb{K}, \mathbb{M} and \mathbb{L}_j all force A_i , it follows that $\mathfrak{b} \models A_i$. We may conclude that $\mathfrak{b} \models X_0$, but $\mathfrak{b} \not\models E$ and $\mathfrak{b} \not\models F$. A contradiction.

References

- Beeson, M.J., 1975, *The nonderivability in intuitionistic formal systems of theorems on the continuity of effective operations*, JSL 40, 321-346.
- Jankov, V.A., 1968, *Constructing a sequence of strongly independent superintuitionistic propositional calculi*, Sov. Math. Dok., 9, 806-807.
- de Jongh, D.H.J., 1970, *A characterization of the intuitionistic propositional calculus*, in Kino, Myhill, Vesley[70], 211-217.
- de Jongh, D.H.J., 1982, *Formulas of one propositional variable in intuitionistic arithmetic*, in: Troelstra & van Dalen[82], 51-64.
- Kino, A., Myhill, J., Vesley, R.E., (eds.), 1970, *Intuitionism and proof theory*, North Holland, Amsterdam.
- Pitts, A., 1992, *On an interpretation of second order quantification in first order intuitionistic propositional logic*, JSL 57, 33-52.
- Renardel de Lavalette, G.R., 1986, *Interpolation in a fragment of intuitionistic propositional logic*, Logic Group Preprint Series 5, Dept. of Philosophy, University of Utrecht, Heidelberglaan 8, 3584 CS Utrecht.
- Rybakov, V.V., 1992, *Rules of Inference with Parameters for Intuitionistic Logic*, JSL 57, 912-923.
- Shavrukov, V., 1993, *Subalgebras of diagonalizable algebras of theories containing arithmetic*, Dissertationes Mathematicae, Polska Akademia Nauk., Mathematical Institute.
- Smorynski, C., 1973, *Applications of Kripke Models*, in Troelstra[73], 324-391.
- Smorynski, C., 1985, *Self-Reference and Modal Logic*, Springer Verlag.
- Solovay, R., 1976, *Provability interpretations of modal logic*, Israel Journal of Mathematics 25, 287-304.
- Troelstra, A.S. (ed.), 1973, *Metamathematical Investigations of Intuitionistic Arithmetic and Analysis*, Springer Lecture Notes 344, Springer, Berlin.
- Troelstra, A.S., and van Dalen, D., (eds.), 1982, *The L.E.J. Brouwer Centenary*

- Symposium*, North Holland, Amsterdam.
- Troelstra, A.S., and van Dalen, D., 1988a, *Constructivism in Mathematics, vol 1*, North Holland, Amsterdam.
- Troelstra, A.S., and van Dalen, D., 1988b, *Constructivism in Mathematics, vol 2*, North Holland, Amsterdam.
- Visser, A., 1982, *On the Completeness Principle*, *Annals of Mathematical Logic* 22, 263-295.
- Visser, A., 1985, *Evaluation, provably deductive equivalence in Heyting's Arithmetic of substitution instances of propositional formulas*, Logic Group Preprint Series 4, Dept. of Philosophy, University of Utrecht, Heidelberglaan 8, 3584 CS Utrecht.
- Zambella, D., 1992, *Shavrukov's Theorem on the subalgebras of diagonalizable algebras for theories containing $I\Delta_0+Exp$* , ILLC Prepublication Series for Mathematical Logic and Foundations ML-92-05, University of Amsterdam. To appear in: *Notre Dame Journal of Formal Logic*.

The ILLC Prepublication Series

Other Prepublications

- X-91-01 Alexander Chagrov, Michael Zakharyashev The Disjunction Property of Intermediate Propositional Logics
- X-91-02 Alexander Chagrov, Michael Zakharyashev On the Undecidability of the Disjunction Property of Intermediate Propositional Logics
- X-91-03 V. Yu. Shavrukov Subalgebras of Diagonalizable Algebras of Theories containing Arithmetic
- X-91-04 K.N. Ignatiev Partial Conservativity and Modal Logics
- X-91-05 Johan van Benthem Temporal Logic
- X-91-06 Annual Report 1990
- X-91-07 A.S. Troelstra Lectures on Linear Logic, Errata and Supplement
- X-91-08 Giorgie Dzhaparidze Logic of Tolerance
- X-91-09 L.D. Beklemishev On Bimodal Provability Logics for Π_1 -axiomatized Extensions of Arithmetical Theories
- X-91-10 Michiel van Lambalgen Independence, Randomness and the Axiom of Choice
- X-91-11 Michael Zakharyashev Canonical Formulas for K4. Part I: Basic Results
- X-91-12 Herman Hendriks Flexibele Categoriale Syntaxis en Semantiek: de proefschriften van Frans Zwarts en Michael Moortgat
- X-91-13 Max I. Kanovich The Multiplicative Fragment of Linear Logic is NP-Complete
- X-91-14 Max I. Kanovich The Horn Fragment of Linear Logic is NP-Complete
- X-91-15 V. Yu. Shavrukov Subalgebras of Diagonalizable Algebras of Theories containing Arithmetic, revised version
- X-91-16 V.G. Kanovei Undecidable Hypotheses in Edward Nelson's Internal Set Theory
- X-91-17 Michiel van Lambalgen Independence, Randomness and the Axiom of Choice, Revised Version
- X-91-18 Giovanna Cepparello New Semantics for Predicate Modal Logic: an Analysis from a standard point of view
- X-91-19 Papers presented at the Provability Interpretability Arithmetic Conference, 24-31 Aug. 1991, Dept. of Phil., Utrecht University
- 1992 *Logic, Semantics and Philosophy of Language* Annual Report 1991
- LP-92-01 Victor Sánchez Valencia Lambek Grammar: an Information-based Categorical Grammar
- LP-92-02 Patrick Blackburn Modal Logic and Attribute Value Structures
- LP-92-03 Szabolcs Mikulás The Completeness of the Lambek Calculus with respect to Relational Semantics
- LP-92-04 Paul Dekker An Update Semantics for Dynamic Predicate Logic
- LP-92-05 David I. Beaver The Kinematics of Presupposition
- LP-92-06 Patrick Blackburn, Edith Spaan A Modal Perspective on the Computational Complexity of Attribute Value Grammar
- LP-92-07 Jeroen Groenendijk, Martin Stokhof A Note on Interrogatives and Adverbs of Quantification
- LP-92-08 Maarten de Rijke A System of Dynamic Modal Logic
- LP-92-09 Johan van Benthem Quantifiers in the world of Types
- LP-92-10 Maarten de Rijke Meeting Some Neighbours (a dynamic modal logic meets theories of change and knowledge representation)
- LP-92-11 Johan van Benthem A note on Dynamic Arrow Logic
- LP-92-12 Heinrich Wansing Sequent Calculi for Normal Modal Propositional Logics
- LP-92-13 Dag Westerståhl Iterated Quantifiers
- LP-92-14 Jeroen Groenendijk, Martin Stokhof Interrogatives and Adverbs of Quantification
- Mathematical Logic and Foundations*
- ML-92-01 A.S. Troelstra Comparing the theory of Representations and Constructive Mathematics
- ML-92-02 Dmitrij P. Skvortsov, Valentin B. Shehtman Maximal Kripke-type Semantics for Modal and Superintuitionistic Predicate Logics
- ML-92-03 Zoran Marković On the Structure of Kripke Models of Heyting Arithmetic
- ML-92-04 Dimiter Vakarelov A Modal Theory of Arrows, Arrow Logics I
- ML-92-05 Domenico Zambella Shavrukov's Theorem on the Subalgebras of Diagonalizable Algebras for Theories containing $\text{IA}_0 + \text{EXP}$
- ML-92-06 D.M. Gabbay, Valentin B. Shehtman Undecidability of Modal and Intermediate First-Order Logics with Two Individual Variables
- ML-92-07 Harold Schellinx How to Broaden your Horizon
- ML-92-08 Raymond Hoofman Information Systems as Coalgebras
- ML-92-09 A.S. Troelstra Realizability
- ML-92-10 V.Yu. Shavrukov A Smart Child of Peano's
- CT-92-01 Erik de Haas, Peter van Emde Boas *Computation and Complexity Theory* Object Oriented Application Flow Graphs and their Semantics
- CT-92-02 Karen L. Kwast, Sieger van Denneheuvel Weak Equivalence: Theory and Applications
- CT-92-03 Krzysztof R. Apt, Kees Doets A new Definition of SLDNF-resolution
- X-92-01 Heinrich Wansing *Other Prepublications* The Logic of Information Structures
- X-92-02 Konstantin N. Ignatiev The Closed Fragment of Dzhaparidze's Polymodal Logic and the Logic of Σ_1 conservativity
- X-92-03 Willem Groeneveld Dynamic Semantics and Circular Propositions, revised version
- X-92-04 Johan van Benthem Modeling the Kinematics of Meaning
- X-92-05 Erik de Haas, Peter van Emde Boas Object Oriented Application Flow Graphs and their Semantics, revised version
- 1993 *Logic, Semantics and Philosophy of Language*
- LP-93-01 Martijn Spaan Parallel Quantification
- LP-93-02 Makoto Kanazawa Dynamic Generalized Quantifiers and Monotonicity
- LP-93-03 Nikolai Pankrat'ev Completeness of the Lambek Calculus with respect to Relativized Relational Semantics
- LP-93-04 Jacques van Leeuwen Identity, Quarrelling with an Unproblematic Notion
- LP-93-05 Jaap van der Does Sums and Quantifiers
- LP-93-06 Paul Dekker Updates in Dynamic Semantics
- LP-93-07 Wojciech Buszkowski On the Equivalence of Lambek Categorical Grammars and Basic Categorical Grammars
- LP-93-08 Zisheng Huang, Peter van Emde Boas Information Acquisition from Multi-Agent resources; abstract
- ML-93-01 Maciej Kandiński *Mathematical Logic and Foundations* Commutative Lambek Categorical Grammars
- ML-93-02 Johan van Benthem, Natasha Alechina Modal Quantification over Structured Domains
- ML-93-03 Mati Pentus The Conjoinability Relation in Lambek Calculus and Linear Logic
- ML-93-04 Andreja Prijatelj Bounded Contraction and Many-Valued Semantics
- ML-93-05 Raymond Hoofman, Harold Schellinx Models of the Untyped λ -calculus in Semi Cartesian Closed Categories
- ML-93-06 J. Zashev Categorical Generalization of Algebraic Recursion Theory
- ML-93-07 A.V. Chagrov, L.A. Chagrova Algorithmic Problems Concerning First-Order Definability of Modal Formulas on the Class of All Finite Frames
- ML-93-08 Raymond Hoofman, Ieke Moerdijk Remarks on the Theory of Semi-Functors
- ML-93-09 A.S. Troelstra Natural Deduction for Intuitionistic Linear Logic
- ML-93-10 Vincent Danos, Jean-Baptiste Joinet, Harold Schellinx
- ML-93-11 Lex Hendriks The Structure of Exponentials: Uncovering the Dynamics of Linear Logic Proofs
- ML-93-12 V. Yu. Shavrukov Inventory of Fragments and Exact Models in Intuitionistic Propositional Logic
- ML-93-13 V. Yu. Shavrukov Remarks on Uniformly Finitely Precomplete Positive Equivalences
- ML-93-14 Dick de Jongh, Albert Visser Undecidability in Diagonalizable Algebras
- CT-93-01 Marianne Kalsbeek Embeddings of Heyting Algebras
- CT-93-02 Sophie Fischer *Computation and Complexity Theory* The Vanilla Meta-Interpreter for Definite Logic Programs and Ambivalent Syntax
- CT-93-03 Johan van Benthem, Jan Bergstra A Note on the Complexity of Local Search Problems
- CT-93-04 Karen L. Kwast, Sieger van Denneheuvel Logic of Transition Systems
- CT-93-05 Erik Aarts The Meaning of Duplicates in the Relational Database Model
- CT-93-06 Krzysztof R. Apt Proving Theorems of the Lambek Calculus of Order 2 in Polynomial Time
- CL-93-01 Noor van Leusen, László Kálmán Declarative programming in Prolog
- CL-93-02 Theo M.V. Janssen *Computational Linguistics* The Interpretation of Free Focus
- X-93-01 Paul Dekker *Other Prepublications* An Algebraic View On Rosetta
- X-93-02 Maarten de Rijke Existential Disclosure, revised version
- X-93-03 Michiel Leezenberg What is Modal Logic?
- X-93-04 A.S. Troelstra (editor) Gorani Influence on Central Kurdish: Substratum or Prestige Borrowing
- X-93-05 A.S. Troelstra (editor) Metamathematical Investigation of Intuitionistic Arithmetic and Analysis, Corrections to the First Edition
- X-93-06 Michael Zakharyashev Metamathematical Investigation of Intuitionistic Arithmetic and Analysis, Second, corrected Edition
- Canonical Formulas for K4. Part II: Cofinal Subframe Logics