

V. KANOVEI

**IST is more than an Algorithm to
prove ZFC Theorems**

ML-94-05, received: June 1994

ILLC Research Report and Technical Notes Series

Series editor: Dick de Jongh

Mathematical Logic and Foundations (ML) Series, ISSN: 0928-3315

Institute for Logic, Language and Computation (ILLC)
University of Amsterdam
Plantage Muidersgracht 24
NL-1018 TV Amsterdam
The Netherlands
e-mail: illc@fwi.uva.nl

IST is more than an algorithm to prove ZFC theorems

V.Kanovei *†

Moscow Transport Engineering Institute
Moscow State University

May 1993

Abstract

There is a sentence in the language of IST, Nelson's internal set theory, which is not equivalent in IST to a sentence in the \in -language. Thus the Reduction algorithm, that converts bounded IST formulas with standard parameters to provably (in IST) equivalent \in -formulas, cannot be extended to all formulas of the IST language.

*This research was partially supported by German grant DFG 436 rus 17/215/93 and Dutch grant NWO PGS 22-262

†kanovei@sci.math.msu.su and kanovei@math.uni-wuppertal.de

Introduction.

Internal set theory IST was invented by Nelson [1977] as an attempt to develop nonstandard mathematics from a unified axiomatic standpoint. This theory has demonstrated its ability to ground various branches of nonstandard analysis, see e.g. van den Berg [1987], F. and M. Diener [1988], F. Diener and Reeb [1989], Reeken [1992].

It is regarded as one of the advantages of IST that there exists a simple algorithm, introduced also by Nelson, to transform sentences in the language of IST to provably equivalent (in the sense of provability in IST) sentences formulated in the ZFC language. This algorithm, together with Nelson's theorem that IST is a conservative extension of ZFC, is used sometimes (see e.g. Nelson [1988]) to give back to the statement that IST is nothing more than a new way to investigate the standard ZFC universe.

This is true, indeed, so far as *bounded* IST formulas are considered. (The mentioned algorithm works for these formulas only.)

It is the aim of this paper to demonstrate that there is a certain, explicitly given sentence in the IST language which is not provably equivalent in IST to a sentence in the \in -language. Thus the IST truth cannot be completely reduced to the ZFC truth.

A sentence of this kind has to be undecidable in IST; actually the sentence we consider belongs to a type of undecidable sentences discovered and studied in Kanovei [1991]. It is as follows:

$$(*) \quad \forall F [\forall^{\text{st}}n (F(n) \text{ is standard}) \longrightarrow \exists^{\text{st}}G \forall^{\text{st}}n (F(n) = G(n))].$$

(n is assumed to range over integers, F and G over functions defined on integers and taking arbitrary values.)

Theorem 1. *Let Φ be an arbitrary \in -sentence. Then the equivalence $\Phi \longleftrightarrow (*)$ is not a theorem of IST unless IST is inconsistent.*

(Take notice that ZFC and IST are equiconsistent.) The idea of the proof is to use a pair of (transitive) models, V and V' , of a sufficiently large fragment of ZFC, elementary equivalent with respect to Φ , and then define their extensions, $*V$ and $*V'$ respectively, models of the corresponding finite fragment of IST, such that $(*)$ is false in $*V$ but true in $*V'$. Then, since $*V$ and $*V'$ are elementary extensions of V and V' respectively with

respect to \in -sentences, Φ is either simultaneously true or simultaneously false in both $*V$ and $*V'$. This proves the theorem.

Acknowledgement. The author is in debt to M.Reeken and S.Albeverio for their interest to this research direction and practical help, and to R.Solovay for a suggestion which made it possible to fix a wrong part in the first version of the proof.

Preliminaries

Theory IST was introduced by Nelson [1977]. The IST language contains, together with equality, the membership predicate \in and the standardness predicate st . Formulas of this language are called *st- \in -formulas* while formulas of the ZFC language are called *\in -formulas*, and also *internal* formulas. Two abbreviations are very useful: $\exists^{st}x \dots$ and $\forall^{st}x \dots$ (there exists standard $x \dots$, for all standard $x \dots$).

IST contains all axioms of ZFC (Separation and Replacement are formulated in the \in -language) together with the following three additional principles or (schemes of) axioms.

Idealization I: $\forall^{stfin} A \exists x \forall a \in A \Phi(x, a) \longleftrightarrow \exists x \forall^{st} a \Phi(x, a)$
for any internal formula $\Phi(x, a)$.

Standardization S: $\forall^{st} X \exists^{st} Y \forall^{st} x [x \in Y \longleftrightarrow x \in X \ \& \ \Phi(x)]$
for any st- \in -formula Φ .

Transfer T: $\exists x \Phi(x) \longrightarrow \exists^{st} x \Phi(x)$
for any internal formula $\Phi(x)$ with standard parameters.

The formula Φ can, of course, contain *arbitrary* parameters in I and S.

Thus $IST = ZFC + I + S + T$.

Definition. Let V be a transitive set. $*V = \langle *V; * \in, *=, *st \rangle$ is an *IST-like* extension of V if and only if, first, axioms I, S, T hold in $*V$, and second, there exists an 1-1 embedding $*$: V onto a subset of $*V$ satisfying

- $x \in y \longleftrightarrow *x * \in *y$ and $x = y \longleftrightarrow *x *= *y$ for all $x, y \in V$,
and

- $*\text{st} X \iff \exists x \in V (*x \doteq X)$ for all $X \in *V$.

It is not assumed, in general, that \doteq coincides with the true equality on $*V$, but \doteq has to be an equivalence relation and satisfy the logic axioms for equality with respect to $*\in$ and $*\text{st}$.

Proof of Theorem 1.

Assume on the contrary that Φ is an \in -sentence such that the equivalence $\Phi \iff (*)$ is a theorem of IST, therefore of a theory

$$\text{IST}' = \text{ZFC}' + \text{I} + \text{S} + \text{T},$$

where ZFC' is a finite fragment of ZFC. Having this fixed, we start to argue in ZFC. The final aim is to obtain a contradiction.

Ground ZFC' models.

It is a consequence of the ZFC Reflection principle that there exist cardinals ϑ of both countable and uncountable cofinality such that \mathbf{V}_ϑ is an elementary submodel of the universe of all sets with respect to Φ and all formulas of ZFC' .

Let ϑ be the least among the countably cofinal while ϑ' among the uncountably cofinal cardinals of this kind. We use the sets $V = \mathbf{V}_\vartheta$ and $V' = \mathbf{V}_{\vartheta'}$ as the ground ZFC' models. Take notice that Φ is either true in both V and V' or false in both V and V' .

The next step is to define IST-extensions (therefore models of IST'), $*V$ and $*V'$, of V and V' respectively, such that $(*)$ is true in $*V'$ but false in $*V$. The extensions are constructed as ultrapowers via a kind of *adequate* ultrafilters of Nelson [1977]. (Original Nelson's construction includes infinite number of successive ultrapowers; we show here that this can be managed an one-step construction.)

The “falsity” extension

Thus we define $*V$ as an ultrapower of V using the *index set*

$$I = \mathcal{P}^{\text{fn}}(V) = \{i \in V : i \text{ is finite}\},$$

and an arbitrary *ultrafilter* U over I containing all sets of the form $I_a = \{i \in I : a \in i\}$, $a \in V$.

We introduce a convenient tool, the quantifier “there exist U -many” by

$$\mathbf{U}i\varphi(i) \text{ if and only if } \{i \in I : \varphi(i)\} \in U.$$

The following is the list of properties of \mathbf{U} implied by the definition of an ultrafilter and (this regards (U5)) the choice of the ultrafilter U .

- (U1) $\varphi \longleftrightarrow \mathbf{U}i\varphi$ whenever i is not free in φ ;
- (U2) if $\forall i[\varphi(i) \longrightarrow \psi(i)]$ then $\mathbf{U}i\varphi(i) \longrightarrow \mathbf{U}i\psi(i)$;
- (U3) $\mathbf{U}i\varphi(i) \ \& \ \mathbf{U}i\psi(i) \longleftrightarrow \mathbf{U}i[\varphi(i) \ \& \ \psi(i)]$;
- (U4) $\mathbf{U}i\neg\varphi(i) \longleftrightarrow \neg\mathbf{U}i\varphi(i)$;
- (U5) if $a \in V$ then $\mathbf{U}i(a \in i)$.

To introduce the extension, we put

$${}^*V_r = \{f : f \text{ is a function, } f : I^r \longrightarrow V\}, \quad \text{for all } r \in \omega.$$

In particular, ${}^*V_0 = \{{}^*z : z \in V\}$, where ${}^*z = \{\langle 0, z \rangle\}$, since $I^0 = \{\emptyset\}$.

The set ${}^*V = \bigcup_{r \in \omega} {}^*V_r$ is what we call *the falsity extension*.

To continue notation, we let, for $F \in {}^*V$, $r(F)$ denote the unique r satisfying $F \in {}^*V_r$. If $F \in {}^*V$, $q \geq r = r(F)$, $\mathbf{i} = \langle i_1, \dots, i_r, \dots, i_q \rangle \in I^q$, then we put $F[\mathbf{i}] = F(i_1, \dots, i_r)$. Note that $F[\mathbf{i}] = F(\mathbf{i})$ whenever $r = q$. We define finally ${}^*z[\mathbf{i}] = z$ for all ${}^*z \in {}^*V_0$ and $\mathbf{i} \in I^r$, $r \geq 0$.

Let $F, G \in {}^*V$ and $r = \max\{r(F), r(G)\}$. We set

$$F \vDash G \text{ if and only if } \mathbf{U}i_r \mathbf{U}i_{r-1} \dots \mathbf{U}i_1 (F[\mathbf{i}] \in G[\mathbf{i}]);$$

$$F \vDash^* G \text{ if and only if } \mathbf{U}i_r \mathbf{U}i_{r-1} \dots \mathbf{U}i_1 (F[\mathbf{i}] = G[\mathbf{i}]);$$

of course \mathbf{i} denotes the sequence i_1, \dots, i_r .

The definition of standardness in *V is given by:

$${}^*\text{st}F \text{ if and only if there exists } x \in V \text{ such that } F \vDash^* x.$$

So up to the relation \vDash^* the level *V_0 is just the standard part of *V .

Let, finally, Ψ be a formula with parameters in *V . We define $r(\Psi) = \max\{r(F) : F \text{ occurs in } \Psi\}$. If in addition $r \geq r(\Psi)$ and $\mathbf{i} \in I^r$, then let $\Psi[\mathbf{i}]$ denote the result of replacing each F that occurs in Ψ by $F[\mathbf{i}]$. Clearly $\Psi[\mathbf{i}]$ is a formula with parameters in V .

Proposition 2. $\langle {}^*V; =, \in, \text{st} \rangle$ is a model of IST' and an IST -like extension of V . Finally, $(*)$ fails in *V .

Proof. The following principal statement plays the key role.

Lemma 3. [Łoś Theorem] Let Ψ be an internal formula with parameters in *V and suppose that $r \geq r(\Psi)$. Then

$$\Psi \text{ is true in } {}^*V \iff \mathbf{U} i_r \dots \mathbf{U} i_1 (\Psi[i_1, \dots, i_r] \text{ is true in } V).$$

Proof of the lemma. The proof goes by induction on the logical complexity of Ψ . We abandon easy parts of the proof, based on properties (U2), (U3), (U4) of the quantifier \mathbf{U} , and consider the induction step \exists . Thus the lemma is to be proved for a formula $\exists x \Psi(x)$ in the assumption that the result holds for $\Psi(F)$ whenever $F \in {}^*V$. We denote $r = r(\Psi)$.

The direction \longrightarrow . Suppose that $\exists x \Psi(x)$ holds in *V . Then $\Psi(F)$ holds in *V for some $F \in {}^*V$. Let $p = \max\{r, r(F)\}$. To convert the reasoning into a more convenient form, we let \mathbf{i} and \mathbf{j} denote sequences

$$\langle i_1, \dots, i_r \rangle (\in I^r) \quad \text{and} \quad \langle i_1, \dots, i_r, \dots, i_p \rangle (\in I^p)$$

respectively. Further let $\mathbf{U} \mathbf{i}$ and $\mathbf{U} \mathbf{j}$ denote sequences of quantifiers

$$\mathbf{U} i_r \dots \mathbf{U} i_1 \quad \text{and} \quad \mathbf{U} i_p \dots \mathbf{U} i_r \dots \mathbf{U} i_1.$$

Thus $\mathbf{U} \mathbf{j} \Psi(F)[\mathbf{j}]$ holds by the induction hypothesis. We note that, for all \mathbf{j} , $\Psi(F)[\mathbf{j}] \longrightarrow \exists x \Psi(x)[\mathbf{j}]$. Hence $\mathbf{U} \mathbf{j} \exists x \Psi(x)[\mathbf{j}]$ is true by (U2). Moreover the formula $\exists x \Psi(x)[\mathbf{j}]$ coincides (graphically) with $\exists x \Psi(x)[\mathbf{i}]$ because $r(\exists x \Psi(x)) = r \leq p$. Hence, deleting the superfluous quantifiers by (U1), we obtain $\mathbf{U} \mathbf{i} \exists x \Psi(x)[\mathbf{i}]$.

The direction \longleftarrow . Let $\Psi(x)$ be $\Psi(x, G, H, \dots)$, where $G, H, \dots \in {}^*V$. Suppose that $\mathbf{U} \mathbf{i} \exists x \Psi(x)[\mathbf{i}]$ holds, that is,

$$\mathbf{U} \mathbf{i} [\exists x \Psi(x, G[\mathbf{i}], H[\mathbf{i}], \dots)] \text{ is true in } V].$$

For each $\mathbf{i} \in I^r$, if there exists some $x \in V$ such that $\Psi(x, G[\mathbf{i}], H[\mathbf{i}], \dots)$ is true in V , then we let $F(\mathbf{i})$ be one of x of such kind; otherwise let $F(\mathbf{i}) = \emptyset$. By definition, $F \in {}^*V_r$ and

$$\forall \mathbf{i} \in I^r [\exists x \Psi(x) [\mathbf{i}] \longrightarrow \Psi(F) [\mathbf{i}]],$$

therefore $\mathbf{U} \mathbf{i} \exists x \Psi(x) [\mathbf{i}] \longrightarrow \mathbf{U} \mathbf{i} \Psi(F) [\mathbf{i}]$ by (U2). Recall that the left-hand side of the last implication has been supposed to be true. So the right-hand side is also true. Then $\Psi(F)$ holds in *V by the induction hypothesis, and we are done.

The just proved lemma easily implies logical equality axioms for ${}^* =$, and Transfer, therefore all ZFC', in *V . Standardization is evident because every set V of the form $V = \mathbf{V}_\emptyset$ has the property that if $Y \subseteq X \in V$ then $Y \in V$. We prove Idealization.

Thus let $\varphi(x, a)$ be an internal formula with parameters in *V . We denote $r = r(\varphi)$ and prove the following:

$$\forall^{\text{stfn}} A \exists x \forall a \in A \varphi(x, a) \longrightarrow \exists x \forall^{\text{st}} a \varphi(x, a)$$

in *V . (The implication \longleftarrow does not need a special consideration because it follows from Standardization that elements of finite standard sets are standard, see Nelson [1977].) Lemma 3 converts the left-hand side to the form:

$$\forall^{\text{fn}} A \subseteq V \mathbf{U} i_r \dots \mathbf{U} i_1 \exists x \forall a \in A (\varphi(x, a) [i_1, \dots, i_r]).$$

Recall that I consists of all finite subsets of V , so we may replace the variable A by i , having in mind that $i \in I$. Further define $\tilde{A} : I^{r+1} \longrightarrow V$ by $\tilde{A}(i_1, \dots, i_r, i) = i$. Then $\tilde{A} \in {}^*V$. The left-hand side takes the form

$$\forall i \mathbf{U} i_r \dots \mathbf{U} i_1 (\exists x \forall a \in \tilde{A} \varphi(x, a)) [i_1, \dots, i_r, i].$$

Changing $\forall i$ by $\mathbf{U} i$, we obtain $\exists x \forall a \in \tilde{A} \varphi(x, a)$ in *V again by the lemma. So, to verify the right-hand side of Idealization, it suffices to prove ${}^*a \in \tilde{A}$ in *V for all $a \in V$. This is equal to

$$\mathbf{U} i \mathbf{U} i_r \dots \mathbf{U} i_1 (a \in \tilde{A} [i_1, \dots, i_r, i]),$$

by the lemma, and then to $\mathbf{U} i \mathbf{U} i_r \dots \mathbf{U} i_1 (a \in i)$ by the definition of \tilde{A} . So apply (U5) and complete the proof of Idealization in *V .

Thus *V is an IST' model. One can easily verify the required properties of the embedding * . To complete the proof of Proposition 2 it remains to show that $(*)$ does not hold in *V .

Let $\langle \kappa_n : n \in \omega \rangle$ be a sequence of ordinals cofinal in ϑ . (We recall that ϑ has countable cofinality.) Let $F \in {}^*V_0$ be defined by

$$F(i) = \{ \langle n, \kappa_n \rangle : \langle n, \kappa_n \rangle \in i \} \quad \text{for all } i \in I.$$

It is true in *V by Lemma 3 that F is a function defined on a subset of integers, and, for every $n \in \omega$, it is also true in *V that $F({}^*n)$ is defined and equal to ${}^*\kappa_n$, hence standard. Thus the left-hand side of $(*)$ is satisfied by F .

The right-hand side cannot be satisfied since it would imply that there exists $g \in V$ such that $g(n) = \kappa_n$ for all n , which is impossible.

Corollary 4. Φ is false in *V , therefore in V .

The “truth” extension

Let ${}^*V'$ be defined the same way as *V above, but starting from V' .

Proposition 5. $\langle {}^*V'; {}^*=, {}^*\in, {}^*\text{st} \rangle$ is a model of IST' and an IST -like extension of V . Finally, $(*)$ holds in ${}^*V'$.

Proof. We check the last statement. Thus let $F \in {}^*V'$ be such that the following is true in ${}^*V'$:

F is a function, every standard $n \in {}^*\omega$ belongs to the domain of F , and $F(n)$ is standard for every standard $n \in {}^*\omega$.

By the definition of standardness, there exists a function $f : \omega \rightarrow V'$ such that $F({}^*n) \stackrel{*}{=} {}^*(f(n))$ for all $n \in \omega$. By the choice of ϑ' (uncountable cofinality) there exists $\kappa < \vartheta'$ such that $f(n) \in \mathbf{V}_\kappa$ for all $n \in \omega$. This easily implies that actually $f \in V'$, and therefore $F({}^*n) \stackrel{*}{=} ({}^*f)({}^*n)$ for all $n \in \omega$, the right-hand side of $(*)$.

Corollary 6. Φ is true in ${}^*V'$, therefore in V' .


Thus finally Φ is true in V' and false in V , a contradiction with the choice of V, V' as models elementary equivalent with respect to Φ .

Question

Does there exist, in ZFC, a transitive set V , a model of a previously fixed finite fragment of ZFC, which has IST-like extensions of both types, those in which $(*)$ holds and those where $(*)$ fails? The answer is affirmative provided there is a cardinal ϑ such that \mathbf{V}_ϑ is a model of the full ZFC (then we may take $V = \mathbf{V}_\vartheta$, where ϑ is the least among such cardinals), but we are unable to get it without extra assumptions. If this is actually impossible, then, perhaps, $(*)$ still corresponds to something in ZFC, not in the direct form mentioned in Theorem 1, of course.

References

- I. van den Berg [1987] *Nonstandard asymptotic analysis* (Lecture Notes in Math. 1249, Springer).
- F.Diener and M.Diener [1988] Some asymptotic results in ordinary differential equations. in: N.Cutland (ed.) *Nonstandard analysis and its applications* (London Math. Soc. Student Texts 10, Cambridge Univ. Press), pp. 282 – 297.
- F.Diener and K.D.Stroyan [1988] Syntactical methods in infinitesimal analysis, in: N.Cutland (ed.) *Nonstandard analysis and its applications* (London Math. Soc. Student Texts 10, Cambridge Univ. Press), pp. 258 – 281.
- F.Diener and G.Reeb [1989] *Analyse non standard* (Herrmann Editeurs).
- V.G.Kanovei [1991] Undecidable hypotheses in Edward Nelson's internal set theory, *Russian Math. Surveys*, 46, pp. 1 – 54.
- E.Nelson [1977] Internal set theory; a new approach to nonstandard analysis, *Bull. Amer. Math. Soc.* 83, pp. 1165 – 1198.
- E.Nelson [1988] The syntax of nonstandard analysis, *Ann. Pure Appl. Log.* 38, 123 – 134.
- M.Reeken [1992] On external constructions in internal set theory. *Expositiones Mathematicae*, 10, 193 – 247.



ILLC Research Reports and Technical Notes

Coding for Research Reports: *Series-Year-Number*, with LP = Linguistics and Philosophy of Language; ML = Mathematical Logic and Foundations; CL = Computational Linguistics; CT = Computation and Complexity Theory; X = Technical Notes.

All previous ILLC-publications are available from the ILLC bureau. For prepublications before 1993, contact the bureau.

- LP-93-01 *Martijn Spaan, Parallel Quantification*
- LP-93-02 *Makoto Kanazawa, Dynamic Generalized Quantifiers and Monotonicity*
- LP-93-03 *Nikolai Pankrat'ev, Completeness of the Lambek Calculus with respect to Relativized Relational Semantics*
- LP-93-04 *Jacques van Leeuwen, Identity, Quarrelling with an Unproblematic Notion*
- LP-93-05 *Jaap van der Does, Sums and Quantifiers*
- LP-93-06 *Paul Dekker, Updates in Dynamic Semantics*
- LP-93-07 *Wojciech Buszkowski, On the Equivalence of Lambek Categorical Grammars and Basic Categorical Grammars*
- LP-93-08 *Zisheng Huang, Peter van Emde Boas, Information Acquisition from Multi-Agent resources; abstract*
- LP-93-09 *Makoto Kanazawa, Completeness and Decidability of the Mixed Style of Inference with Composition*
- LP-93-10 *Makoto Kanazawa, Weak vs. Strong Readings of Donkey Sentences and Monotonicity Inference in a Dynamic Setting*
- LP-93-11 *Friederike Moltmann, Resumptive Quantifiers in Exception Sentences*
- LP-93-12 *Jaap van der Does, On Complex Plural Noun Phrases*
- LP-93-13 *Natasha Alechina, Binary Quantifiers and Relational Semantics*
- LP-93-14 *Mati Pentus, Lambek Calculus is L-complete*
- LP-93-15 *David Ian Beaver, What comes first in Dynamic Semantics*

- ML-93-01 *Maciej Kandulski, Commutative Lambek Categorical Grammars*
- ML-93-02 *Johan van Benthem, Natasha Alechina, Modal Quantification over Structured Domains*
- ML-93-03 *Mati Pentus, The Conjoinability Relation in Lambek Calculus and Linear Logic*
- ML-93-04 *Andreja Prijatelj, Bounded Contraction and Many-Valued Semantics*
- ML-93-05 *Raymond Hoofman, Harold Schellinx, Models of the Untyped λ -calculus in Semi Cartesian Closed Categories*
- ML-93-06 *J. Zashev, Categorical Generalization of Algebraic Recursion Theory*
- ML-93-07 *A.V. Chagrov, L.A. Chagrova, Algorithmic Problems Concerning First-Order Definability of Modal Formulas on the Class of All Finite Frames*
- ML-93-08 *Raymond Hoofman, Ieke Moerdijk, Remarks on the Theory of Semi-Functors*
- ML-93-09 *A.S. Troelstra, Natural Deduction for Intuitionistic Linear Logic*
- ML-93-10 *Vincent Danos, Jean-Baptiste Joinet, Harold Schellinx, The Structure of Exponentials: Uncovering the Dynamics of Linear Logic Proofs*
- ML-93-11 *Lex Hendriks, Inventory of Fragments and Exact Models in Intuitionistic Propositional Logic*
- ML-93-12 *V.Yu. Shavrukov, Remarks on Uniformly Finitely Precomplete Positive Equivalences*
- ML-93-13 *V.Yu. Shavrukov, Undecidability in Diagonalizable Algebras*
- ML-93-14 *Dick de Jongh, Albert Visser, Embeddings of Heyting Algebras*
- ML-93-15 *G.K. Dzhaparidze, Effective Truth*
- ML-93-16 *Maarten de Rijke, Correspondence Theory for Extended Modal Logics*
- ML-93-17 *Alexander Chagrov, Michael Zakharyashev, On the Independent Axiomatizability of Modal and Intermediate Logics*
- ML-93-18 *Jaap van Oosten, Extensional Realizability*
- ML-93-19 *Raymond Hoofman, Comparing Models of the Non-Extensional Typed λ -Calculus*
- ML-93-20 *L.A. Chagrova, Dick de Jongh, The Decidability of Dependency in Intuitionistic Propositional Logic*
- ML-93-21 *Max I. Kanovich, The Relational Knowledge-Base Interpretation and Feasible Theorem Proving for Intuitionistic Propositional Logic*

- ML-93-22 *Andreja Prijatelj, Connectification for n -contraction*
- CT-93-01 *Marianne Kalsbeek, The Vanilla Meta-Interpreter for Definite Logic Programs and Ambivalent Syntax*
 CT-93-02 *Sophie Fischer, A Note on the Complexity of Local Search Problems*
 CT-93-03 *Johan van Benthem, Jan Bergstra, Logic of Transition Systems*
 CT-93-04 *Karen L. Kwast, Sieger van Denneheuvel, The Meaning of Duplicates in the Relational Database Model*
 CT-93-05 *Erik Aarts, Proving Theorems of the Lambek Calculus of Order 2 in Polynomial Time*
 CT-93-06 *Krzysztof R. Apt, Declarative programming in Prolog*
 CT-93-07 *Janusz A. Pomykala, Approximation, Similarity and Rough Constructions, Part I. Elementary Introduction*
- CL-93-01 *Noor van Leusen, László Kálmán, Computational Linguistics*
 CL-93-02 *Theo M.V. Janssen, An Algebraic View On Rosetta*
 CL-93-03 *Patrick Blackburn, Claire Gardent, Wilfried Meyer-Viol, Talking about Trees*
- X-93-01 *Paul Dekker, Existential Disclosure, revised version*
 X-93-02 *Maarten de Rijke, What is Modal Logic?*
 X-93-03 *Michiel Leezenberg, Gorani Influence on Central Kurdish: Substratum or Prestige Borrowing*
 X-93-04 *A.S. Troelstra (editor), Metamathematical Investigation of Intuitionistic Arithmetic and Analysis, Corrections to the First Edition*
 X-93-05 *A.S. Troelstra (editor), Metamathematical Investigation of Intuitionistic Arithmetic and Analysis, Second, corrected Edition*
 X-93-06 *Michael Zakharyashev, Canonical Formulas for K4. Part II: Cofinal Subframe Logics*
- ML-94-01 *Domenico Zambella, Notes on polynomially bounded arithmetic*
 ML-94-02 *Domenico Zambella, End Extensions of Models of Linearly Bounded Arithmetic*
 ML-94-03 *Johan van Benthem, Dick de Jongh, Gerard Renardel de Lavalette, Albert Visser, NNIL, A Study in Intuitionistic Propositional Logic*
 ML-94-04 *Michiel van Lambalgen, Independence Structures in Set Theory*
 ML-94-05 *V. Kanovei, IST is More than an Algorithm to Prove ZFC Theorems*
 LP-94-01 *Dimitar Gelev, Introducing Some Classical Elements of Modal Logic to the Propositional Logics of Qualitative Probabilities*
 LP-94-02 *Andrei Arsov, Basic Arrow Logic with Relation Algebraic Operators*
 LP-94-03 *Jerry Seligman, An algebraic appreciation of diagrams*
 LP-94-04 *Kazimierz Świrydowicz, A Remark on the Maximal Extensions of the Relevant Logic R*
 LP-94-05 *Natasha Kurtonina, The Lambek Calculus: Relational Semantics and the Method of Labelling*
 LP-94-06 *Johan van Benthem, Dag Westerståhl, Directions in Generalized Quantifier Theory*
 LP-94-07 *Nataša Rakić, Absolute Time, Special Relativity and ML*
 CT-94-01 *Harry Buhrman and Leen Torenvliet, On the Cutting Edge of Relativization: the Resource Bounded Injury Method*
 CT-94-02 *Alessandro Panconesi, Marina Papatriantafidou, Philippos Tsigas, Paul Vitányi, Randomized Wait-Free Distributed Naming*
 CT-94-03 *Ming Lee, John Tromp, Paul Vitányi, Sharpening Occam's Razor (extended abstract)*
 CT-94-04 *Ming Lee and Paul Vitányi, Inductive Reasoning*
 CT-94-05 *Tao Jiang, Joel I. Seiferas, Paul M.B. Vitányi, Two heads are Better than Two Tapes*
 CT-94-06 *Guido te Brake, Joost N. Kok, Paul Vitányi, Model Selection for Neural Networks: Comparing MDL and NIC*
 CT-94-07 *Charles H. Bennett, Péter Gács, Ming Li, Paul M.B. Vitányi, Wojciech H. Zurek, Thermodynamics of Computation and Information Distance*

- CT-94-08 Krzysztof R. Apt, Peter van Emde Boas and Angelo Welling, *The STO-problem is NP-hard*
CT-94-09 Klaus Ambos-Spies, Sebastiaan A. Terwijn, Zheng Xizhong, *Resource Bounded Randomness and Weakly Complete Problems*
CT-94-10 Klaus Ambos-Spies, Hans-Christian Neis, Sebastiaan A. Terwijn, *Genericity and Measure for Exponential Time*
X-94-01 Johan van Benthem, *Two Essays on Semantic Modelling*
X-94-02 Vladimir Kanovei, Michiel van Lambalgen, *Another Construction of Choiceless Ultrapower*
X-94-03 Natasha Alechina, Michiel van Lambalgen, *Correspondence and Completeness for Generalized Quantifiers*

Titles in the ILLC Dissertation Series:

- 1993-1 *Transsentential Meditations; Ups and downs in dynamic semantics*, Paul Dekker
1993-2 *Resource Bounded Reductions*, Harry Buhrman
1993-3 *Efficient Metamathematics*, Rineke Verbrugge
1993-4 *Extending Modal Logic*, Maarten de Rijke
1993-5 *Studied Flexibility*, Herman Hendriks
1993-6 *Aspects of Algorithms and Complexity*, John Tromp
1994-1 *The Noble Art of Linear Decorating*, Harold Schellinx
1994-2 *Generating Uniform User-Interfaces for Interactive Programming Environments*, Jan Willem Cornelis Koorn
1994-3 *Process Theory and Equation Solving*, Nicoline Johanna Drost
1994-4 *Calculi for Constructive Communication, a Study of the Dynamics of Partial States*, Jan Jaspars
1994-5 *Executable Language Definitions, Case Studies and Origin Tracking Techniques*, Arie van Deursen