

Institute for Language, Logic and Information


PROVABLE FIXED POINTS IN $\text{IA}_0 + \Omega_1$

revised version

Alessandra Carbone

ITLI Prepublication Series

X-90-11



University of Amsterdam

The ITLI Prepublication Series

- 1986
- 86-01 The Institute of Language, Logic and Information
- 86-02 Peter van Emde Boas A Semantical Model for Integration and Modularization of Rules
- 86-03 Johan van Benthem Categorical Grammar and Lambda Calculus
- 86-04 Reinhard Muskens A Relational Formulation of the Theory of Types
- 86-05 Kenneth A. Bowen, Dick de Jongh Some Complete Logics for Branched Time, Part I Well-founded Time, Forward looking Operators
- 86-06 Johan van Benthem Logical Syntax
- 1987 87-01 Jeroen Groenendijk, Martin Stokhof Type shifting Rules and the Semantics of Interrogatives
- 87-02 Renate Bartsch Frame Representations and Discourse Representations
- 87-03 Jan Willem Klop, Roel de Vrijer Unique Normal Forms for Lambda Calculus with Surjective Pairing
- 87-04 Johan van Benthem Polyadic quantifiers
- 87-05 Víctor Sánchez Valencia Traditional Logicians and de Morgan's Example
- 87-06 Eleonore Oversteegen Temporal Adverbials in the Two Track Theory of Time
- 87-07 Johan van Benthem Categorical Grammar and Type Theory
- 87-08 Renate Bartsch The Construction of Properties under Perspectives
- 87-09 Herman Hendriks Type Change in Semantics: The Scope of Quantification and Coordination
- 1988 LP-88-01 Michiel van Lambalgen *Logic, Semantics and Philosophy of Language:* Algorithmic Information Theory
- LP-88-02 Yde Venema Expressiveness and Completeness of an Interval Tense Logic
- LP-88-03 Year Report 1987
- LP-88-04 Reinhard Muskens Going partial in Montague Grammar
- LP-88-05 Johan van Benthem Logical Constants across Varying Types
- LP-88-06 Johan van Benthem Semantic Parallels in Natural Language and Computation
- LP-88-07 Renate Bartsch Tenses, Aspects, and their Scopes in Discourse
- LP-88-08 Jeroen Groenendijk, Martin Stokhof Context and Information in Dynamic Semantics
- LP-88-09 Theo M.V. Janssen A mathematical model for the CAT framework of Eurotra
- LP-88-10 Anneke Kleppe A Blissymbolics Translation Program
- ML-88-01 Jaap van Oosten *Mathematical Logic and Foundations:* Lifschitz' Realizability
- ML-88-02 M.D.G. Swaen The Arithmetical Fragment of Martin Löf's Type Theories with weak Σ -elimination
- ML-88-03 Dick de Jongh, Frank Veltman Provability Logics for Relative Interpretability
- ML-88-04 A.S. Troelstra On the Early History of Intuitionistic Logic
- ML-88-05 A.S. Troelstra Remarks on Intuitionism and the Philosophy of Mathematics
- CT-88-01 Ming Li, Paul M.B. Vitanyi *Computation and Complexity Theory:* Two Decades of Applied Kolmogorov Complexity
- CT-88-02 Michiel H.M. Smid General Lower Bounds for the Partitioning of Range Trees
- CT-88-03 Michiel H.M. Smid, Mark H. Overmars Maintaining Multiple Representations of Dynamic Data Structures
- Leen Torenvliet, Peter van Emde Boas
- CT-88-04 Dick de Jongh, Lex Hendriks Computations in Fragments of Intuitionistic Propositional Logic
- Gerard R. Renardel de Lavalette
- CT-88-05 Peter van Emde Boas Machine Models and Simulations (revised version)
- CT-88-06 Michiel H.M. Smid A Data Structure for the Union-find Problem having good Single-Operation Complexity
- CT-88-07 Johan van Benthem Time, Logic and Computation
- CT-88-08 Michiel H.M. Smid, Mark H. Overmars Multiple Representations of Dynamic Data Structures
- Leen Torenvliet, Peter van Emde Boas
- CT-88-09 Theo M.V. Janssen Towards a Universal Parsing Algorithm for Functional Grammar
- CT-88-10 Edith Spaan, Leen Torenvliet, Peter van Emde Boas Nondeterminism, Fairness and a Fundamental Analogy
- CT-88-11 Sieger van Denneheuvel, Peter van Emde Boas Towards implementing RL
- X-88-01 Marc Jumelet *Other prepublications:* On Solovay's Completeness Theorem
- 1989 LP-89-01 Johan van Benthem *Logic, Semantics and Philosophy of Language:* The Fine-Structure of Categorical Semantics
- LP-89-02 Jeroen Groenendijk, Martin Stokhof Dynamic Predicate Logic, towards a compositional, non-representational semantics of discourse
- LP-89-03 Yde Venema Two-dimensional Modal Logics for Relation Algebras and Temporal Logic of Intervals
- LP-89-04 Johan van Benthem Language in Action
- LP-89-05 Johan van Benthem Modal Logic as a Theory of Information
- LP-89-06 Andreja Prijatelj Intensional Lambek Calculi: Theory and Application
- LP-89-07 Heinrich Wansing The Adequacy Problem for Sequential Propositional Logic
- LP-89-08 Víctor Sánchez Valencia Peirce's Propositional Logic: From Algebra to Graphs
- LP-89-09 Zhisheng Huang Dependency of Belief in Distributed Systems
- ML-89-01 Dick de Jongh, Albert Visser *Mathematical Logic and Foundations:* Explicit Fixed Points for Interpretability Logic
- ML-89-02 Roel de Vrijer Extending the Lambda Calculus with Surjective Pairing is conservative
- ML-89-03 Dick de Jongh, Franco Montagna Rosser Orderings and Free Variables
- ML-89-04 Dick de Jongh, Marc Jumelet, Franco Montagna On the Proof of Solovay's Theorem
- ML-89-05 Rincke Verbrugge Σ -completeness and Bounded Arithmetic
- ML-89-06 Michiel van Lambalgen The Axiomatization of Randomness
- ML-89-07 Dirk Roorda Elementary Inductive Definitions in HA: from Strictly Positive towards Monotone
- ML-89-08 Dirk Roorda Investigations into Classical Linear Logic
- ML-89-09 Alessandra Carbone Provable Fixed points in $IA_0 + \Omega_1$
- CT-89-01 Michiel H.M. Smid *Computation and Complexity Theory:* Dynamic Deferred Data Structures
- CT-89-02 Peter van Emde Boas Machine Models and Simulations
- CT-89-03 Ming Li, Herman Nueföglise, Leen Torenvliet, Peter van Emde Boas On Space Efficient Simulations
- CT-89-04 Harry Buhrman, Leen Torenvliet A Comparison of Reductions on Nondeterministic Space
- CT-89-05 Pieter H. Hartel, Michiel H.M. Smid A Parallel Functional Implementation of Range Queries
- Leen Torenvliet, Willem G. Vree
- CT-89-06 H.W. Lenstra, Jr. Finding Isomorphisms between Finite Fields
- CT-89-07 Ming Li, Paul M.B. Vitanyi A Theory of Learning Simple Concepts under Simple Distributions and Average Case Complexity for the Universal Distribution (Prel. Version)
- CT-89-08 Harry Buhrman, Steven Homer Honest Reductions, Completeness and Nondeterministic Complexity Classes
- Leen Torenvliet
- CT-89-09 Harry Buhrman, Edith Spaan, Leen Torenvliet On Adaptive Resource Bounded Computations
- CT-89-10 Sieger van Denneheuvel The Rule Language RL/1
- CT-89-11 Zhisheng Huang, Sieger van Denneheuvel Towards Functional Classification of Recursive Query Processing
- Peter van Emde Boas
- X-89-01 Marianne Kalsbeek *Other Prepublications:* An Orey Sentence for Predicative Arithmetic
- X-89-02 G. Wagemakers New Foundations: a Survey of Quine's Set Theory
- X-89-03 A.S. Troelstra Index of the Heyting Nachlass
- X-89-04 Jeroen Groenendijk, Martin Stokhof Dynamic Montague Grammar, a first sketch
- X-89-05 Maarten de Rijke The Modal Theory of Inequality
- X-89-06 Peter van Emde Boas Een Relationele Semantiek voor Conceptueel Modelleren: Het RL-project
- 1990 SEE INSIDE BACK COVER



Instituut voor Taal, Logica en Informatie
Institute for Language, Logic and Information

Faculteit der Wiskunde en Informatica
(Department of Mathematics and Computer Science)
Plantage Muidergracht 24
1018TV Amsterdam

Faculteit der Wijsbegeerte
(Department of Philosophy)
Nieuwe Doelenstraat 15
1012CP Amsterdam

PROVABLE FIXED POINTS IN $\mathcal{I}\Delta_0 + \Omega_1$

revised version

Alessandra Carbone

Department of Mathematics and Computer Science
University of Amsterdam

Department of Mathematics
Graduate School and University Centre, CUNY
33 W. 42nd Str., New York, NY 10036

Received July 1990

revised version of
ITLI ML-89-09

1. INTRODUCTION ¹

This work should be considered as part of the general investigation into the arithmetical system $I\Delta_0 + \Omega_1$. We will present a refinement to $I\Delta_0 + \Omega_1$ of a result stated in [deJongh-Montagna, 1988], on witness comparison formulas having only provable fixed points in PA.

Briefly, let us introduce the arithmetical system and some of its properties: $I\Delta_0 + \Omega_1$ (Cf.[Paris-Wilkie,1987]) is a set of axioms expressing the elementary arithmetic properties of the basic symbols $0, ', +, *, \leq$ (in the following we will refer to the obvious first order language containing these symbols as S) together with the bounded induction schema $I\Delta_0$ (defined in S):

$$\forall x, z (\varphi(x, 0) \wedge \forall y \leq z. (\varphi(x, y) \rightarrow \varphi(x, y'))) \rightarrow \forall y \leq z \varphi(x, z) \quad (\varphi \in \Delta_0)$$

plus the S -sentence Ω_1 expressing $\forall x \exists y. \omega_1(x) = y$ where $\omega_1(x) := |x|$ and $| \cdot |$ is the length function for the binary representation of x .

We note that by the following result of [Verbrugge,1989]

If $NP \neq CO-NP$ then

$$\not\vdash_{I\Delta_0 + \Omega_1} \forall b, c (\exists a (\text{Prf}(a, c) \wedge \forall z \leq a \neg \text{Prf}(z, b)) \rightarrow \text{Pr}(\ulcorner \exists a \text{Prf}(a, c) \wedge \forall z \leq a \neg \text{Prf}(z, b) \urcorner))$$

it seems highly unlikely that the principle of Σ_1 -completeness, i.e.

$$\varphi \rightarrow \text{Pr}(\ulcorner \varphi \urcorner) \text{ for } \varphi \in \Sigma_1$$

¹ Prerequisites: the reader is supposed to be familiar with [Smoryński,1985]; knowledge of [deJongh-Montagna, 1988] will be helpful.

is provable in $I\Delta_0 + \Omega_1$. However, it can be shown that $I\Delta_0 + \Omega_1$ proves *Švejdar's principle* (Cf. [Švejdar, 1983]): i.e.

$$\vdash_{I\Delta_0 + \Omega_1} \text{Pr}(\ulcorner \varphi \urcorner) \rightarrow \text{Pr}(\ulcorner \exists a (\text{Prf}(a, \ulcorner \psi \urcorner) \wedge \forall z \leq a \neg \text{Prf}(z, \ulcorner \varphi \urcorner)) \rightarrow \psi \urcorner)$$

(for all φ, ψ)

(Cf. [Verbrugge, 1989]) and *Visser's principle* (Cf. [Visser, 1989])

$$\vdash_{I\Delta_0 + \Omega_1} \text{Pr}(\ulcorner C(S) \rightarrow s' \urcorner) \rightarrow \text{Pr}(\ulcorner s' \urcorner)$$

where $C(S) = \bigwedge \{ s \rightarrow \text{Pr}(\ulcorner s \urcorner) : s \in S \}$, S is a finite set of Σ_1 -sentences and s' is a Σ_1 -sentence.

In [Paris-Wilkie, 1987], [Buss, 1986] and [Verbrugge, 1989] ample motivation for the general study of $I\Delta_0 + \Omega_1$ is given; therefore we will turn our attention here directly to the more specific aim of this paper.

In [Parikh, 1971] it is shown that for each primitive recursive function g , there is a Σ_1 -sentence s such that $\vdash_{PA} s$ and

$$g(\mu z. \text{Prf}_{PA}(z, \ulcorner \text{Pr}_{PA}(\ulcorner s \urcorner) \urcorner)) < \mu z. \text{Prf}_{PA}(z, \ulcorner s \urcorner) \quad (*)$$

In [deJongh-Montagna, 1988] Parikh's result is analyzed in the modal context R (Cf. [Guaspari-Solovay, 1979]) when g is the identity function; a simpler proof is presented, based on the fact that $(*)$ has *only provable fixed points*. Furthermore, a characterization is given for pairs of modal formulas $B(p)$ and $C(p)$ such that for each arithmetical interpretation $*$, if $\vdash_{PA} p^* \leftrightarrow (\Box B(p) \prec \Box C(p))^*$ then $\vdash_{PA} p^* : \Box B(p) \prec \Box C(p)$ has only provable fixed points in PA . In [deJongh-Montagna, 1989] the result is extended to arbitrary g which are provably recursive in PA .

Our aim is to refine the positive part of the proof of [deJongh-Montagna, 1988], the part in which it is shown that the formulas specified do indeed have only provable fixed points in PA , to a weaker modal system in which the Σ -completeness axiom (i.e. the

corresponding modal version of the Σ_1 -completeness principle) does not hold.

In section 3, it is shown that the modal version of Visser's principle: i.e.

$$(V) \quad \Box(C(S) \rightarrow s') \rightarrow \Box s'$$

where $C(S) = \bigwedge\{s \rightarrow \Box s : s \in S\}$,
 S is a finite set of Σ -formulas
 s' is a Σ -formula,

playing the role of a weak version of Σ -completeness, suffices to obtain the refined theorem we are looking for.

What is provable in the weak modal system including Visser's schema, is clearly provable in $I\Delta_0 + \Omega_1$ under every arithmetical interpretation; therefore, it follows that PA has no witness comparison formulas having only provable fixed points which the system $I\Delta_0 + \Omega_1$ does not already have.

Based on the result obtained in section 3, in section 4 we present the independence between (V) and the modal version of Švejdar's principle

$$(\check{S}v) \quad \Box A \rightarrow \Box(\Box B \Leftarrow \Box A \rightarrow B) \quad \text{for all formulas } A, B.$$

In particular we give a counterexample to show that $(\check{S}v)$ does not imply Visser's schema: that gives an insight to understand why Švejdar's schema cannot play much of a role in the study of formulas having only provable fixed points.

In an appendix we give some proofs, mainly due to Visser [1989], of modal principles derivable from Visser's principle.

2. MODAL SYSTEMS AND KRIPKE SEMANTICS

In this section we will briefly introduce the modal systems that we are going to work with, together with the associated Kripke-semantics.

Formulas of our system are built up from propositional atoms using the boolean connectives $\wedge, \vee, \neg, \rightarrow, \leftrightarrow, \top, \perp$, a unary modality \Box and binary witness comparisons \prec, \preceq , where \prec and \preceq are applicable only to those formulas having \Box as principal connective. The following definition will introduce the list of modal systems.

DEF 2.1:

(a) B^- (*Basic System*) is the modal system L (Pr1 in [Smoryński, 1985]) (including its rules: modus ponens and necessitation) to which the following *order axioms* are added (see [deJongh, 1987]):

- (01) $\Box A \rightarrow (\Box A \preceq \Box B \vee \Box B \prec \Box A)$
- (02) $\Box A \preceq \Box B \rightarrow \Box A$
- (03) $\Box A \preceq \Box B \wedge \Box B \preceq \Box C \rightarrow \Box A \preceq \Box C$
- (04) $\Box A \prec \Box B \leftrightarrow (\Box A \preceq \Box B \wedge \neg(\Box B \preceq \Box A))$

(b) Z^- (Cf. [Švejdar, 1983]) is the system B^- plus Švejdar's schema:

$$(\text{Šv}) \quad \Box A \rightarrow \Box(\Box B \preceq \Box A \rightarrow B) \quad \text{for all formulas } A, B$$

(c) BV^- is the system B^- plus Visser's schema:

$$(V) \quad \Box(C(S) \rightarrow s') \rightarrow \Box s'$$

where $C(S) = \bigwedge \{ s \rightarrow \Box s : s \in S \}$, S is a finite set of Σ -formulas and s' is a Σ -formula

(d) B, BV, Z are respectively the systems B^-, BV^- and Z^- with the rule $\Box E$ (see Definition 2.1.3 Part A) added.

Let $A(p)$ be some formula of B of the form $\Box B(p) \preceq \Box C(p)$. As in [deJongh-Montagna, 1988] we take BC^-, BVC^- and ZC^- to be the systems B^-, BV^- and Z^- respectively, plus the axiom $c \leftrightarrow A(c)$ (analogous notation is used for the systems B, BV and Z). Since a different system is defined for different choice of A it would be more appropriate to name the systems $BC(A)^-, BVC(A)^-$ and $ZC(A)^-$. But, as it will always be clear in the sequel which formula A is

intended, we will refrain from doing so, in order not to unnecessarily complicate the notation.

DEF 2.2: a *model for B^-* is a finite, tree-ordered Kripke-model for L in which witness comparison formulas are treated as atomic formulas and in which every instance of (O1)-(O4) is forced at each node.

DEF 2.3: *models for BV^- , Z^-* are Kripke-models for B^- where respectively (V) , $(\check{S}v)$ is forced at each node.

It is appropriate to remark that, just as is pointed out in [Verbrugge,1989] for the system Z^- , also for BV^- the forcing for witness comparison formulas in BV^- -Kripke-models is *not persistent*, i.e. it does not necessarily hold that if $j \Vdash \Box A \preceq \Box B$ (resp. $j \Vdash \Box A \prec \Box B$) and jRk then $k \Vdash \Box A \preceq \Box B$ (resp. $k \Vdash \Box A \prec \Box B$).

No model-completeness theorem or even a general extension lemma has been established for BV (for Z , Švejdar did establish these [Švejdar,1983]).

3. WITNESS COMPARISON FORMULAS HAVING ONLY PROVABLE FIXED POINTS IN BV

Theorem 3.3 of [deJongh-Montagna,1988] reads:

If $B(p)$ and $C(p)$ are L -formulas (i.e. do not contain witness comparisons), possibly containing propositional variables other than p , then $A(p) \equiv \Box B(p) \preceq \Box C(p)$ has only provable fixed points in R iff

- (i) $\vdash_L B(T)$
- (ii) $\vdash_L \Box^+(\Box B(\perp) \rightarrow \Box C(\perp)) \rightarrow \Box^{k+1}\perp$, for some k
(where \Box^+D abbreviates $D \wedge \Box D$)

Our aim is to obtain a characterization for a witness comparison formula to have only provable fixed points in BV. The result presented in this section constitutes a refinement of the theorem proved by de Jongh and Montagna; the proof that we present is syntactical and based on a different approach characterized by the proof of the following theorem:

THEOREM 3.1: Let $B(p)$ and $C(p)$ be L-formulas. If

(i) $\vdash_L B(\top)$

(ii) $\vdash_L \Box^+(\Box B(\perp) \rightarrow \Box C(\perp)) \rightarrow \Box^{k+1}\perp$, for some k ,

then $A(p) \equiv \Box B(p) \triangleleft \Box C(p)$ has only provable fixed points in BV.

Some preparatory lemmas are needed. In the following we assume that (i) and (ii) of theorem 3.1 hold, the systems BC^- , BVC^- and BVC refer to the $A(p)$ of this theorem. Some results already proved by Visser (Cf. [Visser,1989]) for his principle and used in the proof of the following lemmas are given in the appendix.

LEMMA 3.2: $\vdash_{BC^-} \Box^+ \neg c \rightarrow \Box^{k+1}\perp$

Pf:

1. $\vdash_{B^-} \Box \neg c \rightarrow \Box(c \leftrightarrow \perp)$

$\rightarrow \Box^+((\Box B(c) \leftrightarrow \Box B(\perp)) \wedge (\Box C(c) \leftrightarrow \Box C(\perp)))$

$\rightarrow (\Box^+(\Box B(c) \rightarrow \Box C(c)) \rightarrow \Box^{k+1}\perp)$

(by (b) and the Substitution Lemma(Cf. [Smoryński, 1985]))

2. $\vdash_{BC^-} \Box^+ \neg c \rightarrow \Box^+(\Box B(c) \rightarrow \Box C(c))$

(by obvious properties of \triangleleft)

3. $\vdash_{BC^-} \Box^+ \neg c \rightarrow \Box^{k+1}\perp$

(by 1 and 2)

☒

LEMMA 3.3: $\vdash_L \Box c \rightarrow \Box B(c)$

Pf:

1. $\vdash_L c \rightarrow B(T)$ (by (i))
2. $\vdash_L \Box c \rightarrow \Box B(T)$
3. $\vdash_L \Box c \rightarrow \Box(c \leftrightarrow T)$
4. $\vdash_L \Box c \rightarrow \Box B(c)$ (by 2 and 3)

☒

LEMMA 3.4: $\vdash_L \Box^+ c \rightarrow \Box^+ B(c)$

Pf:

1. $\vdash_L c \rightarrow B(T)$ (by (i))
2. $\vdash_L \Box^+ c \rightarrow \Box^+ B(T)$
3. $\vdash_L \Box^+ c \rightarrow \Box^+(c \leftrightarrow T)$
 $\quad \rightarrow (\Box^+ B(c) \leftrightarrow \Box^+ B(T))$
4. $\vdash_L \Box^+ c \rightarrow \Box^+ B(c)$ (by 2 and 3)

☒

LEMMA 3.5: $\vdash_L \Box^{k+1} \perp \rightarrow (\Box C(\perp) \rightarrow B(\perp))$

Pf: We claim that, if $\vdash_L \Box^+(\Box B \rightarrow \Box C) \rightarrow \Box^{k+1} \perp$, then $\vdash_L \Box^{k+1} \perp \rightarrow (\Box C \rightarrow B)$, where B, C are arbitrary L -formulas.

For suppose not, then a model M exists such that

$M \models \Box^+(\Box B \rightarrow \Box C) \rightarrow \Box^{k+1} \perp$ and $w \Vdash \Box^{k+1} \perp \wedge \Box C$, $w \not\Vdash B$, for some node w in M . Take the submodel of M generated by w and add a tail of nodes below w of such a length that the new model gets a root x of level greater than or equal to $k+1$ (end nodes are counted as having level 0). Clearly none of the nodes added below w can force $\Box B$ but all of them force $\Box^+(\Box B \rightarrow \Box C)$. By hypothesis, $x \Vdash \Box^{k+1} \perp$ and this gives a contradiction, which proves our claim.

By the claim and (ii) it follows that: $\vdash_L \Box^{k+1} \perp \rightarrow (\Box C(\perp) \rightarrow B(\perp))$. ☒

LEMMA 3.6: $\vdash_{BC^-} \Box^+ \neg c \rightarrow \Box^+ B(c)$

Pf:

1. $\vdash_{\perp} \Box^+ \neg c \rightarrow (\Box C(c) \leftrightarrow \Box C(\perp)) \wedge (B(c) \leftrightarrow B(\perp))$
2. $\vdash_{BC^-} \Box^+ \neg c \rightarrow (\Box C(c) \rightarrow B(c))$ (by lemma 3.2 and lemma 3.5)
3. $\vdash_{BC^-} \neg c \rightarrow (\Box B(c) \rightarrow \Box C(c))$ (by obvious properties of \Leftarrow)
4. $\vdash_{BC^-} \Box^+ \neg c \rightarrow (\Box B(c) \rightarrow B(c))$ (by 2 and 3)
5. $\vdash_{BC^-} \Box^+ \neg c \rightarrow \Box(\Box B(c) \rightarrow B(c))$
 $\rightarrow \Box B(c)$ (by formalized Löb)
6. $\rightarrow B(c)$ (by 4) ☒

LEMMA 3.7:

$\vdash_{BV^-} \Box \Box A \vee \Box \Box B \rightarrow \Box(\Box^+(\Box A \prec \Box B) \vee \Box^+(\Box B \Leftarrow \Box A))$

Pf:

1. $\Box \Box A \vee \Box \Box B \rightarrow \Box(\Box A \prec \Box B \vee \Box B \Leftarrow \Box A)$
 $\rightarrow \Box((\Box A \prec \Box B \rightarrow \Box(\Box A \prec \Box B) \wedge \Box B \Leftarrow \Box A \rightarrow \Box(\Box B \Leftarrow \Box A)) \rightarrow$
 $\rightarrow \Box^+(\Box A \prec \Box B) \vee \Box^+(\Box B \Leftarrow \Box A))$
 $\rightarrow \Box(\Box^+(\Box A \prec \Box B) \vee \Box^+(\Box B \Leftarrow \Box A))$ (by (\vee)) ☒

COROLLARY 3.8: $\vdash_{BV^-} \Box A \vee \Box B \rightarrow \Box(\Box A \prec \Box B \rightarrow \Box(\Box A \prec \Box B))$

Pf: Trivial. ☒

LEMMA 3.9: $\vdash_{BVC^-} \Box^{k+2} c \rightarrow \Box^n \Box B(c)$ for each $0 \leq n \leq k+1$

Pf: by downward induction on n :

$$n = k+1: \vdash_{\perp} \Box^{k+2} c \rightarrow \Box^{k+1} \Box c \\ \rightarrow \Box^{k+1} \Box B(c) \quad \text{by lemma 3.3;}$$

$n < k+1$: recall that by induction hypothesis we have

$$\vdash_{BVC^-} \Box^{k+2} c \rightarrow \Box^{n+1} \Box B(c), \text{ i.e.}$$

$$\vdash_{BVC^-} \Box^{k+2} c \rightarrow \Box^n \Box \Box B(c). \text{ So,}$$

1. $\vdash_{BV^-} \Box^{k+2} c \rightarrow (\Box^{n+1} \Box B(c) \rightarrow$
 $\rightarrow \Box^{n+1}(\Box^+(\Box B(c) \Leftarrow \Box C(c)) \vee \Box^+(\Box C(c) \prec \Box B(c)))$
(by lemma 3.7)

2. $\vdash_{BVC} \Box^{k+2}c \rightarrow (\Box^{n+1}\Box B(c) \rightarrow \Box^{n+1}(\Box^+c \vee \Box^+\neg c))$
 $\rightarrow (\Box^{n+1}\Box B(c) \rightarrow \Box^{n+1}B(c))$ (by lemma 3.4 and lemma 3.6)
 $\rightarrow \Box^{n+1}B(c)$ (by modus ponens with the induction hypothesis)
- ☒

We are now ready to prove theorem 3.1:

Pf (theorem 3.1):

1. $\vdash_{BVC} \Box^{k+2}c \rightarrow \Box B(c)$ (by lemma 3.9 where $n=0$)
 $\rightarrow \Box B(c) \preceq \Box C(c) \vee \Box C(c) \prec \Box B(c)$
 (by obvious properties of \preceq)
2. $\vdash_{BVC} \Box(\Box B(c) \preceq \Box C(c)) \vee \Box(\Box C(c) \prec \Box B(c)) \rightarrow \Box^+c \vee \Box^+\neg c$
 $\rightarrow \Box c \vee \Box^+\neg c$
 $\rightarrow \Box^{k+1}c \vee \Box^{k+1}\perp$
 (by lemma 3.2)
 $\rightarrow \Box^{k+1}c$
3. $\vdash_{BVC} \Box^{k+1}c$ (by 1, 2)
4. $\vdash_{BVC} c$ (by $\Box E$)
- ☒

The refinement that we were looking for is an immediate consequence of theorem 3.1:

THEOREM 3.10: Let $B(p)$ and $C(p)$ be L -formulas; then
 $A(p) \equiv \Box B(p) \preceq \Box C(p)$ has only provable fixed points in BV
 iff

- (i) $\vdash_L B(T)$
 (ii) $\vdash_L \Box^+(\Box B(\perp) \rightarrow \Box C(\perp)) \rightarrow \Box^{k+1}\perp$, for some k .

Pf: (\Rightarrow) If c is a fixed point for $A(p)$ then $\vdash_{BVC} c$, therefore $\vdash_{RC} c$ and by lemma 2.3 in [deJongh-Montagna, 1988]
 $\vdash_{R-} \Box^+(c \leftrightarrow A(c)) \rightarrow \Box^{k+1}c$ for some k . Now apply theorem 3.3 in [deJongh-Montagna, 1988].

(\Leftarrow) by theorem 3.1.

☒

By theorem 3.10 and theorem 3.3 (Cf. [deJongh-Montagna, 1988]) it follows that the formulas of the form $A(p) \equiv \Box B(p) \Leftarrow \Box C(p)$ having only provable fixed points in R are exactly the formulas having only provable fixed points in BV. In other words, to obtain the formulas having only provable fixed points we do not need the strong Σ -completeness schema (i.e. $A \rightarrow \Box A$, for every Σ -formula A) but we can replace it by the weaker (V).

Although theorem 3.10 is formulated with *iff* one should note that, unlike with R and PA, $A(p) \equiv \Box B(p) \Leftarrow \Box C(p)$ having only provable fixed points in $I\Delta_0 + \Omega_1$ for all arithmetical interpretations does not imply that A(p) has only provable fixed points in BV, since arithmetic completeness even of L is unknown for $I\Delta_0 + \Omega_1$ (see [Verbrugge, 1989]). At the present, theorem 3.10 does imply that each formula of R having only provable fixed points in PA has only provable fixed points in $I\Delta_0 + \Omega_1$ when arithmetical interpretations are restricted to sentences. The restriction to sentences is essential; otherwise Visser's principle loses its validity (see [Visser, 1989]).

4. INDEPENDENCE OF VISSER'S AND ŠVEJDAR'S SCHEMAS

As already announced in the introduction, it can be shown that $I\Delta_0 + \Omega_1$ proves Švejdar's principle. Because the principle appears as a weak version of the Σ -completeness axiom it may be of some interest to study its relations with Visser's principle: in this section we will prove the independence of the two principles.

First of all we show that Švejdar's schema does not imply Visser's schema, i.e. $Z \not\vdash (V)$. To prove that, consider the formula $\Box^3 p \Leftarrow \Box^2 p$ having only provable fixed points in R, as proved in [deJongh-Montagna, 1988]. By theorem 3.10 it follows that this formula has only provable fixed points in BV. On the other hand², note that

² the argument was suggested to the author by F.Montagna.

$\Box^3 p \prec \Box^2 p$ cannot have only provable fixed points in Z , because by Švejdar's essential reflexivity interpretation of $\Box A \prec \Box B$ as "there exists a proof of A using axioms with smaller Gödel numbers than in any proof of B " (Cf. [Švejdar, 1983]) that would mean that for the fixed point c in PA , $\Box^2 c$ would have a proof in PA using axioms with smaller Gödel numbers than any proof of $\Box c$ would use. This is impossible because being a provable Σ -sentence, $\Box c$ wouldn't need any but the axioms of Q and we could take those as the zero base of our interpretation. This proves our claim.


At this point it may be of interest to remark that the formula $\Box^2 p \prec \Box p$ has only provable fixed points in Z .

The following argument is due to Visser: in BC^- it is provable that $\Box^2 c \rightarrow \Box(\Box c \prec \Box^2 c \vee \Box^2 c \prec \Box c)$. Thus, in ZC^- , $\Box^2 c \rightarrow \Box c$ is provable, from which with Löb in ZC , immediately c follows. Under the same arithmetical interpretation used in the previous argument, the result is not very surprising: it is well known that there are theorems provable in PA and not in Q . From these observations we can see that Švejdar's schema can by itself hardly be useful in studying formulas having only provable fixed points in BV . Recall also that in the proof of theorem 3.10, the schema ($\check{S}v$) is not used.

To obtain our second claim, that Visser's schema does not imply Švejdar's schema (i.e. $BV \not\vdash (\check{S}v)$), it is enough to exhibit a countermodel of BV to the formula $\Box p \rightarrow \Box(\Box q \prec \Box p \rightarrow q)$ (i.e. an instance of ($\check{S}v$) where p and q are propositional variables)³.

Let A be the formula $\Box p \rightarrow \Box(\Box q \prec \Box p \rightarrow q)$ and consider the following A -sound model $\langle \{1, 2, 3, \dots\}, R, \Vdash \rangle$ where the forcing relation is restricted to subformulas of A :

³ Observe that $\vdash_{BV} \Box A \rightarrow \Box \Box(\Box B \prec \Box A \rightarrow B)$, for all formulas A, B . The proof is an immediate consequence of lemma 3.7.



where E and F stand for $\Box p \prec \Box q$ and $\Box q \prec \Box p$ respectively.

From the forcing relation indicated in the figure note that: 2 does not force p or q; 4 does not force p or $E \rightarrow \Box E$, but does force $F \rightarrow \Box F$; for kR5 and $k=5$, k does not force p, q, E, F, but does force $E \rightarrow \Box E$ and $F \rightarrow \Box F$. In particular note that 4 does not satisfy $\Box p \rightarrow \Box(\Box q \prec \Box p \rightarrow q)$.

Observe that the role of node 1 is crucial to obtain a model forcing all the instances of Visser's principle; consider the formula $\Box \neg p$ and suppose that node 1 did not exist. It is easy to check that $6 \not\models \Box((E \rightarrow \Box E) \rightarrow \Box \neg p) \rightarrow \Box \Box \neg p$.

We claim that under a suitable forcing extension given to the model, every instantiation of Visser's principle holds on the model. Before

giving the procedure to define the appropriate forcing relation, let us fix some notation and definition which will be used in the sequel. We write P to denote the set of all propositional variables except p and q ; S^0 for $\{p, q, \Box p, \Box q, \Box p \leq \Box q, \Box q < \Box p\}$; S^{2m+1} to denote the closure of $S^{2m} \cup P$ under the propositional connectives and \Box (obviously P is effective only when $m=0$); S^{2m+2} for $S^{2m+1} \cup \{\Box A \leq \Box B, \Box A < \Box B \mid \Box A, \Box B \in S^{2m+1}\}$.

DEF 4.1: let k, k' nodes of $\langle \{1,2,3,\dots\}, R, \Vdash \rangle$; we write:
 $\Box A <_k \Box B$ iff $\exists k' ((k'Rk \text{ or } k'=k) \text{ and } k' \Vdash \Box A \text{ and } k' \nVdash \Box B)$
 $\Box A \leq_k \Box B$ iff $k \Vdash \Box A$ and $\forall k' ((k'Rk \text{ or } k'=k) \text{ and } k' \Vdash \Box B \text{ then } k' \Vdash \Box A)$

Here is the procedure to construct the forcing relation:

stage 0: for all $r \in P$ fix

$k \Vdash r$ iff $k \Vdash p$ for all nodes k

stage $2m+1$: automatically and uniquely define a forcing relation for all members of the closure S^{2m+1} ;

stage $2m+2$: call (as in [deJongh, 1987]) a boxed formula $\Box A$ *old* if $\Box A \in S^{2m}$ and *new* if $\Box A \in S^{2m+1} \setminus S^{2m}$. To give an extension of the forcing relation to S^{2m+2} , it is enough to define the forcing on witness comparison formulas $\Box A \leq \Box B$ and $\Box A < \Box B$ (belonging to S^{2m+2}) for $\Box A, \Box B$ both new, $\Box A$ old and $\Box B$ new, and for $\Box A$ new and $\Box B$ old. Before giving the way to construct the forcing let us recall two definitions occurring in [deJongh, 1987]:

- (i) $k \Vdash \Box A < \Box B$ iff $\Box A <_k \Box B$ or, $\Box A \leq_k \Box B$ and $\Box A$ old, $\Box B$ new
- (ii) $k \Vdash \Box A \leq \Box B$ iff $\Box A <_k \Box B$ or, $\Box A \leq_k \Box B$ and $\Box B$ new.

We are now ready to present the procedure, to repeat for all nodes k . Here it is:

If $k \in \{1, 2, 3, 4, 5\}$ and $k \Vdash \Box A$ and $k \nVdash \Box B$

then let $k \Vdash \Box A < \Box B$ and $k \Vdash \Box A \leq \Box B$

else fix the forcing on $\Box A < \Box B, \Box B < \Box A, \Box A \leq \Box B$ and $\Box B \leq \Box A$ as defined in (i) and (ii), respectively.

Apply the procedure repeatedly (i.e for all $m \in \mathbb{N}$) so as to cover all formulas, and call the resulting model M .

Note that points 1 and 3 satisfy the same formulas since they are always treated alike by the construction.

CLAIM 1: $\forall s \in \Sigma. 5 \Vdash s \Rightarrow k \Vdash s$ where $k \in \{1, 2, 4\}$

Pf: suppose $s \in \Sigma$ and $5 \Vdash s$; by cases:

$s \equiv \Box B$: by the previous observation points 1 and 3 force the same formulas, therefore the claim;

$s \equiv \Box B \prec \Box C$: by stage $2m+2$ of construction and definition (i);

$s \equiv \Box B \preceq \Box C$: by definition (ii) on stage $2m+2$ of construction;

$s \equiv$ "boolean combination of Σ -formulas": by the previous cases. \square

Using claim 1 and definition 4.1, it is easy to check that stage $2m+2$ excludes the existence of two boxed formulas $\Box A, \Box B$ for which $\Box A \prec \Box B$ and $\Box B \preceq \Box A$ are both forced at node 4.

CLAIM 2: $5 \Vdash C(S)$ for all finite sets S of Σ -formulas

Pf: straightforward from claim 1. \square

CLAIM 3: (*Persistency property*) let $\Box A, \Box B$ be two boxed formulas such that at least one of them is new at some stage $m \geq 1$;

if $(k \Vdash \Box A \prec \Box B$ and $kRk')$ then $k' \Vdash \Box A \prec \Box B$ and

if $(k \Vdash \Box A \preceq \Box B$ and $kRk')$ then $k' \Vdash \Box A \preceq \Box B$

Pf: immediate from the forcing procedure and the following consequences of definition 4.1:

if $(\Box A \prec_k \Box B$ and $kRk')$ then $\Box A \prec_{k'} \Box B$ and

if $(\Box A \preceq_k \Box B$ and $kRk')$ then $\Box A \preceq_{k'} \Box B$. \square

Note that the only witness comparison formulas that do not satisfy the persistency property are E and F (see definition of forcing at nodes 3 and 4).

CLAIM 4: all instances of Visser's schema are forced in each node of M .

Pf: obviously points 1, 2, 3 satisfy the claim; moreover notice that Visser's principle is always satisfied at level 1 in any Kripke model since each $C(S)$ is always satisfied at terminal nodes, therefore 4 and 5 satisfy the claim. By induction we check the tail of points k :

$k=6$: suppose there exist $C(S)$ and s' such that $6 \Vdash \Box s'$ and $\forall k$ (if $6Rk$ then $k \Vdash C(S) \rightarrow s'$); it follows that $\exists h$ ($6Rh$ and $h \not\Vdash s'$ and $h \Vdash C(S)$); but $k \Vdash C(S)$ for $k \in \{1, 2, 3, 5\}$ therefore h must be 4. By claim 1 we get a contradiction.

$k+1$: (with $k+1 > 6$) assume the claim holding for all h such that $k+1Rh$ and suppose there exist $C(S)$ and s' such that $k+1 \Vdash \Box s'$ and $\forall h$ (if $k+1Rh$ then $h \Vdash C(S) \rightarrow s'$); it follows that $\exists h$ ($k+1Rh$ and $h \not\Vdash s'$ and $h \Vdash C(S)$); this node must be k since, by induction hypothesis, every instance of Visser's schema holds at k , so $k \Vdash \Box s'$. Therefore $k \not\Vdash C(S)$, i.e. for some $s \in S$, $k \Vdash s$ but $k \not\Vdash \Box s$. By cases:

$s \equiv \Box B$: $k \Vdash \Box s$, a contradiction;

$s \equiv \Box B \prec \Box C$: $k \Vdash \Box B$ and B can be neither p nor q since $\Box p$ and $\Box q$ are not forced at any point $kR6$. Therefore by claim 3, the forcing on witness comparison formulas must be persistent and this gives a contradiction.

$s \equiv \Box B \preceq \Box C$: similar to the previous case;

$s \equiv$ "boolean combination of Σ -formulas": by the previous cases.

☒

To show that M is a model for BV it suffices to prove the following

CLAIM 5: for all formulas A ,
if $M \vDash \Box A$ then $M \vDash A$,

Pf: trivial.

☒

APPENDIX: SOME THEOREMS PROVED BY VISSER'S PRINCIPLE

In [Visser,1989] the following theorems, proved using the principle (V), are pointed out:

- (V1) $\Box \bigvee s \rightarrow \Box \bigvee s^+$
(V2) $\Box(\Box A \rightarrow \bigvee s) \wedge \Box(\bigvee s^+ \rightarrow A) \rightarrow \Box A$
(V3) $\Box(C(S) \rightarrow (A \rightarrow s')) \rightarrow \Box A \rightarrow \Box s'$
(V4) $\Box(C(S) \rightarrow (\Box s' \rightarrow s')) \rightarrow \Box s'$

where S is a finite set of Σ -formulas, $C(S) = \bigwedge \{ s \rightarrow \Box s : s \in S \}$, $S^+ = \{ s \wedge \Box s : s \in S \}$ and s' a Σ -formula.

We will give the proof of them in the modal system BV^- :

(V1):

1. $\Box \bigvee s \rightarrow \Box(C(S) \rightarrow \bigvee \Box^+ s)$
2. $\Box(C(S) \rightarrow \bigvee \Box^+ s) \rightarrow \Box(\bigvee \Box^+ s)$ (by (V))
3. $\Box \bigvee s \rightarrow \Box(\bigvee \Box^+ s)$ (by 1 and 2)

(V2):

1. $\Box(\Box A \rightarrow \bigvee s) \rightarrow \Box(\Box \Box A \rightarrow \Box \bigvee s)$
 $\rightarrow \Box(\Box \Box A \rightarrow \Box(\bigvee \Box^+ s))$ (by (V1))
2. $\Box(\bigvee \Box^+ s \rightarrow A) \rightarrow \Box(\Box \bigvee \Box^+ s \rightarrow \Box A)$
3. $\Box(\Box A \rightarrow \bigvee s) \wedge \Box(\bigvee \Box^+ s \rightarrow A) \rightarrow \Box(\Box \Box A \rightarrow \Box A)$ (by 1 and 2)
 $\rightarrow \Box \Box A$ (by formalized Löb)
 $\rightarrow \Box s$
 $\rightarrow \Box(\bigvee \Box^+ s)$ (by (V1))
 $\rightarrow \Box A$

(V3):

1. $\Box(C(S) \rightarrow (A \rightarrow s')) \rightarrow \Box(A \rightarrow (C(S) \rightarrow s'))$
 $\rightarrow \Box A \rightarrow \Box(C(S) \rightarrow s')$
 $\rightarrow \Box A \rightarrow \Box s'$ (by (V))

(V4):

1. $\Box(C(S) \rightarrow (\Box s' \rightarrow s')) \rightarrow \Box(\Box(C(S) \rightarrow (\Box s' \rightarrow s')))$
 $\rightarrow \Box(\Box \Box s' \rightarrow \Box s')$ (by (V3))
 $\rightarrow \Box \Box s'$ (by formalized Löb)
 $\rightarrow \Box(C(S) \rightarrow \Box s')$
 $\rightarrow \Box(C(S) \rightarrow s')$
 $\rightarrow \Box s'$ (by (V))

REFERENCES

1. Buss, S., 1986. *Bounded arithmetic*, Bibliopolis, Napoli.
2. de Jongh, D.H.J., 1987. "A simplification of a completeness proof of Guaspari and Solovay", *Studia Logica*, vol.46, 187-192.
3. de Jongh, D.H.J., Montagna, F., 1989. "Much shorter proofs", *Zeitschr. für Math. Log. und Grundlagen d.Math*, vol.35, 247-260.
4. de Jongh, D.H.J., Montagna, F., 1988. "Provable fixed points", *Zeitschr. für Math. Log. und Grundlagen d.Math*, vol.34, 229-250.
5. Guaspari, D., Solovay, R., 1979. "Rosser sentences", *Annals of Math. Logic*, vol.16, 81-99.
6. Parikh, R.J., 1971. "Existence and feasibility in arithmetic", *Journal of Symbolic Logic*, vol.36, 494-508.
7. Paris, J., Wilkie, A., 1987. "On the scheme of induction for bounded arithmetic formulas", *Annals of Pure and Applied Logic*, vol.35, 261-302.

8. Smoryński, C., 1985. *Self-reference and modal logic*, Springer-Verlag, Berlin.
9. Švejdar, V., 1983, Modal analysis of generalized Rosser sentences, *Journal of Symbolic Logic*, vol.48, 986–999.
10. Verbrugge, L.C., 1989. "Σ-completeness and bounded arithmetic", *ITLI Prepublication Series*, ML-89-05, Department of Mathematics and Computer Science, University of Amsterdam.
11. Visser, A., 1989. "On the Σ_1 -conservativity of Σ_1 -completeness", *Logic Group Preprint Series*, no. 52, Department of Philosophy, University of Utrecht.

ACKNOWLEDGEMENTS

The present result have been found during my visit to (the Department of Mathematics and Computer Science of the University of) Amsterdam and in direct interaction with the research by Albert Visser on Σ_1 -conservativity of Σ_1 -completeness of $I\Delta_0 + \Omega_1$. My special thanks are given to Dick de Jongh for his nice way of supervising me during my presence in Amsterdam and for many helpful remarks that he made during the research and the drafting of the paper. I am grateful to Rineke Verbrugge and Albert Visser for a number of corrections and suggestions. Finally, I would like to thank Franco Montagna for the stimulating discussions on provable fixed points.

The ITLI Prepublication Series

1990

Logic, Semantics and Philosophy of Language

LP-90-01 Jaap van der Does
LP-90-02 Jeroen Groenendijk, Martin Stokhof
LP-90-03 Renate Bartsch
LP-90-04 Aarne Ranta
LP-90-05 Patrick Blackburn
LP-90-06 Gennaro Chierchia
LP-90-07 Gennaro Chierchia
LP-90-08 Herman Hendriks
LP-90-09 Paul Dekker

LP-90-10 Theo M.V. Janssen

Mathematical Logic and Foundations

ML-90-01 Harold Schellinx
ML-90-02 Jaap van Oosten
ML-90-03 Yde Venema
ML-90-04 Maarten de Rijke
ML-90-05 Domenico Zambella

Computation and Complexity Theory

CT-90-01 John Tromp, Peter van Emde Boas
CT-90-02 Sieger van Denneheuvel

Gerard R. Renardel de Lavalette

CT-90-03 Ricard Gavaldà, Leen Torenvliet
Osamu Watanabe, José L. Balcázar

CT-90-04 Harry Buhrman, Leen Torenvliet

Other Prepublications

X-90-01 A.S. Troelstra

X-90-02 Maarten de Rijke

X-90-03 L.D. Beklemishev

X-90-04

X-90-05 Valentin Shehtman

X-90-06 Valentin Goranko, Solomon Passy

X-90-07 V. Yu. Shavrukov

X-90-08 L.D. Beklemishev

X-90-09 V. Yu. Shavrukov

X-90-10 Sieger van Denneheuvel

Peter van Emde Boas

X-90-11 Alessandra Carbone

A Generalized Quantifier Logic for Naked Infinitives
Dynamic Montague Grammar
Concept Formation and Concept Composition
Intuitionistic Categorical Grammar
Nominal Tense Logic
The Variability of Impersonal Subjects
Anaphora and Dynamic Logic
Flexible Montague Grammar
The Scope of Negation in Discourse,
towards a flexible dynamic Montague grammar
Models for Discourse Markers

Isomorphisms and Non-Isomorphisms of Graph Models
A Semantical Proof of De Jongh's Theorem
Relational Games
Unary Interpretability Logic
Sequences with Simple Initial Segments

Associative Storage Modification Machines
A Normal Form for PCSJ Expressions

Generalized Kolmogorov Complexity
in Relativized Separations
Bounded Reductions

Remarks on Intuitionism and the Philosophy of Mathematics,
Revised Version

Some Chapters on Interpretability Logic

On the Complexity of Arithmetical Interpretations of Modal Formulae
Annual Report 1989

Derived Sets in Euclidean Spaces and Modal Logic

Using the Universal Modality: Gains and Questions

The Lindenbaum Fixed Point Algebra is Undecidable

Provability Logics for Natural Turing Progressions of Arithmetical
Theories

On Rosser's Provability Predicate

An Overview of the Rule Language RL/1

Provable Fixed points in $\mathcal{I}\Delta_0 + \Omega_1$, revised version